

Accepted Manuscript

Use of canonical variate analysis to differentiate onion cultivars by mineral content as measured by ICP-AES

Gemma A. Chope, Leon A. Terry

PII: S0308-8146(09)00013-2
DOI: [10.1016/j.foodchem.2008.12.090](https://doi.org/10.1016/j.foodchem.2008.12.090)
Reference: FOCH 8173

To appear in: *Food Chemistry*

Received Date: 4 June 2008
Revised Date: 4 December 2008
Accepted Date: 29 December 2008



Please cite this article as: Chope, G.A., Terry, L.A., Use of canonical variate analysis to differentiate onion cultivars by mineral content as measured by ICP-AES, *Food Chemistry* (2009), doi: [10.1016/j.foodchem.2008.12.090](https://doi.org/10.1016/j.foodchem.2008.12.090)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Use of canonical variate analysis to differentiate onion cultivars by mineral content**
2 **as measured by ICP-AES.**

3
4 Gemma A. Chope and Leon A. Terry*

5 Plant Science Laboratory, Cranfield University, Bedfordshire, MK43 0AL, UK.

6 * Corresponding author. Tel.: +44-7500-766-490

7 *E-mail address:* l.a.terry@cranfield.ac.uk (L.A. Terry).

8
9 **Abstract**

10 Three onion cultivars *viz.* Renate, Ailsa Craig and SS1 were characterised according to
11 their mineral content. The concentrations of the macronutrients phosphorus, potassium,
12 calcium, manganese and sulphur and the micronutrients iron, boron, manganese, copper
13 and zinc were analysed in freshly harvested and stored onion bulbs using ICP-AES
14 (Inductively coupled plasma – atomic emission spectroscopy). Onions were treated pre-
15 harvest with additional sulphur (100 kg ha⁻¹) and/or calcium (300 kg ha⁻¹) applied in four
16 combinations at the time of seed drilling, however these treatments did not affect the total
17 concentrations of sulphur or calcium in the harvested bulbs. The data were subjected to
18 canonical variate analysis in order to determine the most appropriate variate to
19 discriminate between cultivars. Two canonical variates were sufficient to differentiate
20 between the three cultivars, with the first canonical variate describing differences in
21 micronutrients between the genotypes and the second separating the cultivars by
22 differences in sulphur concentration.

Keywords: *Allium cepa*; calcium; multivariate analysis; sulphur.

1. Introduction

The elemental composition of the edible portion of some *Allium* species has been used as a method for defining geographic origin (Ariyama, Horita, & Yasui, 2004; Ariyama & Yasui, 2006; Ariyama, Aoyama, Mochizuki, Homura, Kadokura, & Yasui, 2007). In addition, analysis of 63 major and trace elements was shown to allow differentiation between conventionally and organically grown onions cv. Hysam from Denmark (Gundersen, Bechmann, Behrens, & Stürup, 2000). However, apart from these works and others (Bibak *et al.*, 1998; Alvarez, Marcó, Arroyo, Greaves, & Rivas, 2003; (Rodríguez Galdon, Oropeza González, Rodríguez Rodríguez, & Díaz Romero, 2008) there has been little research concerning the elemental composition of bulb onions. Most work concerning manipulation of mineral nutrition of onions has concentrated on the effect on quality parameters, particularly the effects on pungency (enzymatically produced pyruvate after maceration), flavour precursors and firmness. Only a few studies have performed analysis to determine whether the changes in mineral nutrition resulted in an effect on total mineral concentrations in the bulb. Field, hydroponic and tissue culture trials worldwide have shown that attempting to manipulate the sulphur content of onions by varying the sulphur supply during growth has had mixed results and is highly dependent on a range of factors. These factors include cultivar, the extent of variation of sulphur supply, and other seasonal and environmental influences including climatic conditions and soil type (Randle, 1992; Randle, Bussard, & Warnock, 1993; Hamilton,

Yoo, & Pike, 1998; Kopsell, Randle, & Eiteman, 1999; Lancaster, Farrant, & Shaw, 2001; Coolong, Kopsell, Kopsell, & Randle, 2004; O'Donoghue *et al.*, 2004). In general, increasing the sulphur supply increases firmness, pungency, alk(en)yl cysteine sulfoxides (ACSOs) and dry matter when the treatment is applied constantly (e.g. in a hydroponic system), or when applied as a field treatment towards the end of the growing season when bulbing has been initiated. The effect of the addition of calcium chloride to the crop has also been investigated in the USA (Coolong & Randle, 2008). Additional calcium nutrition resulted in increased firmness of bulbs at harvest, although this effect did not persist throughout storage. Reduced pyruvate concentration was observed in CaCl_2 -treated onions, but this effect only occurred in one out of two growing seasons. Work carried out on UK-grown onions was unable to replicate these results (B. Smith & T. Crowther, pers. comm, 2005).

Given previous findings, the aim of the research presented herein was to determine the effects of field application of additional calcium and/or sulphur treatments on the mineral content of onion bulbs grown from seed, and to describe genotypic differences in the mineral content of onion bulbs of short, intermediate and long-storing onion cultivars *viz.* SS1, Ailsa Craig and Renate respectively using canonical variate analysis.

2. Experimental

2.1 Plant material and storage regime

Short, intermediate and long-storing onion (*Allium cepa* L.) cvs.; SS1, Ailsa Craig and Renate, respectively, were spring-drilled on 18 March 2003 in four rows per bed using a tape seeder at a rate of 35 seeds m⁻² and grown in a sandy loam soil field at FB Parrish & Son (Beds., UK) using standard agronomic practices. Two treatments of additional sulphur and/or calcium at rates of 100 kg ha⁻¹ of sulphur and 300 kg ha⁻¹ of calcium were applied in four combinations including a negative control (plus Ca minus S, plus Ca and S, minus Ca plus S and minus Ca and S). Sulphur, in the form of agricultural gypsum, was applied uniformly over the plot area at the time of drilling. Calcium was applied evenly by hand in the form of 77% calcium chloride flakes (Kemira, Cheshire, UK). Onion bulbs were harvested (cv. SS1 on 19 August 2003; cvs. Renate and Ailsa Craig on 2 September 2003) into standard 25 kg plastic nets and dried in bin driers with ambient air for five weeks (cv. SS1) and three weeks (cvs. Ailsa Craig and Renate) as per standard practice in the UK. The dry aerial parts and roots were removed, and any diseased or damaged bulbs discarded prior to storage. Bulbs were held under industry standard controlled atmosphere (CA) conditions (3 kPa CO₂ and 5 kPa O₂; Smittle, 1988) using an Oxystat 2 CA system, attached to an Oxystat 2002 Controller, and Type 770 fruit store analyser (David Bishop Instruments, Sussex, UK). This system was self-calibrating every 24 h against 5% CO₂ in N₂ (British Oxygen Co., Surrey, UK) as previously described by Chope, Terry & White (2007). Bulbs were stored at 2 ± 1°C inside two rigid polypropylene fumigation chambers (88 x 59 x 59 cm). Relative humidity was not measured.

91

92 *2.2 Experimental design*

93

94 The experiment was conducted as a completely randomised design with the
95 assumption that the storage containers were identical and the samples were taken
96 randomly. Bulbs were divided equally between the two storage containers. Bulbs were
97 removed from storage at regular intervals. Samples of cv. Renate were taken after days
98 0, 40 and 230, cv. Ailsa Craig after days 0, 81 and 129, and cv. SS1 after days 0, 40 and
99 81. At each sampling date (except time 0) for each cultivar, three bulbs from each
100 treatment were sampled from each storage chamber. At sampling time 0, four sets of six
101 bulbs were sampled. Samples from each set of three bulbs from each sampling time and
102 treatment were combined and lyophilised prior to mineral analysis (except at time 0
103 where sets of 6 bulbs were combined); therefore, values represent the mean of the
104 replicates for each sampling date and cultivar.

105

106 *2.3 Sample preparation*

107

108 The dry outer skins were removed, and then a vertical wedge of tissue was taken from
109 the basal section of the bulb. Each sample was weighed and immediately snap-frozen in
110 liquid nitrogen, and stored at -40°C until it was lyophilised. Dry weight measurements
111 were made on lyophilised samples.

112

113 *2.4 Mineral analysis*

Lyophilised bulb tissue ($n = 60$) was ashed and then digested with 1 ml concentrated nitric acid (ADAS, 1985). The intensity of ion response was measured by ICP-AES (Inductively coupled plasma – atomic emission spectroscopy; Ultima 2, Jobin Yvon, London, UK). Results were expressed as total ion concentrations of boron, calcium, copper, iron, magnesium, manganese, phosphorus, potassium, sodium, sulphur and zinc by comparison with external standards.

2.5 Statistical analysis

The effect of the application of additional calcium and sulphur to onion plants in the field on the concentration of each mineral analysed was assessed using ANOVA. The calcium and sulphur treatments did not affect total bulb sulphur or calcium content at a probability level of 0.05. Therefore, the data sets for all calcium and sulphur treatments were pooled. A main effect of cultivar was identified by univariate analysis of variance (ANOVA). The groups were further defined using canonical variate analysis (CVA) (Gardner, Gower & Le Roux, 2006), which bases the analysis on grouping the data so that the variance between groups (cultivars) is maximized and the variance within groups is minimised. The circular 95% confidence limits represent the confidence interval for the population rather than the population mean and have radii of 2.448. The basic radius value is the square root of the chi-square distribution value at the 5% significance level, for a distribution with 2 degrees of freedom (as displaying the CVA in 2 dimensions).

All statistical analyses were carried out using Genstat for Windows Version 7.1.0.198 (VSN International Ltd., Herts., UK).

3. Results and discussion

The variation in mineral content between cultivars was greater than the variation caused by storage time (Fig. 1). Bulbs of each cultivar were stored for different lengths of time due to differences in storage life; however, the sampling points represent the beginning, middle and end of the storage period for each.

INSERT FIGURE 1.

To enable comparison with other published data on the elemental composition of onions, the data was calculated on a fresh weight basis using the mean dry weights of each cultivar (Table 1).

INSERT TABLE 1.

Free mineral content should not be expected to change on a dry weight basis during storage yet mineral content will change as a proportion of fresh weight due to weight loss. Therefore, the different concentrations recorded at various storage durations are more likely to represent the natural variation within the population rather than real changes in mineral concentration. Onion cv. Ailsa Craig bulbs showed the greatest

variability over time (Fig. 1), and this may be attributed to the fact that it is a not a commercial cultivar and maintenance of the breeding line is poor (B. Smith, pers. comm, 2004). Previously published data (Hansen, Wyse, & Sorensen, 1979; Holland, Unwin, & Buss, 1991; U.S. Department of Agriculture, 2004) is comparable with that recorded in this investigation (Table 1); however, onion cv. Renate bulbs contained approximately 2-fold the concentration of calcium, copper, iron, potassium and sodium than the values cited by the literature (U.S. Department of Agriculture, 2004) for unspecified cultivars. The data are, however, within the same range as that reported for calcium, copper, iron, manganese, potassium and zinc in cvs. 438 Granex and Yellow Granex, using a technique based on total reflection X-ray fluorescence and ultrasound-based extraction procedure (Alvarez *et al.*, 2003), and for those reported for boron, copper, manganese, potassium and sodium in onion cv. Hysam measured using high resolution ICP-MS (Bibak *et al.*, 1998). In addition, the data reported here are similar to those reported for copper, iron, manganese and zinc in the fresh leaves of other *Allium* L. species (*viz.* *A. schoenoprasum* L., *A. nutans* L., *A. angulosum* L., *A. fistulosum* L., *A. montanum* Schmidt, *A. odorum* L. and *A. flavescens* Bess.) measured using ICP-MS (Golubev, Golubkina, & Gorbunov, 2003). The reason for differences occurring between reported values for mineral concentrations in onion is likely to be due to a combination of factors including the preharvest growing conditions, postharvest treatment, analytical methods, and, as has been demonstrated here, genotype.

Onion cv. SS1 bulbs had a greater concentration of total sulphur per gram dry weight than cv. Ailsa Craig and Renate (Fig. 1). This is perhaps surprising as cv. SS1 was the least pungent of the three cultivars (Chope, Terry, & White, 2006), and pungency

is determined by the availability of sulphur-containing flavour precursors (ACSOs). Pungency is calculated on a fresh weight basis to reflect the edible product. However, all mineral analyses were performed on lyophilised tissue, and when the concentrations were calculated on a fresh weight basis (using the mean dry weight percentage), free sulphur concentration was least in cv. SS1 bulbs (Table 1). In addition, it has been shown that there are genotypic differences in the partitioning of sulphur accumulated by the onion plant. Randle, Kopsell, Kopsell, & Snyder (1999), showed that total bulb sulphur accumulation was poorly correlated with pungency in three onion cultivars. They concluded that a key contributing factor in determination of the pungency of an onion depends upon the capability of the plant to partition sulphur as sulphate in the vacuole, thus reducing the amount of sulphur incorporate into the ACSO biosynthetic pathway.

All elements measured (except for calcium) were present in significantly different ($P<0.005$) concentrations in onion cvs. Renate, Ailsa Craig and SS1 bulbs (Fig. 1). There was no consistent pattern according to which cultivar had the highest and lowest concentration of each element, although cv. Ailsa Craig did not have the highest concentration of any of the elements measured. However, since ANOVA is a univariate technique which allows comparisons to be made between cultivars based on a single variate it does not give an insight into how the cultivars are grouped or which is the most important element in defining groups. Canonical variate analysis (CVA) allows these types of conclusions to be drawn, and has previously been used to discriminate between cultivars of other crop species (Cole & Phelps, 1979). Another type of multivariate analysis, principal component analysis (PCA) has been recently applied to data on mineral and trace elements in onion bulbs of six different cultivars from various seed

origins cultivated under the same agronomic, climatic and soil conditions (Rodríguez Galdon *et al.*, 2008). Significant differences in mineral composition were reported between cultivars, although relatively high genotypic overlap was evident.

Good separation of the cultivars was obtained using CVA. The majority of the variation, 89.23 %, was explained by the first canonical variate (CV1), which separated Renate from cvs. Ailsa Craig and SS1, while 10.77 % of the variation was explained by the second canonical variate. Onion cv. Renate bulbs differed from cvs. Ailsa Craig and SS1, as shown by the high positive score for Renate in CV1 (5.901), versus negative scores for Ailsa Craig (-2.702) and SS1 (-2.134), respectively. Canonical variate 2 (CV2) discriminated between cv. SS1 and cvs. Ailsa Craig and Renate, as shown by the positive score for SS1 for CV2 (1.536), versus negative scores for Ailsa Craig (-1.424) and Renate (-0.118), respectively. In the case of CV1 there was less variation between the cultivars in terms of the macronutrients measured (*viz.* phosphorus, potassium, calcium, manganese and sulphur) than for the micronutrients measured (*viz.* iron, boron, manganese, copper and zinc), as demonstrated by the higher scores for CV1 for the macronutrients compared to the micronutrients (Table 2).

INSERT TABLE 2.

A plot of the means of the data relative to the first two canonical variates (Fig. 2) showed that there was minimal overlapping between the 95% confidence limits for each cultivar. The highest loading for CV1 was assigned to boron, and onions cv. Renate had a high positive value for CV1, signifying a high boron concentration compared to cvs.

Ailsa Craig and SS1. The highest loading for CV2 was assigned to sulphur, indicating that onion bulbs cv. SS1 contained a higher concentration of sulphur per dry weight than cvs. Renate and Ailsa Craig, which was also demonstrated using ANOVA.

INSERT FIGURE 2

The sulphur and calcium treatments, applied at the time of seed drilling, did not affect bulb sulphur or calcium content. Similarly, fertilisation with calcium (in the form of calcium carbonate) or magnesium (in the form of magnesium sulphate) a few days prior to planting did not affect the concentration of magnesium or calcium in onion cv. Super-Kitamomjii bulbs grown in Japan (Ariyama, Nishada, Noda, Kadokura, & Yasui, 2006). Conversely, Coolong & Randle (2008) recently reported that application of calcium chloride to low pungency Vidalia cv. Georgia Boy onions increased bulb calcium concentration in each of two growing seasons. The final rate of application was 115 and 230 kg ha⁻¹, applied at 8, 12, 16 and 20 weeks after transplant, ensuring a regular supply throughout the growing season. It may be that this staggered approach to application was the reason why mineral content was affected. It is also possible that the method of cultivation i.e. the use of transplants rather than direct drilling could have affected the results. However, sulphur concentration was only increased in one growing season following application of 155 or 230 kg ha⁻¹ ammonium sulphate staggered at 6 and 10 weeks after transplant. It is evident from the results presented herein that the sulphur and calcium applied was not present in the soil in a sufficiently available form, or was applied too early to the crop. There is also the possibility that the sulphur and calcium were taken

up early on in the growth period, and retained in the upper part of the plant, however, it has been shown that ACSOs are translocated from the leaves to the storage scales during bulbing (Mallor & Thomas, 2008). In addition, it may have been that the soil conditions supplied sufficient sulphur and calcium nutrition to plants so that the applied treatments did not have a measurable effect.

4. Conclusion

The work presented herein has shown that it is possible to separate the three cultivars in this experiment on the basis of mineral content using just two canonical variates. It was also apparent that when onions of different cultivars are grown on the same site (i.e. same soil and climatic conditions), there are still significant differences in the mineral composition of the bulbs. It is therefore likely that the source of this variation is genotypic. Additional calcium and sulphur fertilisation applied at the time of drilling had no significant effect on the concentrations of these minerals in the bulbs. Therefore, it is important to consider the mode and timing of application of these treatments, and it should not be assumed that mere application to the soil results in greater bioavailability.

Acknowledgements

This work was performed as part of a project funded by the Horticultural Development Council; 'Understanding the mechanisms behind onion bulb dormancy in relation to the potential of improved onion storage' (CP20). Bulb material was supplied

from HortLink project HL0164LFV (Defining quality assurance for sweet onions with rapid biosensor analysis) funded by the UK Government (Department for Environment, Food and Rural Affairs; Defra) and UK industry representatives. Thanks are given to both Brian Smith and David O'Connor. Mineral analysis was carried out at Warwick HRI (Warks, UK).

References

- ADAS. (1985). *The analysis of agricultural materials, MAFF Reference book 427*. HMSO, London.
- Alvarez, J., Marcó, L. M., Arroyo, J., Greaves, E. D. & Rivas, R. (2003). Determination of calcium, potassium, manganese, iron, copper and zinc levels in representative samples of two onion cultivars using total reflection X-ray fluorescence and ultrasound extraction procedure. *Spectrochimica Acta Part B: Atomic Spectroscopy*, 58(12), 21873-2189.
- Ariyama, K., Horita, H & Yasui, A. (2004). Application of inorganic element ratios to chemometrics for determination of the geographic origin of Welsh onions. *Journal of Agricultural and Food Chemistry*, 52(2), 5893-5809.
- Ariyama, K., Nishida, T., Noda, T., Kadokura, M. & Yasui, A. (2006). Effects on fertilisation, crop year, variety, and provenance factors on mineral concentrations in onions. *Journal of Agricultural and Food Chemistry*, 54(9), 3341-3350.

- Ariyama, K. & Yasui, A. (2006). The determination technique of the geographic origin of Welsh onions by mineral composition and perspectives for the future. *Japan Agricultural Research Quarterly*, 40(4), 333-339.
- Ariyama, K., Aoyama, Y., Mochizuki, A., Homura, Y., Kadokura, M. & Yasui, A. (2007). Determination of the geographic origin of onions between three main production areas in Japan and other countries by mineral composition. *Journal of Agricultural and Food Chemistry*, 52(19), 347-354.
- Bibak, A., Behrens, A., Stürup, S., Knudsen, L. and Gundersen, V. (1998). Concentrations of 63 major and trace elements in Danish agricultural crops measured by inductively couple plasma mass spectrometry. I. Onion (*Allium cepa* Hysam). *Journal of Agricultural and Food Chemistry*, 46(8), 3139-3145.
- Chope, G. A, Terry, L. A. & White, P. J. (2006). Effect of controlled atmosphere storage on abscisic acid concentration and other biochemical attributes of onion bulbs. *Postharvest Biology and Technology*, 39(3), 233-242.
- Chope, G. A., Terry, L. A. & White, P. J. (2007). The effect of the transition between controlled atmosphere and regular atmosphere storage on bulbs of onion cultivars SS1, Carlos and Renate. *Postharvest Biology and Technology*, 44(3), 228-239.
- Cole, R. A. & Phelps, K. (1979). Use of canonical variate analysis in the differentiation of swede cultivars by gas-liquid chromatography of volatile hydrolysis products. *Journal of the Science of Food and Agriculture*, 30(7), 669-676.
- Coolong, T. W., Kopsell, D. A., Kopsell, D. E. & Randle, W. M. (2004). Nitrogen and sulphur influence nutrient usage and accumulation in onion. *Journal of Plant Nutrition*, 27(9), 1667-1686.

- Coolong, T. W. & Randle, W. M. (2008). The effects of calcium chloride and ammonium sulphate on onion bulb quality at harvest and during storage. *HortScience*, 43(2), 465-471.
- Food and Agriculture Organisation of the United Nations. (2007). FAOSTAT. [online]. Available at: <http://faostat.fao.org/site/346/Desktop/default.aspx?PageID=346> (Accessed 15 June 2007).
- Gardner, S., Gower, J. C & Le Roux, N. J. (2006). A synthesis of canonical variate analysis, generalised canonical correlation and Procrustes analysis. *Computational Statistics and Data Analysis*, 50(1), 107-134.
- Golubev, F. V., Golubkina, N. A. & Gorbunov, Y. N. (2003). Mineral composition of wild onions and their nutritional value. *Applied Biochemistry and Microbiology*, 39(5), 532-535.
- Gunderson, V., Ellegaard Bechmann, I., Behrens, A. and Stürup, S. (2000). Comparative investigation of concentrations of major and trace elements in organic and conventional Danish agricultural crops. I. Onions (*Allium cepa* Hysam) and Peas (*Pisum sativum* Ping Pong). *Journal of Agricultural and Food Chemistry*, 48(12), 6094-6102.
- Hamilton, B. K., Yoo, K. S. & Pike, L. M. (1998). Changes in pungency of onions by soil type, sulphur nutrition and bulb maturity. *Scientia Horticulturae*, 74(4), 249-256.
- Hansen, R. G., Wyse, B. W. & Sorensen, A. W. (1979). *Nutritional quality index of foods*. AVI Publishing Company, Connecticut, USA.

- Holland B., Unwin, I. D. & Buss, D. H. (1991). *Vegetables, Herbs and Spices*. Fifth supplement to *McCance and Widdowson's The Composition of Foods* (4th Edition). Royal Society of Chemistry, Cambridge.
- Kopsell, D. E., Randle, W. M. & Eiteman, M. A. (1999). Changes in the S-alk(en)yl cysteine sulfoxides and their biosynthetic intermediates during onion storage. *Journal of the American Society for Horticultural Science*, 124(2), 177-183.
- Lancaster, J. E., Farrant, J. & Shaw, M. L. (2001). Sulfur nutrition affects cellular sulfur, dry weight distribution, and bulb quality in onion. *Journal of the American Society for Horticultural Science*, 126(2), 164-168.
- Mallor, C. & Thomas, B. (2008). Resource allocation and the origin of flavour precursors in onion bulbs. *Journal of Horticultural Science and Biotechnology*, 83(2), 191-198.
- O'Donoghue, E. M., Somerfield, S. D., Shaw, M., Bendall, M., Hedderly, D., Eason, J. & Sims, I. (2004). Evaluation of carbohydrates in Pukekohe Longkeeper and Grano cultivars of *Allium cepa*. *Journal of Agricultural and Food Chemistry*, 52(17), 5383-5390.
- Randle, W. M. (1992). Sulfur nutrition affects nonstructural water-soluble carbohydrates in onion germplasm. *HortScience*, 27(1), 52-55.
- Randle, W. M., Bussard, M. L. & Warnock, D. F. (1993). Ontogeny and sulfur fertility affect leaf sulfur in short-day onions. *Journal of the American Society for Horticultural Science*, 118(6), 762-765.

- Randle, W. M., Kopsell, D. E., Kopsell, D. A. & Snyder, R. L. (1999). Total sulphur and sulphate accumulation in onion is affected by sulphur fertility. *Journal of Plant Nutrition*, 22(1), 45-51.
- Rodríguez Galdón, B., Oropeza González, R., Rodríguez Rodríguez, E. & Díaz Romero, C. (2008). Comparison of mineral and trace elements in onion cultivars (*Allium cepa* L.). *Journal of the Science of Food and Agriculture*, 88(9), 1554-1561.
- Smittle, D. A. (1988). Evaluation of storage methods for 'Granex' onions. *Journal of the American Society for Horticultural Science*, 133(6), 877-880.
- U.S. Department of Agriculture. (2004). *USDA nutrient database for standard reference*. Vol. 2004. [online]. Available at: <http://www.nal.usda.gov/fnic/foodcomp> (Accessed: 20 October 2004).
- White, P. J. & Broadley, M. R. (2005). Biofortifying crops with essential mineral elements. *Trends in Plant Science*, 10(12), 586-593.

Figures

Fig. 1.





Fig. 2.



382 **Tables**

383 Table 1.

Element	Renate	Ailsa Craig	SS1	Raw onion ¹	Sweet onion ¹	Raw onion ²	Raw onion ³	RDA ⁴ (mg)	RNI ⁵ (mg)
B	0.58	0.07	0.12	NS	NS	NS	NS	NS	NS
Ca	60.34	45.34	29.22	22.00	20.00	25.00	27.06	1000-1200	700
Cu	0.58	0.08	0.13	0.04	0.06	0.05	NS	0.9	1.2
Fe	0.70	0.34	0.31	0.19	0.26	0.30	0.53	5.0-18.0	11.4
K	270.85	187.64	149.24	144.00	119.00	160.00	157.06	1600-3500	3500
Mg	10.81	7.42	6.39	10.00	9.00	4.00	NS	310-420	300
Mn	0.23	0.13	0.11	0.13	0.08	0.10	NS	1.8-2.3	>1.4
Na	14.57	5.19	3.84	3.00	8.00	3.00	NS	500-2400	1600
P	29.25	19.86	15.92	27.00	27.00	30.00	35.88	700	550
S	67.97	50.66	38.21	NS	NS	51.00	NS	NS	NS
Zn	0.45	0.28	0.27	0.16	0.13	0.20	NS	8.0-11.0	9.5

384

385 NS – not specified

386 ¹Data from USDA database (U.S. Department of Agriculture, 2004) cvs. not specified.

387 ²Data from Holland, Unwin, & Buss (1991) cv. not specified.

388 ³Hansen, Wyse, & Sorensen (1979) cv. not specified.

389 ⁴RDA=Recommended Daily Allowance (US) (White & Broadley, 2005).

390 ⁵RNI=Reference Nutrient Intake (UK) (White & Broadley, 2005).

391 Table 2.

Variate	Canonical variate 1	Canonical variate 2
Boron	0.16033	0.04664
Calcium	-0.00049	-0.00047
Copper	0.03610	0.01124
Iron	0.02923	-0.03678
Potassium	-0.00011	-0.00010
Magnesium	-0.00540	0.01503
Manganese	0.13895	0.01173
Sodium	0.00185	-0.00143
Phosphorus	-0.00499	-0.00149
Sulphur	0.00105	-0.09950
Zinc	-0.01398	0.03573

392

Figure Captions

Fig. 1. Elemental content of onion cvs. Renate, Ailsa Craig and SS1 bulbs after postharvest curing (day 0, black bars, $n = 4$), and at early (light grey bars, $n = 8$) and late (dark grey bars, $n = 8$) time points during controlled atmosphere storage (cv. Renate: days 40 and 230, cv. Ailsa Craig: days 81 and 129, and cv. SS1: days 40 and 81). Standard error bars are shown.

Fig. 2. The scores for the first two canonical variates of the three cultivars. The lines around the groups represent the 95% confidence interval for each group. $\circ$ = cv. SS1, $\times$ = cv. Renate, $+$ = cv. Ailsa Craig.

Table Captions

Table 1. The mean elemental composition of onion cvs. Renate, Ailsa Craig and SS1 bulbs ($n = 20$) compared with data for raw and sweet onions from the USDA database (U.S. Department of Agriculture, 2004) compared to the RDA and RNI values. Values are presented as $\text{mg } 100\text{g}^{-1}$ unless otherwise stated).

Table 2. The mean scores of the first two canonical variates of the minerals analysed.



DSPACE

<https://dspace.org/>

Use of canonical variate analysis to differentiate onion cultivars by mineral content as measured by ICP-AES

Chope, Gemma Amy; Terry, Leon A.

2009-08-01T00:00:00Z

Elsevier

<http://dx.doi.org/10.1016/j.foodchem.2008.12.090>

Gemma A. Chope, Leon A. Terry, Use of canonical variate analysis to differentiate onion cultivars by mineral content as measured by ICP-AES. Food Chemistry, Vol. 115, Issue 3, 1 August 2009, p.1108-1113