

Accelerating abscission of macadamia nuts using ethephon: are there implications for nut quality?

Noluthando Noxolo Aruwajoye^{a,b}, Olaoluwa Omoniyi Olarewaju^c, Abimbola E. Oluwalana-Sanusi^{d,e}, Asanda Mditshwa^a, Lembe Samukelo Magwaza^f and Samson Zera Tesfay^a

^aDiscipline of Crop and Horticultural Science, School of Agricultural, Earth and Environmental Sciences, University of KwaZulu-Natal, Pietermaritzburg, South Africa; ^bDepartment of Civil Engineering, Durban University of Technology, Pietermaritzburg, South Africa; ^cDepartment of Nature Conservation, Faculty of Natural Sciences, Mangosuthu University of Technology, Durban, South Africa; ^dDepartment of Physical and Earth Sciences, Sol Plaatje University, Kimberley, South Africa; ^eCentre for Global Change, Sol Plaatje University, Kimberley, South Africa; ^fPlant Science Laboratory, Cranfield University, Bedfordshire, UK

ABSTRACT

Macadamia is a valuable nut due to its high content of healthy monounsaturated fats, dietary fibre, antioxidants, and essential minerals such as magnesium-B and potassium. One of the major issues affecting industries around the globe is the delayed nut abscission and a prolonged and expensive harvest. A delay in harvesting also reduces the quality of the kernel. To address this challenge, ethephon is used to accelerate abscission, helping to prevent delays in harvesting. Ethephon can cause up to a 15-fold increase in nut abscission within 4 weeks compared to untreated trees. We provide a brief overview of the use of ethephon and its effects on nut quality. The major side-effects of ethephon used to manage macadamia harvesting are associated with ethylene, which is released from ethephon during the acceleration of plant senescence. This process impacts nut quality and shelf life by modulating lipid composition and nut biochemistry. Additionally, ethephon influences the appearance of the nut and the expression and regulation of specific plant genes. This review bridges the information gap between the application of ethephon for harvesting and its effects on the nut quality.

ARTICLE HISTORY

Received 4 June 2024
Accepted 1 November 2024

KEYWORDS

Abscission; ethephon; ethylene; kernel quality; *macadamia integrifolia* and *macadamia tetraphylla*

Introduction

Macadamia is a high-value nut that is cultivated in many countries, including Australia, South Africa, Kenya, USA, China, Guatemala, Malawi, Zimbabwe, and Brazil (Bandason et al., 2022). In 2022, annual revenue was 300 million USD compared with 7.3 million USD in 1996 (Bouarakia et al., 2023). South Africa and Australia accounted for 31% and 23% of exports, respectively (International Nut and Dried Fruit Council, 2022).

Macadamias are native to Australia, where there are four species, two of which produce edible nuts. *Macadamia integrifolia* and *Macadamia tetraphylla* are known as the smooth shell and the rough shell, respectively (Dailey & Vuong, 2015; Nagao et al., 1992). These species are part of the *Proteaceae* botanical family (Canneddu et al., 2016). These nuts stand out for their content of monounsaturated fats, dietary fibre, protein, total sugar, antioxidants, as well as essential minerals such as magnesium and potassium (Figure 1) (Buthelezi et al., 2019; Y. Liu et al., 2022; Rahman et al., 2021; Schlörmann et al., 2017; Yang et al., 2009). The nut in shell contains about 25%

moisture on a dry basis and requires drying. Initially dried by natural convection to 10% moisture, it then undergoes forced convection drying to reduce moisture to around 3.5% on a dry basis (Navarro & Rodrigues, 2016).

Though native to Australia, commercial production of macadamia was initiated in Hawaii during the mid-1920s (O'Connor et al., 2019), and cultivars are now grown across the world. Global production reached 78,415 metric tons in 2022 (Figure 2). In 2021, both global exports and imports reached 37,622 metric tons (International Nut and Dried Fruit Council, 2022). Production and trade are influenced by regional harvesting schedules. In Australia, the harvest takes place from March to September (Brinkhoff & Robson, 2021), whereas, in South Africa, harvest takes place from February to August (SAMAC, 2021). Australia and South Africa export up to 99% of their crop to the USA, China, Japan, and Vietnam (Shabalala et al., 2022). The crop is harvested by collecting nuts from the orchard floor following natural nut abscission (D. A. Walton & Wallace, 2009). Harvesting is affected by cultivar type, rapid fruit abscission, lack of harvesting

CONTACT Samson Zera Tesfay  tesfay@ukzn.ac.za  Discipline of Crop and Horticultural Science, School of Agricultural, Earth and Environmental Sciences, University of KwaZulu-Natal, Private Bag X01, Scottsville, 3209, Pietermaritzburg South Africa

© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.




Figure 1. Nutritional composition of macadamia nuts (Buthelezi et al., 2019; Y. Liu et al., 2022; Rahman et al., 2021; Schlörmann et al., 2017; Yang et al., 2009).




Figure 2. Global macadamia production in metric tons of kernels between 2012 and 2022. Data from the International Nut and Dried Fruit Council (2022).

equipment, inadequate on-farm storage, the inability of processors to receive the product, and weather (Trueman et al., 2002; D. A. Walton & Wallace, 2009). Delayed abscission has been identified as a significant factor contributing to the pattern of commercial nut drop of some cultivars (Trueman et al., 2002). The nuts can remain on the trees for a long period, adversely affecting the quality of the kernels. Hence, there is a need to hasten the abscission process. Macadamia has a high oil content of fatty acids and is susceptible to rancidity and rapid deterioration (Gama et al., 2018). D. A. Walton and Wallace (2009) reported that, when the harvest is delayed, deterioration is more rapid in later-harvested nuts than in those harvested earlier.

Kernels from late-harvested nuts have high peroxide values, indicating that lipid oxidation is initiated several months before harvest whilst nuts are still attached to the tree (Gama et al., 2020).

It is crucial to harvest nuts soon after maturity to reduce the loss of quality (Gama et al., 2020; D. A. Walton & Wallace, 2009). However, variations in abscission necessitate the use of cultural practices to hasten nut abscission. This can be accomplished by shaking the trees or by applying ethephon (Quinlan et al., 2008). The combined use of ethephon and shaking has been proven to be the most efficient protocol (Trueman et al., 2002). Ethephon is applied before shaking to reduce the nut removal force, but the use of ethephon alone is the most practical method (D.

Walton & Wallace, 2008). Ethephon releases ethylene after application and accelerates nut abscission (Trueman, 2010). Challenges from delayed abscission and multiple harvests can be mitigated through the use of ethephon.

Therefore, we reviewed current knowledge regarding the effect of ethephon on the harvesting of macadamia. The paper explores possible side-effects of ethephon on kernel quality and shelf life. The findings of this review will help producers make better decisions to maintain profitable macadamia production.

Stages of nut development

The stages of nut development, including ripening and ageing, are closely tied to the tree's growth cycle. This cycle includes floral initiation, flowering, nut set, nut development, oil accumulation, and nut abscission (Figure 3) (Stephenson et al., 2003). Mature trees produce over 10,000 inflorescences (racemes), bearing about 100–300 flowers (Prasannath et al., 2021). Each tree has close to 1–3 million flowers. However, yield does not reflect flower production. Howlett et al. (2015) reported that 0.3% of the flowers resulted in harvestable nuts. This is because a high percentage of flowers drop because they are not fertilised (du Queens et al., 2018). Factors contributing to this issue include inadequate pollination, poor weather, and nutrient deficiencies (De Silva et al., 2022). Identifying the causes of poor nut set can potentially improve yields. Macadamia is partially self-

incompatible pollinated and pollinated by bees. Some cultivars are self-compatible; however, they can have higher yields after cross-pollination. Cross-pollination increases yield by up to 95% compared to self-pollination (da Silva Santos et al., 2020; Trueman, 2013). Pollination can be limited by non-pollinator-friendly management and high pesticide use (Grass et al., 2018). Once the fruit have set, nut growth increases until the nuts are mature and have a hard shell (Nortjé, 2018). Subsequently, sugars in the kernels are converted to oil (Nortjé, 2018). Kernels with high sugar content are characterised as immature (Gama et al., 2018; Wall & Gentry, 2007). The highest quality kernels have an oil content of 72–78% (Wall, 2013). Following oil accumulation, the mature nut drops from the trees. However, this stage varies among cultivars and seasons (Trueman, 2013). The extended nut drop poses a challenge to the industry. In the next section, we explore the benefits of plant growth regulators, such as ethephon.

Use of ethephon and other plant growth regulators

Abscission occurs naturally in plants, resulting in the shedding or separation of leaves, flowers, fruit and branches from the parent plant (Meir et al., 2019). The process takes place at the abscission zone and is divided into four phases. The first step includes the differentiation of cells that are present within the abscission zone. This is followed by the second step




Figure 3. The phenological stages of the macadamia.

where these cells acquire the competence to respond to abscission signals. This is followed by loosening of the cell wall and separation (nut drop). Finally, a protective layer is formed over the abscission zone (J. Kim, 2014; G. Liu et al., 2020a; Patharkar & Walker, 2018; Rai et al., 2021). Two models of abscission have been suggested: senescence-triggered abscission and correlatively driven abscission (Ackerman & Samach, 2015; Bangerth, 2000; Giulia et al., 2013). Senescence-triggered abscission is a natural process that occurs in ripe fruit and leaves where shedding is programmed (Giulia et al., 2013). In this model, as the leaf or fruit matures, ethylene production is triggered, in turn reducing auxin production and basipolar transport of indolyl acetic acid to the abscission zone. This leads to increased sensitivity of the tissue to ethylene, ultimately initiating abscission (Bangerth, 2000). However, this model focuses on mature fruit and doesn't account for the abscission of young fruit. Correlative-driven abscission involves the natural shedding of young fruit, driven by the interplay between neighbouring fruit and nearby shoots (Giulia et al., 2013). This process reflects the regulation of fruit load based on nutritional factors, ensuring effective resource utilisation (Giulia et al., 2013). A fruitlet with a stronger ability to attract and utilise nutrients is more likely to accumulate greater

resources compared with neighbour with a lower capacity. The fruitlet possessing higher 'sink strength' has a better chance of surviving (Ackerman & Samach, 2015).

Furthermore, the movement of auxins within fruitlets, encounters specific points termed 'autoinhibitory junctions' along the stem or peduncle (Larson et al., 2022). These junctions act as points of interaction where auxin flow from dominant fruitlets can hinder the movement of auxin from less dominant ones. The reduced downward movement of auxin through the pedicel and heightened sensitivity to ethylene in the abscission zone triggers the abscission of fruitlets (Hagemann et al., 2015). Although ethylene contributes to the final stages of fruit detachment, it is not the initial catalyst for the process, as is the case with abscission prompted by senescence (Bangerth, 2000). The ability of ethephon to release ethylene has made its use practical in accelerating the abscission of macadamia nuts. Ethylene released from ethephon causes the cell inside the pedicel abscission zone to produce cell wall degrading enzymes that are responsible for breaking down the pedicel abscission zone which leads to the detachment of leaves, flowers, fruit, and branches from the parent plant (Hagemann et al., 2015). A systematic process of ethylene-induced abscission is illustrated in Figure 4. Briefly, when




Figure 4. The ethylene-induced abscission process in macadamia (adapted from Yamada et al., 2015).

ethylene is released, it acts as a signal to initiate the synthesis of wall-degrading enzymes, such as cellulase and polygalacturonase. These enzymes are instrumental in breaking down the primary cell wall and the middle lamella, a process crucial for facilitating cellular separation. This step is followed by the breakdown of the cell wall, which results in cell separation and organ shedding (Yamada et al., 2015).

Evaluation of the application of ethephon for macadamia nut abscission

The high costs associated with multiple harvest times and cultivars exhibiting late-season abscission are the major motivations for the use of ethephon to promote nut abscission. Studies done by Trueman et al. (2002) and Trueman (2003) demonstrated that ethephon significantly reduced the harvest duration for macadamia nuts. In the study by Trueman et al. (2002), ethephon-treated plots achieved up to 96% nut drop within 4 weeks, compared to 62% in untreated controls. The study by Trueman (2003) found that ethephon caused up to 86.7% nut drop within the same period, whereas untreated controls saw only 43.2% nut drop (Table 1). These findings collectively demonstrated that ethephon effectively accelerated nut abscission, resulting in a shorter overall harvest period and improved harvesting efficiency. Reports investigating abscission without the use of ethephon revealed that fruit maturation and the timing of abscission are independent in macadamia (Trueman et al., 2000). This necessitates the use of ethephon on cultivars exhibiting delayed abscission. Ethephon applied to 'A16' (a macadamia cultivar exhibiting late-season abscission)

effectively induced abscission and had no adverse effects on quality but decreased yields (Trueman, 2003). Ethephon reduced macadamia nut yields significantly, with decreases up to 2.39 kg in unshaken trees mid-season and 2.02 kg (Trueman, 2003). Richardson and Dawson (1993) observed lower yields in the following season when ethephon was applied 2 or 4 weeks after nut maturity (Richardson & Dawson, 1993). This response was primarily attributed to the loss of leaves caused by the chemical (Table 2) (Trueman, 2003). Philips (1970) reported that a decrease in yield is associated with leaf loss. Burns et al. (2003) found that ethephon led to 100% leaf drop after 10 days, providing further evidence of its negative impact on the canopy. When ethephon is applied to plants, it reduces the number of leaves available for photosynthesis. Consequently, the plant's capacity to assimilate carbon dioxide (CO₂) is diminished, in turn resulting in lower yields (Trueman, 2003). However, a recent study by G. Liu et al. (2020b) demonstrated that defoliation does not always lead to yield loss (G. Liu et al., 2020b). Optimized leaf removal can enhance light distribution, subsequently increasing canopy photosynthesis (G. Liu et al., 2020b).

Factors influencing the efficacy of ethephon

In acidic solutions, ethephon remains stable, whereas it hydrolyses in plant tissues with higher pH levels (Torres & Asín, 2022). This hydrolysis releases ethylene, which, upon external application, triggers ethylene-dependent processes such as flower or fruit abscission (Torres & Asín, 2022). In flowers and climacteric fruit, ethephon is exogenously applied but

Table 1. Influence of ethephon concentration on nut drop in various macadamia cultivars.

Ethephon concentration (mg/L)	Cultivar	Untreated trees nut drop	Ethephon treated nut drop %	References
400	'Beaumont'	6%	87%	Richardson and Dawson (1993)
500	'Beaumont'	5%	95%	Kadman and Ben-Tal (1983)
600	'A16'	43.2%	83.6% ± 3.8%	Trueman (2003)
900	'A16'	43.2%	78.9% ± 3.4%	Trueman (2003)
1000	'Ikaika'	–	60%	Nagao and Sakai (1987)
1200	'A16'	43.2%	86.7% ± 3.4%	Trueman (2003)
1600	'A16'	18.29%	93.77%	Jing et al. (2018)

Table 2. Effects of ethephon application on leaf drop, yield, and nut biochemistry in macadamia trees, based on various studies.

Parameter	Effect	Reference
 Leaf drop	Leaf abscission was observed in ethephon treated trees, particularly among older leaves within the canopy, though no specific counts were made.	Trueman et al. (2000)
 Yield	Ethephon application was linked to leaf drop, although it was not quantitatively measured. Application of ethephon at 400 mg/L reduced subsequent yield to 48 nuts per tree, compared to 121 nuts per tree in untreated controls. There was no observed negative impact on the yield following ethephon application.	Trueman (2003) Richardson and Dawson (1993) Jing et al. (2018)
 Nut biochemistry	No significant changes were noted in kernel oil content. However, an increase in nut-in-shell (NIS) weight was observed post-ethephon treatment, although this change was not statistically significant. Ethephon application significantly influenced fatty acid composition, total phenolics, and flavonoid content in nuts.	Trueman (2003) Aruwajoye et al. (2023a) Aruwajoye et al. (2023b)

induces endogenous ethylene biosynthesis. Two distinct systems have been acknowledged in regulating ethylene production. The first is System 1, where ethylene suppresses its biosynthesis during vegetative growth. System 2 operates exclusively during climacteric fruit ripening in specific species and is characterised by an autocatalytic ethylene biosynthesis process that intensifies production (Argueso et al., 2007). Autocatalytic ethylene production involves ethylene-induced upregulation of enzymes such as 1-aminocyclopropane-1-carboxylate synthase and 1-aminocyclopropane-1-carboxylate oxidase, which play a crucial role in the rate-limiting reaction of ethylene biosynthesis (Gorny & Kader, 1997). Marei and Crane (1971) reported that exogenous ethylene stimulates the tissue to produce more ethylene. However, a recent study by Reig et al. (2016) on loquat fruit suggests that the ethylene released by ethephon might have a transient effect, lacking the ability to trigger the autocatalytic biosynthesis of ethylene and, as a result, the subsequent ripening process. The response to ethephon can be inconsistent due to cultivar, weather, chemical rate, time of application and spray additives (Jing et al., 2018; Torres et al., 2021). Some cultivars exhibit a differential response to ethephon. For example, kiwifruit has two commercially important species (*Actinidia deliciosa* and *A. chinensis*). *A. chinensis* exhibit higher endogenous ethylene production (Ampa et al., 2016; Asiche et al., 2016). This may be attributed to genetics and higher levels of ethylene biosynthesis enzymes. Temperature affects the degradation of ethephon, with degradation more rapid at higher temperatures (Ali et al., 2012; Walters & Lopez, 2018). Ethephon applied under low temperatures has been associated with poor abscission, whereas in high temperatures, the growth

regulator may cause over-thinning or defoliation (Yuan, 2007). Moreover, research has optimised the use of ethephon for nut abscission, focusing on the solution's pH and effective concentration (Jing et al., 2018). However, as a strong acid, ethephon can alter the final pH of the solution, potentially impacting its effectiveness (Camberato et al., 2014). Higher efficiencies are attained if the age of the nut is also considered (Nagao & Sakai, 1988). The sensitivity to ethephon varies with the stage of nut development. Nuts that are small (3–7 weeks after flowering) and mature nuts nearing maturity (28–30 weeks after flowering) respond to ethephon (Jones, 2002). In contrast, mid-size nuts are not sensitive to the chemical (Jones, 2002). This suggests that the timing to ethephon application can affect the rate of nut drop.

Application of ethephon on nut trees

Ethephon has many benefits but also undesirable side-effects (Table 3; Atli et al., 2021; Grijalva-Contreras et al., 2011; Kamiab et al., 2020; Wang et al., 2016a). Ethephon induces synchronised flowering but also causes massive defoliation (Jhade et al., 2019). Ethephon can induce fruit abscission or loosening, but this can come at the expense of defoliation (Burns et al., 2008). For example, ethephon applied to table olives reduced fruit detachment force but defoliated the plants (Burns et al., 2008). Khalil et al. (2016) reported that high rates of ethephon caused leaf and nut drop and decreased total kernel weight of pecan nuts (Khalil et al., 2016). Similarly, the use of ethephon reduced the size and number of pistachio nuts, reducing market value (Rahemi & Ramezani, 2007). Ethephon maintained kernel quality when walnuts were harvested within 10–14 days of treatment

Table 3. The effect of ethephon on selected tree nuts.

Tree nut	Concentration of ethephon	Benefits	Side-effects	References
Almonds (<i>Prunus dulcis</i>)	75–300 mg/L	Delayed bloom.	Up to 3-fold yield reduction.	Grijalva-Contreras et al. (2011)
Pecan (<i>Carya illinoensis</i>)	200, 400, and 600 mg/L	Increased leaf chlorosis and kernel quality.	Reduced total kernel weight, and leaf defoliation.	Khalil et al. (2015)
Pecan (<i>Carya illinoensis</i> (Wangenh.) C. Koch)	600 mg/L	Increased fruit set.	Decreased total kernel weight.	Khalil et al. (2016)
Walnuts (<i>Juglans regia</i> L)	40, 70, and 100 mL/100 L	Increase in hull-dehiscence.	Slight yellowing of the leaves.	Lang and Evans (2010)
Walnuts (<i>Juglans regia</i> L)	0, 10, 500, and 8000 mg/L	Low concentration enhances resistance and retards the decay of fresh in-husk nuts.	High concentration promotes senescence and decay of the fruit.	Wang et al. (2016)
Pistachos (<i>Pistacia vera</i> L)	100 and 200 mg/L	Increased size of the nut, floral bud retention, weight of kernel, and decreased % of seedless nuts.	Reduced number of fruit per cluster.	Rahemi and Ramezani (2007)
Pistachos (<i>Pistacia vera</i> L)	1000, 3000, and 5000 mg/L	Improved cluster breaking force from the branch and endocarp removal force from the seed.	–	Atli et al. (2020)
Pistachos (<i>Pistacia vera</i> L)	75, 150, and 200 mg/L	Decreased flower bud abscission and increased yield, nuts per ounce, blank and split nuts percentage improved.	–	Kamiab et al. (2020)
Cashew (<i>Anacardium occidentale</i> L)	50 mg/L	Increased nut yield and the number of fruit set and decreased fruit drop.	–	Lakshmipathi et al. (2014)

but decreased quality when harvesting was delayed (Lang & Evans, 2010). In some cases, the side-effects of ethephon are minor (Table 3; Lang & Evans, 2010). Ethephon has been studied for its potential application on almond trees since the flowers are vulnerable to frost damage (Omidifard & Gharaghani, 2022). Grijalva-Contreras et al. (2011) assessed the effect of ethephon on delaying bloom and found that application of ethephon at the 10% leaf drop resulted in a delay of bloom by 7, 8 and 9 days at 75, 150 or 300 mg L⁻¹.

Factors affecting the post-harvest quality of macadamias

Nuts, including macadamia, are susceptible to oxidative rancidity and rapid deterioration due to their high oil content and fatty acid composition (Gama et al., 2018). Pre-harvest, harvest, and post-harvest conditions affect nut biochemistry and quality. Gama et al. (2020) reported that macadamias harvested later in the season had a shorter shelf life and lower oil quality than those harvested earlier (higher free fatty acid levels). Similarly, oil extracted from early harvested almonds had superior quality (lower K232 and K270 values and a higher K232/K270 ratio) compared to late-harvested almonds (Kazantzis et al., 2003). Based on Kazantzis et al. (2003) findings, the degradation of oil quality was attributed to the increase in UV absorption coefficients K232 and K270, which are respectively linked to the formation of conjugated hydroperoxides from lipid oxidation and the presence of hydroperoxide breakdown products. On the other hand, in *Torreya grandis*, later harvested nuts had lower free fatty acid levels and peroxide values compared to early harvested nuts (Y. Wang et al., 2021). The findings suggest that the optimal time of harvesting varies with the species and its susceptibility to lipid oxidation. The quality of all nuts, including macadamia, can be revealed by various indices such as oil content, lipid oxidation, moisture content, and rancidity (Buthelezi et al., 2019). Oil content is the most critical quality index in macadamia because the nut has unsaturated fatty acids levels greater than 80% (Buthelezi et al., 2019; Rengel et al., 2015).

The effects of ethephon on the post-harvest quality of macadamias

Ethylene released from ethephon can result in changes in nut quality. The abscission process is regulated by a complex series of structural, physiological, biochemical, and molecular changes, which influence nut biochemistry (Qiu et al., 2020). Aruwajoye et al. (2023b) examined the impact of ethephon, designed to stimulate nut detachment, on the fatty acid composition of macadamias ('Beaumont' and '788') during post-harvest storage. Ethephon increased stearic acid and

saturated fatty acids (SFAs) in 'Beaumont' and decreased unsaturated fatty acids (USFAs) (Aruwajoye et al., 2023b). Similar results were observed in *Chlorella vulgaris* (*C. vulgaris*) (S. H. Kim et al., 2016). Exogenous ethylene increased stearic acid and total saturated fatty acids in walnuts when used for hulling (Farooq et al., 2023). Ethylene-induced fruit abscission in oil palm and increased quality as reflected by lower Free Fatty Acids (FFA) and higher oxidative stability (Chew et al., 2021). The treated fruits had a higher triacylglycerol, which is also found in macadamias (Chew et al., 2021). These observations emphasise the effect of ethylene on the nutrition and storage of macadamias.













The application of ethephon has been associated with sugar accumulation and changes in phenolic metabolism in fruits and nuts (Aruwajoye et al., 2023a; Li et al., 2002; Wang et al., 2016a). Wang et al. (2016a) indicated that ethephon improved the storage life of walnuts by slowing the role of decay, inhibiting browning-related enzymatic activities, and preserving kernel quality (Wang et al., 2016a). However, at higher ethephon concentrations, a reduction in phenolic content and an increase in polyphenol oxidase activity were observed. This highlights the importance of considering the applied concentration of ethephon.

Effect of ethephon on gene expression and regulation

The impact of ethephon is also evident in its influence on the expression and regulation of specific genes. For instance, Y. Hu et al. (2022) found that exposing *Torreya grandis* nuts to exogenous ethylene enhanced cracking rates, elevated squalene content, and reduced β -sitosterol levels. This indicates an accelerated post-ripening process that influences the biosynthesis of bioactive compounds through ethylene-induced gene expression in the mevalonate pathway, particularly involving TgHMGS and TgHMGR. Ethephon used to induce abscission in *Camellia oleifera* resulted in the upregulation of CoACO1 and CoACO2, which are genes that encode for 1-aminocyclopropane-1-carboxylate oxidase (ACO) enzymes (X. Hu et al., 2021). This is important to note because *Camellia oleifera* has similarities to macadamia nuts; both contain high levels of monounsaturated fats and have a similar fatty acid profile. While the genetic and biochemical responses of *Camellia oleifera* to ethephon are reported (X. Hu et al., 2021), similar responses in macadamia nuts treated with ethephon would need to be investigated.

Moreover, a study by Z. Zhang et al. (2023) found that exogenous ethylene treatment in *Torreya grandis* significantly increased amino acid levels, particularly enhancing umami amino acids

Table 4. The effect of ethylene on the amino acid content of nuts (adapted from Farooq et al., 2023 and Z. Zhang et al., 2023).

Amino acids	Walnuts	<i>Torreya grandis</i> nuts
 Proline	–	Decrease
 Aspartic acid	Decrease	Increase
 Serine	Increase	Increase
 Glutamine acid	Increase	Increase
 Glycine	Decrease	Increase
 Tyrosine	Increase	Not significant
 Arginine	Increase	Increase
 Histidine	Increase	Increase
 Lysine	Decrease	Not significant
 Alanine	Decrease	Not significant
 Phenylalanine	Decrease	Decrease
 Isoleucine	Decrease	Not significant
 Leucine	Decrease	Not significant

like glutamic and aspartic acids (the effect of ethephon on other amino acids is shown in Table 4). This occurred through the upregulation of biosynthesis genes and downregulation of degradation enzymes, ultimately improving post-harvest nut quality. These findings collectively suggest that exogenous ethylene treatment has the potential to improve both the physical and biochemical aspects of *Torreya grandis* nuts, contributing to enhanced post-harvest nut quality.

Conclusions

Ethephon in macadamia production has emerged as a critical tool, promoting uniform nut abscission and easier harvesting. However, a limitation arises when this growth regulator hastens the ripening of the macadamia nuts. Ethephon modulates lipid composition and nut biochemistry. The chemical also affects nut appearance, as well as the expression and regulation of specific genes. Future research should focus on determining optimal application rates and timing for different cultivars. These studies should assess physical, biochemical, and genetic responses. A better protocol for the use of ethephon will help the expansion and viability of this important nut crop.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was funded by the National Research Foundation (NRF) of South Africa [Grant Number 142060].

Authors' contribution statement

The study was conceptualized by S. Z. Tesfay and N. N. Aruwajoye. The first draft was written by S. Z. Tesfay, N. N. Aruwajoye, and O. O. Olarewaju. The final manuscript was prepared by S. Z. Tesfay, N. N. Aruwajoye, O. O. Olarewaju, and A. E. Oluwalana-Sanusi. Supervision was provided by S. Z. Tesfay, L. S. Magwaza, and A. Mditshwa. Visualization was handled by N. N. Aruwajoye, O. O. Olarewaju, and A. E. Oluwalana-Sanusi.

Data availability statement

Data sharing does not apply to this article since no new data were generated or analysed.

References

- Ackerman, M., & Samach, A. (2015). Doubts regarding carbohydrate shortage as a trigger toward abscission of specific apple (*Malus domestica*) fruitlets. *New Negatives in Plant Science*, 1, 46–52. <https://doi.org/10.1016/j.neps.2015.06.003>
- Ali, T., John, A. C., Subramanian, J., & Peter, K. (2012). Berberine prolongs life span and stimulates locomotor activity of drosophila melanogaster. *American Journal of Plant Sciences*, 3(7A), 1037–1040. <https://doi.org/10.4236/ajps.2012.327123>
- Ampa, K., Ikeura, H., Saito, T., Okawa, K., Ohara, H., & Kondo, S. (2016). Effects of pre-harvest application of ethephon or abscisic acid on 'Kohi' kiwifruit (*Actinidia chinensis*) ripening on the vine. *Scientia Horticulturae*, 209, 255–260. <https://doi.org/10.1016/j.scientia.2016.06.026>
- Argueso, C. T., Hansen, M., & Kieber, J. J. (2007). Regulation of ethylene biosynthesis. *Journal of Plant Growth Regulation*, 26(2), 92–105. <https://doi.org/10.1007/s00344-007-0013-5>

- Aruwajoye, N. N., Mditshwa, A., Magwaza, L. S., Ngidi, M. S. C., & Tesfay, S. Z. (2023a). Effect of preharvest ethephon application on selected biochemical components and polyphenol oxidase activity in macadamia nuts. *Horticulturae*, 9(10), 1101. <https://doi.org/10.3390/horticulturae9101101>
- Aruwajoye, N. N., Mditshwa, A., Magwaza, L. S., Ngidi, M. S. C., & Tesfay, S. Z. (2023b). Impact of preharvest ethephon foliar spray on the postharvest fatty acid profile and dietary indicators of macadamia nuts. *Agriculture*, 13(10), 1898. <https://doi.org/10.3390/agriculture13101898>
- Asiche, W. O., Mworia, E. G., Oda, C., Mitalo, O. W., Owino, W. O., Ushijima, K., Nakano, R., & Kubo, Y. (2016). Extension of shelf-life by limited duration of propylene and 1-MCP treatments in three kiwifruit cultivars. *The Horticulture Journal*, 85(1), 76–85. <https://doi.org/10.2503/hortj.MI-066>
- Atli, H. S., Ilikçioğlu, E., & Çelik, F. (2021). Effect of preharvest ethephon application on harvest mechanization of Uzun and Siirt pistachio varieties.
- Bandason, W., Parwada, C., & Mushunje, A. (2022). Macadamia nuts (*Macadamia intergrifolia*) value chain and technical efficiency among the small-scale farmers in Zimbabwe. *Research on World Agricultural Economy*, 3(4), 25–35. <https://doi.org/10.36956/rwae.v3i4.700>
- Bangerth, F. (2000). Abscission and thinning of young fruit and their regulation by plant hormones and bioregulators. *Plant Growth Regulation*, 31(1/2), 43–59. <https://doi.org/10.1023/A:1006398513703>
- Bouarakia, O., Anders, M., Linden, V. M., Grass, I., Westphal, C., Taylor, P. J., & Foord, S. H. (2023). Reduced macadamia nut quality is linked to wetter growing seasons but mitigated at higher elevations. *Journal of Agriculture and Food Research*, 12, 100569. <https://doi.org/10.1016/j.jafr.2023.100569>
- Brinkhoff, J., & Robson, A. J. (2021). Block-level macadamia yield forecasting using spatio-temporal datasets. *Agricultural and Forest Meteorology*, 303, 108369. <https://doi.org/10.1016/j.agrformet.2021.108369>
- Burns, J. K., Ferguson, L., Glozer, K., Krueger, W. H., & Rosecrance, R. C. (2008). Screening fruit loosening agents for black ripe processed table olives. *Horticulture Science*, 43(5), 1449–1453. <https://doi.org/10.21273/HORTSCI.43.5.1449>
- Burns, J. K., Pozo, L. V., Arias, C. R., Hockema, B., Rangaswamy, V., & Bender, C. L. (2003). Coronatine and abscission in citrus. *Journal of the American Society for Horticultural Science*, 128(3), 309–315. <https://doi.org/10.21273/JASHS.128.3.0309>
- Buthelezi, N. M. D., Tesfay, S. Z., Ncama, K., & Magwaza, L. S. (2019). Destructive and non-destructive techniques used for quality evaluation of nuts: A review. *Scientia Horticulturae*, 247, 138–146. <https://doi.org/10.1016/j.scienta.2018.12.008>
- Camberato, D. M., Camberato, J. J., & Lopez, R. G. (2014). Determining the effect of carrier water pH and bicarbonate concentration on final pH of plant growth regulator solutions. *Horticulture Science*, 49(9), 1176–1182. <https://doi.org/10.21273/HORTSCI.49.9.1176>
- Canneddu, G., Júnior, L. C. C., & de Almeida Teixeira, G. H. (2016). Quality evaluation of shelled and unshelled macadamia nuts by means of near-infrared spectroscopy (NIR). *Journal of Food Science*, 81(7), C1613–C1621. <https://doi.org/10.1111/1750-3841.13343>
- Chew, C. L., Tan, B. A., Low, J. Y. S., Mohd Hakimi, N. I. N., Kua, S. F., & Lim, C. M. (2021). *Exogenous ethylene application on postharvest oil palm fruit bunches improves crude palm oil quality*. Food Science & Nutrition.
- Dailey, A., & Vuong, Q. V. (2015). Optimization of aqueous extraction conditions for recovery of phenolic content and antioxidant properties from macadamia (*Macadamia tetraphylla*) skin waste. *Antioxidants*, 4(4), 699–718. <https://doi.org/10.3390/antiox4040699>
- da Silva Santos, R., de Oliveira Milfont, M., Silva, M. M., Carneiro, L. T., & Castro, C. C. (2020). Butterflies provide pollination services to macadamia in northeastern Brazil. *Scientia Horticulturae*, 259, 108818. <https://doi.org/10.1016/j.scienta.2019.108818>
- De Silva, A. L., Kämper, W., Wallace, H. M., Ogbourne, S. M., Hosseini Bai, S., Nichols, J., & Trueman, S. J. (2022). Boron effects on fruit set, yield, quality and paternity of macadamia. *Agronomy*, 12(3), 684. <https://doi.org/10.3390/agronomy12030684>
- du Queens, F. N., de Queensland, S. N., & Joubert, A. J. (2018). Macadamia. In: Monselise, S.P. (ed.) *CRC handbook of fruit set and development* (pp. 247–252). CRC Press.
- Farooq, A., Hussain, S. Z., Bhat, T. A., Naseer, B., & Shafi, F. (2023). Walnut fruit: Impact of ethylene assisted hulling on in vitro antioxidant activity, polyphenols, PUFAs, amino acids and sensory attributes. *Food Chemistry*, 404, 134763. <https://doi.org/10.1016/j.foodchem.2022.134763>
- Gama, T., Wallace, H. M., Trueman, S. J., & Hosseini-Bai, S. (2018). Quality and shelf life of tree nuts: A review. *Scientia Horticulturae*, 242, 116–126. <https://doi.org/10.1016/j.scienta.2018.07.036>
- Gama, T., Wallace, H. M., Trueman, S. J., Jones, K., & Hosseini-Bai, S. (2020). Late-dropping macadamia nuts have reduced shelf life. *Scientia Horticulturae*, 268, 109378. <https://doi.org/10.1016/j.scienta.2020.109378>
- Giulia, E., Alessandro, B., Mariano, D., Andrea, B., Benedetto, R., & Angelo, R. (2013). Early induction of apple fruitlet abscission is characterized by an increase of both isoprene emission and abscisic acid content. *Plant Physiology*, 161(4), 1952–1969. <https://doi.org/10.1104/pp.112.208470>
- Gorny, J. R., & Kader, A. A. (1997). Low oxygen and elevated carbon dioxide atmospheres inhibit ethylene biosynthesis in preclimacteric and climacteric apple fruit. *Journal of the American Society for Horticultural Science*, 122(4), 542–546. <https://doi.org/10.21273/JASHS.122.4.542>
- Grass, I., Meyer, S., Taylor, P. J., Foord, S. H., Hajek, P., & Tscharnkte, T. (2018). Pollination limitation despite managed honeybees in South African macadamia orchards. *Agriculture, Ecosystems & Environment*, 260, 11–18. <https://doi.org/10.1016/j.agee.2018.03.010>
- Grijalva-Contreras, R. L., Martínez-Díaz, G., Macías-Duarte, R., & Robles-Contreras, F. (2011). Effect of ethephon on almond bloom delay, yield, and nut quality under warm climate conditions in northwestern Mexico. *Chilean Journal of Agricultural Research*, 71(1), 34. <https://doi.org/10.4067/S0718-58392011000100004>
- Hagemann, M. H., Winterhagen, P., Hegele, M., & Wünsche, J. N. (2015). Ethephon induced abscission in mango: Physiological fruitlet responses. *Frontiers in Plant Science*, 6, 706. <https://doi.org/10.3389/fpls.2015.00706>
- Howlett, B. G., Nelson, W. R., Pattermore, D. E., & Gee, M. (2015). Pollination of macadamia: Review and opportunities for improving yields. *Scientia Horticulturae*, 197, 411–419. <https://doi.org/10.1016/j.scienta.2015.09.057>
- Hu, X., Yang, M., Gong, S., Li, H., Zhang, J., Sajjad, M., Ma, X., & Yuan, D. (2021). Ethylene-regulated immature fruit abscission is associated with higher expression of

- CoACO genes in *Camellia oleifera*. *Royal Society Open Science*, 8(6), 202340. <https://doi.org/10.1098/rsos.202340>
- Hu, Y., Suo, J., Jiang, G., Shen, J., Cheng, H., Lou, H., Yu, W., Wu, J., & Song, L. (2022). The effect of ethylene on squalene and β -sitosterol biosynthesis and its key gene network analysis in *Torreya grandis* nuts during post-ripening process. *Food Chemistry*, 368, 130819. <https://doi.org/10.1016/j.foodchem.2021.130819>
- International Nut and Dried Fruit Council. (2022). Macadamias. In *Nuts and dried fruits statistical yearbook 2021/2022*, (pp. 30–33). The International Nut and Dried Fruit Council.
- Jhade, R., Sawarkar, S., Ambulkar, P., Pannase, S., & Alawa, S. (2019). Foliar application of ethephon (etheph) for quick defoliation in pomegranate (*Punica granatum* L.) var Bhagwa. *IJCS*, 7(2), 1497–1499.
- Jing, M., Jin, L., He, X.-Y., Chen, L.-L., Li, T., & Ni, S.-B. (2018). Research on enhancing abscission of macadamia nuts with ethephon. *Journal of Southwest University*, 40(7), 18–24.
- Jones, V. P. (2002). *Macadamia integrated pest management: IPM of insects and mites attacking macadamia nuts in Hawaii*. University of Hawaii.
- Kadman, A., & Ben-Tal, Y. (1983). Inducing macadamia nutdrop with ethephon. *Horticulture Science*, 18(2), 240–242. <https://doi.org/10.21273/HORTSCI.18.2.240>
- Kamiab, F., Rasouli, P., & ZamaniBahramabadi, E. (2020). Application of some bloom thinning techniques to overcome alternate bearing in pistachio (*Pistacia vera* L.). *Horticulture, Environment and Biotechnology*, 61(1), 31–39. <https://doi.org/10.1007/s13580-019-00187-9>
- Kazantzis, I., Nanos, G. D., & Stavroulakis, G. G. (2003). Effect of harvest time and storage conditions on almond kernel oil and sugar composition. *Journal of the Science of Food & Agriculture*, 83(4), 354–359. <https://doi.org/10.1002/jsfa.1312>
- Khalil, S., Mexal, J., Abdur, R., Said, W., Khalil, I., Zahid, H., & Khan, A. (2015). Management of alternate bearing dilemma in pecan through foliar ethephon thinning in New Mexico, USA. 2nd ICSAE 2015, International Conference on Sustainable Agriculture and Environment, September 30–October 03, 2015, I & II, Konya, Turkey. Proceedings book.
- Khalil, S., Mexal, J., Khalil, I., Wahab, S., Rehman, A., Hussain, Z., Khan, A., Khan, A., & Khattak, M. (2016). Foliar ethephon fruit thinning improves nut quality and could manage alternate bearing in pecan. *Pharmaceutical Chemistry Journal*, 3, 150–156.
- Kim, J. (2014). Four shades of detachment: Regulation of floral organ abscission. *Plant Signaling & Behavior*, 9(11), e976154. <https://doi.org/10.4161/15592324.2014.976154>
- Kim, S. H., Lim, S. R., Hong, S. J., Cho, B. K., Lee, H., Lee, C. G., & Choi, H. K. (2016). Effect of ethephon as an ethylene-releasing compound on the metabolic profile of *Chlorella vulgaris*. *Journal of Agricultural and Food Chemistry*, 64(23), 4807–4816. <https://doi.org/10.1021/acs.jafc.6b00541>
- Lakshmipathi, J., Adiga, D., Kalaivanan, D., Mohana, G., & Meena, R. (2014). Effect of growth regulators on leaf area and yield of cashew (*Anacardium occidentale* L.) var Bhaskara. *Ecology Environment & Conservation*, 20, 9–11.
- Lang, M., & Evans, K. (2010). Advancing hull maturity of Australian walnuts. *Australian Nutgrower*, 24(4), 14–17.
- Larson, J. E., Kon, T. M., & Malladi, A. (2022). Apple fruitlet abscission mechanisms. *Horticultural Reviews*, 49, 243–274. <https://doi.org/10.1002/9781119851981.ch5>
- Li, Y., Yang, L., Li, Y., & Ye, Y. (2002). Influence of ethephon sprayed at different stages on growth, agronomic traits and drought resistance of sugarcane. *Sugarcane*, 9(1), 12–18.
- Liu, D., Li, J., Li, Z., & Pei, Y. (2020a). Hydrogen sulfide inhibits ethylene-induced petiole abscission in tomato (*Solanum lycopersicum* L.). *Horticulture Research*, 7(1), 1–11. <https://doi.org/10.1038/s41438-019-0237-0>
- Liu, G., Yang, Y., Liu, W., Guo, X., Xue, J., Xie, R., Ming, B., Wang, K., Hou, P., & Li, S. (2020b). Leaf removal affects maize morphology and grain yield. *Agronomy*, 10(2), 269. <https://doi.org/10.3390/agronomy10020269>
- Liu, Y., Fan, L., & Li, J. (2022). Flavor and compositional analysis of macadamia nuts during long-term storage. *Journal of Food Processing and Preservation*, 46(5), e16540. <https://doi.org/10.1111/jfpp.16540>
- Marei, N., & Crane, J. C. (1971). Growth and respiratory response of fig (*Ficus carica* L. cv. Mission) fruits to ethylene. *Plant Physiology*, 48(3), 249–254. <https://doi.org/10.1104/pp.48.3.249>
- Meir, S., Philosoph-Hadas, S., Riov, J., Tucker, M. L., Patterson, S. E., & Roberts, J. A. (2019). Re-evaluation of the ethylene-dependent and-independent pathways in the regulation of floral and organ abscission. *Journal of Experimental Botany*, 70(5), 1461–1467. <https://doi.org/10.1093/jxb/erz038>
- Nagao, M. A., Hirae, H. H., & Stephenson, R. A. (1992). Macadamia: Cultivation and physiology. *Critical Reviews in Plant Sciences*, 10(5), 441–470. <https://doi.org/10.1080/07352689209382321>
- Nagao, M. A., & Sakai, W. S. (1987). Effect of ethephon on abscission of macadamia fruits. In: Proceedings 14th Annual Meeting Plant Growth Regulator Society of America, Honolulu, HI, Aug. 2–6, 1987, pp. 70.
- Nagao, M. A., & Sakai, W. S. (1988). Influence of nut age on ethephon-induced abscission of macadamia. *Scientia Horticulturae*, 36(1–2), 103–108. [https://doi.org/10.1016/0304-4238\(88\)90012-X](https://doi.org/10.1016/0304-4238(88)90012-X)
- Navarro, S. L., & Rodrigues, C. E. (2016). Macadamia oil extraction methods and uses for the defatted meal byproduct. *Trends in Food Science & Technology*, 54, 148–154. <https://doi.org/10.1016/j.tifs.2016.04.001>
- Nortjé, G. P. (2018). Orchard management and fertilisation of macadamias. *FarmBiz*, 4(1), 28–31.
- O'Connor, K., Kilian, A., Hayes, B., Hardner, C., Nock, C., Baten, A., Alam, M., & Topp, B. (2019). Population structure, genetic diversity and linkage disequilibrium in a macadamia breeding population using SNP and silicoDarT markers. *Tree Genetics & Genomes*, 15(2), 1–16. <https://doi.org/10.1007/s11295-019-1331-z>
- Omidifard, Z., & Gharaghani, A. (2022). Chilling and heat requirements and bloom frost tolerance of *Prunus scoparia* (Spach) CK schneid and *P. elaeagnifolia* Spach. *Trees*, 36(5), 1559–1572. <https://doi.org/10.1007/s00468-022-02311-2>
- Patharkar, O. R., & Walker, J. C. (2018). Advances in abscission signaling. *Journal of Experimental Botany*, 69(4), 733–740. <https://doi.org/10.1093/jxb/erx256>
- Philips, A. L. (1970). Título en español. *Journal of Agriculture of the University of Puerto Rico*, 54(3), 504–507. <https://doi.org/10.46429/jaupr.v54i3.10986>
- Prasannath, K., Shivas, R. G., Galea, V. J., & Akinsanmi, O. A. (2021). Novel botrytis and cladosporium species associated with flower diseases of macadamia in Australia. *Journal of Fungi*, 7(11), 898. <https://doi.org/10.3390/jof7110898>
- Qiu, Z. L., Wen, Z., Yang, K., Tian, T., Qiao, G., Hong, Y., & Wen, X. P. (2020). Comparative proteomics profiling

- illuminates the fruitlet abscission mechanism of sweet cherry as induced by embryo abortion. *International Journal of Molecular Sciences*, 21(4), 1200. <https://doi.org/10.3390/ijms21041200>
- Quinlan, K., Treverrow, N., O'Hare, P., Slaughter, G., Mason, R., Wallace, H. M., & Walton, D. (2008). *Adoption of quality management systems in macadamia. Report series: MC03008*.
- Rahemi, M., & Ramezani, A. (2007). Potential of ethephon, NAA, NAD and urea for thinning pistachio fruitlets. *Scientia Horticulturae*, 111(2), 160–163. <https://doi.org/10.1016/j.scienta.2006.10.002>
- Rahman, A., Wang, S., Yan, J., & Xu, H. (2021). Intact macadamia nut quality assessment using near-infrared spectroscopy and multivariate analysis. *Journal of Food Composition & Analysis*, 102, 104033. <https://doi.org/10.1016/j.jfca.2021.104033>
- Rai, A. C., Halon, E., Zemach, H., Zviran, T., Sisai, I., Philosoph-Hadas, S., Meir, S., Cohen, Y., & Irihimovitch, V. (2021). Characterization of two ethephon-induced ida-like genes from mango and elucidation of their involvement in regulating organ abscission. *Genes*, 12(3), 439. <https://doi.org/10.3390/genes12030439>
- Reig, C., Martínez-Fuentes, A., Mesejo, C., Rodrigo, M. J., Zacarías, L., & Agustí, M. (2016). Loquat fruit lacks a ripening-associated autocatalytic rise in ethylene production. *Journal of Plant Growth Regulation*, 35(1), 232–244. <https://doi.org/10.1007/s00344-015-9528-3>
- Rengel, A., Pérez, E., Piombo, G., Ricci, J., Servent, A., Tapia, M. S., Gibert, O., & Montet, D. (2015). Lipid profile and antioxidant activity of macadamia nuts (*macadamia integrifolia*) cultivated in Venezuela. *Natural Science*, 07(12), 535–547. <https://doi.org/10.4236/ns.2015.712054>
- Richardson, A. C., & Dawson, T. E. (1993). Enhancing abscission of mature macadamia nuts with ethephon. *New Zealand Journal of Crop and Horticultural Science*, 21(4), 325–329. <https://doi.org/10.1080/01140671.1993.9513789>
- SAMAC. (2021). *Macadamia industry 2020*. Retrieved November 28, 2021, from <https://www.samac.org.za/industry-overview/>
- Schlörmann, W., Dinc, T., Lorkowski, S., & Gleis, M. (2017). Influence of roasting on the chemopreventive potential of macadamia nuts. *SDRP Journal of Food Science & Technology*, 2(1). <https://doi.org/10.25177/JFST.2.1.5>
- Shabalala, M., Toucher, M., & Clulow, A. (2022). The macadamia bloom—what are the hydrological implications? *Scientia Horticulturae*, 292, 110628. <https://doi.org/10.1016/j.scienta.2021.110628>
- Stephenson, R. A., Gallagher, E. C., & Doogan, V. J. (2003). Macadamia responses to mild water stress at different phenological stages. *Australian Journal of Agricultural Research*, 54(1), 67–75. <https://doi.org/10.1071/AR02108>
- Torres, E., & Asín, L. (2022). Physiological study of ethephon- and ACC-Induced fruitlet and leaf abscission in peach trees under different conditions of temperature. *Journal of Plant Growth Regulation*, 42(7), 1–10. <https://doi.org/10.1007/s00344-022-10888-8>
- Torres, E., Giné-Bordonaba, J., & Asín, L. (2021). Thinning flat peaches with ethephon and its effect on endogenous ethylene production and fruit quality. *Scientia Horticulturae*, 278, 109872. <https://doi.org/10.1016/j.scienta.2020.109872>
- Trueman, S. J. (2003). Yield responses to ethephon for unshaken and mechanically shaken macadamia. *Australian Journal of Experimental Agriculture*, 43(9), 1143–1150. <https://doi.org/10.1071/EA02048>
- Trueman, S. J. (2010). Benzyladenine delays immature fruit abscission but does not affect final fruit set or kernel size of macadamia. *African Journal of Agricultural Research*, 5(12), 1523–1530.
- Trueman, S. J. (2013). The reproductive biology of macadamia. *Scientia Horticulturae*, 150, 354–359. <https://doi.org/10.1016/j.scienta.2012.11.032>
- Trueman, S. J., McConchie, C., & Turnbull, C. (2002). Ethephon promotion of crop abscission for unshaken and mechanically shaken macadamia. *Australian Journal of Experimental Agriculture*, 42(7), 1001–1008. <https://doi.org/10.1071/EA01164>
- Trueman, S. J., Richards, S., McConchie, C., & Turnbull, C. (2000). Relationships between kernel oil content, fruit removal force and abscission in macadamia. *Australian Journal of Experimental Agriculture*, 40(6), 859–866. <https://doi.org/10.1071/EA00004>
- Wall, M. M. (2013). Improving the quality and safety of macadamia nuts. In *Improving the safety and quality of nuts*. Harris, Linda J. (ed.) (pp. 274–296). Woodhead Publishing.
- Wall, M. M., & Gentry, T. S. (2007). Carbohydrate composition and color development during drying and roasting of macadamia nuts (*macadamia integrifolia*). *LWT-Food Science and Technology*, 40(4), 587–593.
- Walters, K. J., & Lopez, R. G. (2018). Ethephon foliar sprays are influenced by carrier water alkalinity and ambient air temperature at application. *Horticulture Science*, 53(12), 1835–1841. <https://doi.org/10.21273/HORTSCI13426-18>
- Walton, D. A., & Wallace, H. M. (2009). Delayed harvest reduces quality of raw and roasted macadamia kernels. *Journal of the Science of Food & Agriculture*, 89(2), 221–226. <https://doi.org/10.1002/jsfa.3429>
- Walton, D., & Wallace, H. (2008). Review of quality management in the macadamia industry. Adoption of quality management systems in macadamia. Report series: MC03008, 11.
- Wang, J., Liang, S., Ma, H., Zhang, P., & Shi, W. (2016). Effects of ethephon on fresh in-husk walnut preservation and its possible relationship with phenol metabolism. *Journal of Food Science*, 81(8), C1921–C1927. <https://doi.org/10.1111/1750-3841.13370>
- Wang, Y., Yao, X., Yang, L., Fei, X., Cao, Y., Wang, K., & Guo, S. (2021). Effects of harvest time on the yield, quality and active substance of *Torreya grandis* nut and its oil. *Journal of Oleo Science*, 70(2), 175–184. <https://doi.org/10.5650/jos.ess20155>
- Yamada, Y., Koibuchi, M., Miyamoto, K., Ueda, J., & Uheda, E. (2015). Breakdown of middle lamella pectin by OH during rapid abscission in a zolla. *Plant, Cell & Environment*, 38(8), 1555–1564. <https://doi.org/10.1111/pce.12505>
- Yang, J., Liu, R. H., & Halim, L. (2009). Antioxidant and antiproliferative activities of common edible nut seeds. *LWT-Food Science and Technology*, 42(1), 1–8. <https://doi.org/10.1016/j.lwt.2008.07.007>
- Yuan, R. (2007). Effects of temperature on fruit thinning with ethephon in 'golden delicious' apples. *Scientia Horticulturae*, 113(1), 8–12. <https://doi.org/10.1016/j.scienta.2007.01.005>
- Zhang, Z., Chen, W., Tao, L., Wei, X., Gao, L., Gao, Y., Suo, J., Yu, W., Hu, Y., Yang, B., Jiang, H., Farag, M. A., Wu, J., & Song, L. (2023). Ethylene treatment promotes umami taste-active amino acids accumulation of *Torreya grandis* nuts postharvest by comparative chemical and transcript analyses. *Food Chemistry*, 408, 135214. <https://doi.org/10.1016/j.foodchem.2022.135214>



DSPACE

<https://dspace.org/>

Accelerating abscission of macadamia nuts using ethephon: are there implications for nut quality?

Aruwajoye, Noluthando Noxolo; Olarewaju, Olaoluwa Omoniyi; Oluwalana-Sanusi, Abimbola E.; Mditshwa, Asanda; Magwaza, Lembe Samukelo ...

2025-03

Taylor and Francis

<https://doi.org/10.1080/14620316.2024.2434614>

Aruwajoye NN, Olarewaju OO, Oluwalana-Sanusi AE, et al., (2025) Accelerating abscission of macadamia nuts using ethephon: are there implications for nut quality?. The Journal of Horticultural Science and Biotechnology, Volume 100, Issue 2, March 2025, pp. 153-163