

Accepted Manuscript

Title: Temperature and water activity effects on production of T-2 and HT-2 by *Fusarium langsethiae* strains from north European countries

Authors: A. Medina, N. Magan

PII: S0740-0020(10)00248-0

DOI: [10.1016/j.fm.2010.09.012](https://doi.org/10.1016/j.fm.2010.09.012)

Reference: YFMIC 1511

To appear in: *Food Microbiology*


Received Date: 22 July 2010

Revised Date: 27 September 2010

Accepted Date: 28 September 2010

Please cite this article as: Medina, A., Magan, N. Temperature and water activity effects on production of T-2 and HT-2 by *Fusarium langsethiae* strains from north European countries, *Food Microbiology* (2010), doi: 10.1016/j.fm.2010.09.012

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Temperature and water activity effects on production of T-2 and HT-2 by
Fusarium langsethiae strains from north European countries**

Medina, A.,* and Magan, N.

Applied Mycology Group, Cranfield Health, Vincent Building, Cranfield University, Cranfield,
Bedfordshire. MK43 0AL, UK.

* Corresponding author:

Ángel Medina-Vayá

Applied Mycology Group,

Cranfield Health, Vincent Building, Cranfield University,

Cranfield, Bedfordshire, MK43 0AL, U.K.

a.medinavaya@cranfield.ac.uk

Tel: 01234-758313

Fax: 01234-758380

Abstract

This study has examined the effect of ecophysiological factors, water activity (a_w , 0.995-0.90) and temperature (10-37°C), on the T-2 and HT-2 toxins production by *Fusarium langsethiae*. Two dimensional profiles for optimum and marginal conditions have been built for two strains from each of four northern European countries (UK, Norway, Sweden, Finland) on an oat-based medium. This showed that the optimum a_w and temperature conditions for T-2 + HT-2 production was between 0.98-0.995, and 20-30°C respectively. Kruskal-Wallis analysis of ranks showed a statistically significant differences between the different a_w levels examined ($P<0.001$) but no significant effect of the temperatures examined. The ratio of HT-2/T-2 was investigated and non-uniform distribution of HT-2 toxin was found under different ecological conditions. No statistically significant differences were found for the mean toxin production between strains from the different countries. Intra-strain differences in toxin production was only found for those from Finland (P -value=0.0247). The growth/no growth and toxin/no toxin conditions in relation to a_w x temperature have been constructed for the first time. This knowledge will be useful in developing prevention strategies to minimise T-2 and HT-2 toxin contamination by strains of *F. langsethiae* on important small grain cereals.

Key words: Type A trichothecenes, *Fusarium*, mycotoxins, ecological conditions

1. Introduction

Fusarium langsethiae has been isolated from infected oats, wheat and barley in central and northern Europe (Torp & Adler, 2004; Torp & Nirenberg, 2004). This species has been implicated in the production of high levels of T-2 and HT-2 mycotoxins in cereals in Norway (Langseth & Rundberget, 1999; Torp and Langseth, 1999) and in oats in the UK (Edwards, 2007). *F. langsethiae* has been isolated from infected symptomless oat and wheat grains which makes detection of contamination often very difficult. Its pathogenicity on these cereals has been recently demonstrated (Imathiu *et al.*, 2009).

T-2 and HT-2 toxins are type-A trichothecenes produced by different *Fusarium* species such as *F. acuminatum*, *F. sporotrichioides*, *F. poae*, and the recently described species *F. langsethiae* (Bottalico, 1998; Torp & Adler, 2004; Torp & Nirenberg, 2004). T-2 toxin is produced by *Fusarium* species and is rapidly metabolized to HT-2 toxin which is also the main metabolite *in vivo* (Eriksen & Alexander, 1998; Visconti, 2001). Studies on the metabolism of T-2 (Matsumoto *et al.*, 1978) suggested that the liver is the major organ for its metabolism, although other tissues are capable of metabolic modification of this toxin. Hepatic carboxylesterases have been shown to be responsible for the specific deacetylation of T-2, resulting in HT-2 as the major metabolite (Matsumoto *et al.*, 1978; Johnsen *et al.*, 1988). T-2 toxin, the most toxic Type A trichothecene, is a potent inhibitor of DNA, RNA, protein synthesis and mitochondrial function, and shows immunosuppressive and cytotoxic effects both *in vivo* and *in vitro* (Visconti *et al.*, 1991; Canady *et al.*, 2001; Visconti, 2001). A recent survey conducted in order to evaluate the risk of dietary exposure to *Fusarium* toxins by the population of EU member states, showed that T-2 and HT-2 toxins are quite common contaminants in cereals in the EU (Schothorst & van Egmond, 2004).

This resulted in special attention being paid to the toxic effects of T-2 in the Joint FAO/WHO Expert Committee on Food Additives (JECFA) where the safety of certain mycotoxins in food was evaluated (WHO/FAO, 2001). JECFA concluded that the toxic effects of

T-2 and its metabolite HT-2 could not be differentiated, and that the *in vivo* toxicity of T-2 might be due partly to toxic effects of HT-2. Therefore, the provisional maximum tolerable daily intake (PMTDI) for these toxins was fixed at 60 ng/kg body weight per day, including intake of T-2 and HT-2, alone or in combination (WHO/FAO, 2001). Recently, the European Food Safety Authority (EFSA) published a report on the toxicity of these trichothecenes and they concluded that the toxicity of T-2 toxin *in vivo* is considered to include that of HT-2 toxin and the results of studies with T-2 toxin are used to approximate the effects of HT-2 toxin (Schuhmacher-Wolz et al., 2010). The European Commission (EC) has established, with Regulations No. 856/2005 and No. 1881/2006, admissible levels of several *Fusarium* toxins in cereals and cereal-based products which became effective from 1 July, 2006. Maximum admissible levels for T-2 and HT-2 toxins in unprocessed cereals and cereal products are currently under discussion (Commission Regulation (EC) No 856/2005; Commission Regulation (EC) No 1881/2006).

It is now accepted that mycotoxin production is predominantly dependent on nutritional and ecological factors. Of the abiotic ecological factors, the water availability (water activity; a_w) and temperature are important factors which impact on both growth and mycotoxin production (Magan & Lacey, 1984; Sanchis & Magan, 2004; Magan & Aldred, 2007; Magan et al., 2010). While information is available on the relationship between these factors and profiles for growth and deoxynivalenol production by *F. culmorum* and *F. graminearum*, there is practically no information for *F. langsethiae* (Hope et al., 2005). As this species has become very important in northern Europe in a range of small grains it is important to understand the ecology of this species and whether any intra- or inter-strain differences may exist. Recently, we reported on the effect of a_w x temperature on growth of 8 strains, two each from the U.K., Norway, Sweden and Finland on an oat-based medium (Medina & Magan, 2010). This showed that there were no statistical differences in terms of a_w and temperature tolerances of the strains in terms of growth. The a_w and temperature optima were 0.98-0.995

and 25°C and growth limits were established at 0.92-0.93 a_w and 37°C and 5°C, respectively. There is no comparable information on how these interacting factors may affect T-2 and HT-2 production and whether intra- or inter-strain differences exist. This is critical in developing models to predict environmental conditions which represent a high risk and those that represent a low risk for contamination with these two mycotoxins.

The objectives of this study were thus to (a) determine the effect of a_w x temperature interactions on T-2, HT-2 and totals for two strains from four different northern European countries (U.K., Norway, Sweden, Finland) on an oat-based medium, (b) to evaluate any intra- or inter-strain differences and (c) develop contour maps of the optimum and marginal conditions for the production of these two mycotoxins and compare these with limits for growth reported recently (Medina & Magan, 2010).

2. Materials and methods

2.1. Strains.

Eight *F. langsethiae* strains from different northern European countries were examined. The isolates were from the UK (2004/57, 2004/59); Norway (44P, 88E); Sweden (560, 562) and Finland (05010, 05014).

2.2. Medium preparation and fungal culture.

Milled oats were prepared by homogenisation for 5 mins in a Waring laboratory science homogeniser model 7009G (Waring Laboratory Science, CT, USA). Mixtures of 2% (w/v) oat flour in water were prepared and 2% (w/v) agar added. Water used to prepare the medium was modified with glycerol to the required water activity levels (a_w ; 0.995, 0.98, 0.95, 0.93, 0.90 and 0.88). The culture media were prepared by autoclaving for 20 minutes at 121°C. The medium was vigorously shaken and poured into 9 cm diameter Petri dishes.

The eight strains of *F. langsethiae* were inoculated using 7 day old cultures by taking agar discs (4 mm diameter) with a cork borer from the growing margin of the colonies and inoculating the treatment plates centrally. The a_w treatments were incubated at different temperatures (35, 30, 25, 20, 15 and 10 °C) for 10 days. All experiences were carried out with three replicates per treatments, and in some cases repeated twice.

2.3.Reagents and standards.

Trichothecene standards, including T-2 and HT-2 were supplied by Sigma (Sigma-Aldrich, UK). T-2 standard was dissolved in acetonitrile at a concentration of 2.0 mg mL⁻¹ and stored at -26 °C in a sealed vial until use. HT-2 standard solution in acetonitrile was purchased at a concentration of 100.2 µg mL⁻¹. Working standards (50, 20, 10, 5, 2.5, 1, 0.5 and 0.2 µg mL⁻¹) were prepared by appropriate dilution of known volumes of the stock solutions with acetonitrile and used to obtain calibration curves for LC-DAD analysis. Acetonitrile and methanol were purchased from Fisher Scientific (Fisher Scientific UK Ltd., UK). All solvents were HPLC grade. Pure water was obtained from a Milli-R/Q water system (Millipore, Billerica, MA, USA) and used when water was required.

2.4.Sample preparation.

Using a cork borer, six-seven discs of agar weighing approx. 0.75 grams were removed from the fungal cultures and placed in previously weighed 2 mL volume safe-lock Eppendorf tubes. A total of 3 replicates per treatment were collected, weighed, and immediately frozen at -20°C and stored.

2.5.Extraction procedure.

Extraction was made using the methodology described by Medina et al. (2010). Briefly, samples were thawed and extracted by mixing the agar plugs with a one ml mixture of

methanol:water (80:20 v/v). The tubes were shaken at 150 rev min^{-1} at 25°C in the dark in an orbital shaker for 90 min. They were then centrifuged at 1150 g for 15 min and 750 μL of supernatant transferred to a 2 mL chromatography silanized amber vial. Extracts were completely dried in a stream of nitrogen. Dry extracts were redissolved in 300 μL of acetonitrile:water (50:50 v/v).

2.6.HPLC-DAD analysis.

The HPLC equipment consisted of an Agilent 1100 Series HPLC system equipped with a UV diode-array detector set at 200 nm (Agilent Technologies, Palo Alto, CA, USA). The column was a Phenomenex® Gemini C₁₈, 150 mm $\times$ 4.6 mm, 3 μm (Agilent Technologies, Palo Alto, CA, USA) preceded by a Phenomenex® Gemini C₁₈ 3 mm, 3 μm guard cartridge. Signals were processed by Agilent ChemStation software Ver. B Rev: 03.01 [317] (Agilent Technologies, Palo Alto, CA, USA).

Analyses were performed in the gradient mode. Solvent A was water and solvent B acetonitrile. Gradient conditions were initiated by holding for the first 3 minutes with 30% B, after this the conditions were changed linearly to 55% B for 18 min. The composition was then changed to 99% B in one minute and maintained for 5 min as a cleaning step in order to improve the results. After cleaning, conditions were returned to the initial 30% B. The flow rate of the mobile phase was 1 mL min^{-1} and injection volume was 50 μL .

2.7.Statistical analysis and profiling.

Statistical analysis was performed using the package JMP® 8 (SAS Institute Inc., 2008. Cary, NC, USA) and package STATISTICA 8 (StatSoft® Inc., 2007. Tulsa, OK, USA).

Data on toxin production were tested for normality using the Shapiro-Wilk test. Due to non-normality of the toxin concentration data, analysis was performed using non-parametric tests for testing whether distributions across factor levels were centered at the same location.

Differences between independent groups, using temperature and water activity as factors, were examined by the Kruskal-Wallis analysis of ranks. Analysis was performed for all data sets, including the 8 strains, and also for each strain individually. In order to compare strains from the same origin, t-tests were used.

Profiling graphs were performed using Sigma Plot v.10.0 (Systat Software Inc. Hounslow, London, UK).

3. Results

3.1. Effects of temperature and water activity on T-2 and HT-2 production.

Effect of a_w in the production of both mycotoxins individually is shown in Figure 1. This shows that reducing a_w from 0.995 to 0.95 produced a decrease in the mean production of both T-2 and HT-2 by a factor of approx. 17 and 20 respectively. No toxin production was found when the a_w treatment was <0.93 . The effect of the a_w was studied for the whole data set by Kruskal-Wallis analysis of ranks and this showed a statistically significant differences between the different a_w levels examined ($p < 0.001$). Statistical analysis of each strain gave the same results, showing the influence of water activity on all strains examined. In this case the p-values ranged from 0.0014 to 0.0078.

An example of the effect of temperature on T-2 and HT-2 production is shown in Figure 2. There was an up to 50% reduction in production of both the toxins at 10°C when compared with that at 20°C and 0.995 a_w . The Swedish strains appeared to be somewhat different as HT-2 production was higher at 10 than 20°C. The effect of the temperature was studied for the whole data set, including the 8 strains, by Kruskal-Wallis analysis of ranks and no significant differences were found. Strain by strain analysis did not show any significance of temperature as a factor with all the p-values >0.6078 . Despite this lack of statistical differences, there was a general trend in the behavior of *F. langesthiae* strains. Thus beyond the maximum production point a decrease of temperature resulted in a decrease in toxin production.

Response surfaces representing mean total mycotoxin (T-2 + HT-2) production for two strains in relation to a_w (from 0.93 to 0.995) x temperature (10-30°C) are shown for each country in Figure 3. Generally, the means of the two English and Finnish strains showed very similar contour maps for production which represent conditions of similar production levels. Maximum production were reached when the temperature was 25°C and the a_w was 0.98, and when the temperature was 20°C and the a_w 0.995 respectively. Also, in both cases, high mycotoxin levels were observed at 30°C at a slightly reduced a_w of 0.98.

Swedish strains appeared to produce higher concentrations of both T-2 and HT-2. In this case maximum production occurred when a_w and temperatures were 0.995 and 25°C respectively. Furthermore an increase was observed at 30°C at 0.98 a_w . The Norwegian strains produced the lowest concentrations of both mycotoxins with maximum production at 20°C and 0.98 a_w .

The representation of growth/no growth and toxin/no toxin (T-2 + HT-2) boundaries of *F. langsethiae* for different water availabilities and temperatures are shown in Figure 4. This demonstrates that toxin production by strains of *F. langsethiae* occurred over a narrower range of environmental conditions. While growth occurred at 0.90 a_w under the best temperature conditions, toxin production was only possible at 0.93 a_w . At marginal temperatures the difference between growth and toxin production was reduced.

3.2.Examination of HT-2/T-2 ratios.

The ratio of HT-2 against T-2 toxin in each treatment condition was calculated using the formula:

$$R = \text{HT-2} / \text{T-2}$$

Results obtained showed a non-uniform distribution of HT-2 toxin at all the ecological conditions. Overall, a substantial increase in R was observed in all strains studied under different ecological conditions.

In experiments where the temperatures were higher (20, 25 and 30°C) the R increased when a_w was decreased to 0.95 or 0.93. When temperatures were lower (10 and 15°C) the R was higher when a_w levels were 0.98 and 0.995. An example of this behaviour is shown in Figure 5. Thus, under marginal conditions HT-2 toxin increased in *F. langsethiae* strains although growth rates were reduced by between 0.5 and 2.5 mm/day.

3.3. Intra-strain differences based on source of strains.

Homoscedasticity was checked using the Levene's and Brown-Forsythe's tests. Since variances were not equal in some pairs, two different t-tests, one assuming equal and the other unequal variances, were applied where appropriate to find possible differences in T-2 and HT-2 production in strains from the same country of origin.

Significant differences regarding toxin production were found between strains from Finland (P -value=0.0247). No differences were observed between the strains from the other three countries.

3.4. Inter-strain differences between countries of origin.

Kruskal-Wallis analysis of ranks using the country of origin as a factor was applied to the data in order to find possible differences among countries. Since the p -value was 0.3487, no significant differences were found regarding the mean toxin production between strains from different countries of origin.

4. Discussion

This is the first study to examine the effect of ecological factors on the production of T-2 and HT-2 and the T-2/HT-2 ratio by *F. langsethiae* strains from a range of countries. This has provided new data which suggests that the variation between strains may be quite small, although there are some exceptions.

Changes in water availability appear to be a major controlling factor in affecting T-2 and HT-2 production. Generally maximum toxin production occurred where mycelia growth was under either a slight a_w or temperature stress. These results agree with data from other studies which have compared growth and toxin production by *Fusarium* species (Magan & Lacey, 1984; Magan et al., 2002; Sanchis & Magan, 2004).

Overall strains from England, Finland and Norway exhibited similar tolerances over a_w conditions of 0.995-0.93 for toxin production. Swedish strains showed a narrower toxin production window with less produced at 0.98 a_w , and practically almost none produced at 0.95 a_w . Evidences of genetic variability has been reported in this species (Yli-Mattila et al., 2004), supporting the possible existence of variability in ecophysiological performance at an intraspecific level.

Interestingly, for changes in temperature, no statistically significant differences were found. These results showed that *F. langsethiae* strains can produce T-2 and HT-2 toxins over a wide temperature range of 10 to 30°C, which were tested in this study. Some production may occur at 5°C with much longer incubation times but this was not tested.

Our results show that the optimum a_w and temperature conditions for T-2 + HT-2 production was between 0.98-0.995, and 20-30°C respectively. The optimum conditions for Type A trichothecenes have been previously described in different substrates for related species such as *F. sporotrichioides* and *F. poae*. Other authors have studied the production of type A trichothecenes by these species and concluded that moderate rather than warm temperatures were optima for these toxins. Also the optimum production conditions varied depending on the substrate and toxic metabolite. As an example, *F. sporotrichioides*-infected maize, wheat and rice grains contained more type A trichothecenes when a_w was 0.99 and were incubated at 20°C (Miller, 1994; Mateo et al., 2002). Recently Kokkonen et al. (2010) have studied toxin production of *F. langsethiae* on a mixed cereal medium under 3 environmental regimes (0.996, 0.96 a_w) and described maximum production of T-2 and HT-2

at a_w 0.996 and 15°C. However, they did not include 0.98 a_w which we found to be optimum for production.

In their reports, JECFA and EFSA concluded that the toxicity of T-2 toxin *in vivo* is considered to include that of HT-2 toxin and the results of studies with T-2 toxin are used to approximate the effects of HT-2 toxin (WHO/FAO, 2001; Schuhmacher-Wolz et al., 2010). Because of this, there has been interest in the total sum of both toxins. However, we have also examined the ratio of these two toxins. This suggested that this ratio may change as water stress conditions are imposed. Thus, we observed that more HT-2 toxin was produced when *F. langsethiae* was under intermediate stress conditions, where some reduction in growth occurs, but represent conditions which still allow toxin production. More investigation is needed to evaluate whether HT-2 toxin is directly produced by the strains, or, whether under ecological stress the fungus itself may degrade T-2, the main toxin produced, to HT-2. There are no studies of the metabolic pathways of HT-2 in *F. langsethiae* but for *F. sporotrichioides*, a close related species, a clear pathway has been proposed for the production of T-2 and also HT-2 (Meek et al., 2003). On the other hand, degradation of T-2 to HT-2 has been described in *F. graminearum*, *F. nivale*, *F. solani*, *F. sporotrichioides* and *Calonectria nivalis* (Vlastimil et al., 2008) suggesting the possibility of similar metabolic pathways in *F. langsethiae*.

The data obtained in this study on an oat-based nutritional matrix show that *F. langsethiae* produces mainly T-2 toxin. This finding is in contrast to other studies that describe a higher amount of natural samples contaminated with HT-2 and also higher concentrations (Langseth & Runberget, 1999; Edwards, 2009). However, these studies never examined the effect of water or temperature stress.

In oat-based culture medium HT-2 was observed in small quantities. Our findings suggest that the predominant toxin produced may be T-2 under most environmental conditions. The amount of HT-2 could increase because of the imposed ecological stress increasing the HT-2/T-2 ratio. Recent data supports our findings describing a switch from T-2 to

HT-2 as the major toxin produced depending on the environmental conditions (Kokkonen *et al.*, 2010). Partially, this may be due to T-2 being rapidly transformed to HT-2 by other microorganisms, by the fungus itself or by the cereal. This possibility has been suggested by Lattanzio *et al.* (2009). These authors studied the natural bio-transformation of T-2 into HT-2. Forty two per cent of the initial T-2 level was transformed naturally by the effect of carboxylesterase (CXE) enzymatic activity after 120 minutes. Different transformation speeds have been observed in different cereals; pointing out that the difference between them could be the amount of CXE or the expression of isoenzymes having different affinities for T-2. This enzymatic activity is also present in animal liver and is responsible for degradation *in vivo* (Matsumoto *et al.*, 1978). This enzymatic effect could explain why in natural cereal samples the toxin that is generally found is HT-2 instead of T-2. These differences regarding enzymatic activities could be on the basis of differences that have been suggested in some studies pointing out higher contamination with *F. langsethiae* and T-2/HT-2 toxin in conventional than organically produced oats (Edwards, 2009).

Differences in toxin contamination of cereals depending on the country of origin have also been found. Scudamore *et al.* (2009) in a recent four year study of contamination of oats with T-2/HT-2 found highest contamination in samples from the UK and Ireland while levels from Scandinavia were usually lower. No differences were found regarding the toxin production ability of strains of different origin. This supports the findings in the present study that there is very little difference in ecology and toxin production by *F. langsethiae* strains regardless of country of origin. This needs to be further tested by examining strains which have also been found in Germany and France.

Recently Parikka *et al.* (2007) found differences on the percentage of infected kernels depending on the cultivation practices. In this case Finnish cereals cultivated using direct drilling had higher *F. langsethiae* contamination while tilled plots were less contaminated.

Combining all this it is clear that different mycotoxin levels in grains would depend on different infection rates by *F. langsethiae* in the plots and prevailing weather conditions.

In summary, the present study has detailed, for the first time, information on the influence of interacting ecophysiological factors on T-2 and HT-2 toxin by strains of *F. langsethiae*. Water availability appears to be very important in determining contamination levels with these toxins. In contrast, they are produced over a wide temperature range. Often oats are harvested late in the season when conditions are wet. Thus inefficient drying may allow *F. langsethiae* to continue to colonise and increase contamination post-harvest. The contour maps of the growth/no growth and toxin/no toxin boundaries are useful for determining whether a high risk of contamination might occur post-harvest, and perhaps also in the field during ripening, in small grain cereals generally, especially oats. Thus, this study provides useful base line data on the conditions which represent a high and low risk for contamination by these mycotoxins which are becoming of growing importance in Europe.

5. Acknowledgements

We are very grateful to Prof. S. Edwards, Dr. M. Torp, Dr. M. Olsen and Dr. M. Jestoi for the supply of the strains, and to Balwinder Cook for her technical assistance with HPLC analysis. A. Medina wants to acknowledge funding from the Spanish Ministry of Science and Innovation (Programa Nacional de Movilidad de Recursos Humanos del Plan Nacional de I+D+I 2008-2011). Authors are also very grateful with Dr. M. Solfrizzo for his valuable critical discussions about T-2 and HT-2 chemistry.

6. References

- Bottalico, A. (1998). *Fusarium* diseases of cereals: Species complex and related mycotoxin profiles in Europe. *Journal of Plant Pathology*, 80, 85-103.
- Canady, R. A., Coker, R. D., Egan, S. K., Krska, R., Olsen, M., Resnik, S. & Schlatter, J. (2001). T-2 and HT-2 toxins. In: WHO (Ed.), Safety Evaluation of Certain Mycotoxins in Food, FAO Food and Nutrition Paper 74, WHO Food Additives Series 47, Geneva, Switzerland, pp. 557-680.
- Commission Regulation (EC) No 856/2005 of 6 June, 2005 amending Regulation (EC) No 466/2001 as regards *Fusarium* toxin. *Off J Eur Union* **L143** 3.
- Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Off J Eur Union* **L364** 5.
- Edwards, S. G. (2007). Investigation of *Fusarium* mycotoxins in UK barley and oat production. In *HGCA Project Report No. 415*. London: HGCA.
- Edwards, S. G. (2009). *Fusarium* mycotoxin content of UK organic and conventional oats. *Food Additives and Contaminants A*, 26, 1063-1069.
- Eriksen, G. S. & Alexander, J. (eds.) (1998). *Fusarium* toxins in cereals - a risk assessment. Nordic Council of Ministers; Tema Nord 1998, 502, Copenhagen pp. 7-44.
- Johnsen, H., Odden, E., Johnsen, B. A., Beyum, A. & Amundsen, E. (1988). Cytotoxicity and effects of T2-toxin on plasma proteins involved in coagulation, fibrinolysis and kallikrein-kinin system. *Archives of Toxicology*, 61, 237-240.R.
- Hope, R., Aldred, D. & Magan, N. (2005). Comparison of environmental profiles for growth and deoxynivalenol production by *Fusarium culmorum* and *F. graminearum* on wheat grain. *Letters in Applied Microbiology*, 40, 295-300.
- Imathiu, S. M., Ray, R. V., Back, M., Hare, M. C., & Edwards, S. G. (2009). *Fusarium langsethiae* pathogenicity and aggressiveness towards oats and wheat in wounded and unwounded in vitro detached leaf assays. *European Journal of Plant Pathology*, 124, 117-126.

- 367 Kokkonen, M., Ojala, L., Parikka, P., Jestoi, M. (2010). Mycotoxin production of selected
 368 *Fusarium* species at different culture conditions. *International Journal of Food Microbiology*,
 369 143, 17-25.
- 370 Langseth, W. & Rundberget, T. (1999). The occurrence of HT-2 toxin and other trichothecenes
 371 in Norwegian cereals. *Mycopathologia*, 147, 157-165.
- 372 Lattanzio, V. M. T., Solfrizzo, M. & Visconti, A. (2009). Enzymatic hydrolysis of T-2 toxin for the
 373 quantitative determination of total T-2 and HT-2 toxins in cereals. *Analytical and Bioanalytical*
 374 *Chemistry*, 395, 1325-1334.
- 375 Magan, N., Aldred, D., Mylona, K. & Lambert, R.J.W. (2010). Limiting mycotoxins in stored
 376 wheat. *Food Additives and Contaminants A*, 27, 644-650.
- 377 Magan, N., & Aldred, D. (2007). Post-harvest control strategies: Minimizing mycotoxins in the
 378 food chain. *International Journal of Food Microbiology*, 119, 131-139
- 379 Magan N & Lacey J (1984). Water relations of some *Fusarium* species from infected wheat ears
 380 and grain. *In: Transactions of the British Mycological Society* 83, pp. 281-285.
- 381 Magan, N. & Lacey, J. (1985). The effect of water activity and temperature on mycotoxin
 382 production by *Alternaria alternata* in culture and on wheat grain. *In: Lacey, J. (ed).*
 383 *Trichothecenes and other mycotoxins*. J. Wiley, Chichester, pp 243-256.
- 384 Magan, N., Hope, R., Colleate, A. & Baxter, E. S. (2002). Relationship between growth and
 385 mycotoxin production by *Fusarium* species, biocides and environment. *European Journal of*
 386 *Plant Pathology*, 108, 685-690.
- 387 Mateo, J.J., Mateo, R. & Jimenez, M. (2002). Accumulation of type A trichothecenes in maize,
 388 wheat and rice by *Fusarium sporotrichioides* isolates under diverse culture conditions.
 389 *International Journal of Food Microbiology*, 72, 115-123.
- 390 Matsumoto, H., Ito, T. & Ueno, Y. (1978). Toxicological approaches to the metabolites of
 391 fusaria. XII. Fate and distribution of T-2 toxin in mice. *Japanese Journal of Experimental*
 392 *Medicine*, 48, 393-399.

- 393 Medina, A., Valle-Algarra, F. M., Jimenez, M. & Magan, N. (2010). Different sample treatment
394 approaches for the analysis of T-2 and HT-2 toxins from oats-based media. *J Chromatograph B*,
395 878, 2145-2149.
- 396 Medina, A. & Magan, N. (2010). Comparisons of water activity and temperature impacts on
397 growth of *Fusarium langsethiae* strains from northern Europe on oat-based media.
398 *International Journal of Food Microbiology*, 142, 365-369.
- 399 Meek, B., Peplow, A. W., Ake, Jr. C., Phillips, T. D. & Beremand, M.N. (2003). Tri1 encodes the
400 cytochrome P450 monooxygenase for C-8 Hydroxylation during trichothecene biosynthesis in
401 *Fusarium sporotrichioides* and resides upstream of another new Tri gene. *Applied and*
402 *Environmental Microbiology*, 69, 1607-1613.
- 403 Miller, J.D. (1994). Epidemiology of *Fusarium* diseases of cereals. *In*: Miller, J.D. & Trenholm,
404 H.L. (eds) *Mycotoxins in Grain: Compounds other than Aflatoxins*. Eagon Press, St. Paul, MN,
405 USA, pp 19-36.
- 406 Parikka, P., Hietaniemi, V., Rämö, S. & Jalli, H. (2007). The effect of cultivation practices on
407 *Fusarium langsethiae* infection of oats and barley. *In*: Vogelgsang, S., Jalli, M., Kovács, G. &
408 Vida, G. (Eds). *Proceedings of the COST SUSVAR. Fusarium subgroup meeting: Fusarium*
409 *diseases in cereals-potential impact from sustainable cropping systems*. 1-2 June 2007.
410 Velence, Hungary
- 411 Sanchis, V. & Magan, N. (2004). Environmental profiles for growth and mycotoxin production.
412 *In*: Magan, N. & Olsen, M. (eds.). *Mycotoxins in Food: Detection and Control*. Cambridge:
413 Woodhead Publishing Ltd. pp. 174-189.
- 414 Schothorst, R. C., & van Egmond, H.P. (2004). Report from SCOOP task 3.2.10 "collection of
415 occurrence data of *Fusarium* toxins in food and assessment of dietary intake by the population
416 of EU member states" Subtask: trichothecenes. *Toxicology Letters*, 153, 133-143.

- Schuhmacher-Wolz, U., Heine, K. & Schneider, K. (2010). Report on toxicity data on trichothecene mycotoxins HT-2 and T-2 toxins. CT/EFSA/CONTAM/2010/03. Available online: http://www.efsa.europa.eu/en/scdocs/scdoc/65e.htm?WT.mc_id=EFSAHL01&emt=1
- Scudamore, K.A., Patel, S. & Edwards, S. G. (2009). HT-2 toxin and T-2 toxin in commercial cereal processing in the United Kingdom, 2004-2007. *World Mycotoxin Journal* 2, 357-365.
- Torp, M. & Adler, A. (2004). The European Sporotrichiella 309 Project: A polyphasic approach to the biology of new *Fusarium* species. *International Journal of Food Microbiology*, 95, 257-266.
- Torp, M. & Langseth, W. (1999). Production of T-2 toxin by a *Fusarium* resembling and *Fusarium poae*. *Mycopathologia*, 147, 89-96.
- Torp, M. & Nirenberg, H. I. (2004). *Fusarium langsethiae* sp. nov. on cereals in Europe. *International Journal of Food Microbiology*, 95, 247-256.
- Visconti, A., Minervini, F., Lucivero, G. & Gambatesa, V. (1991). Cytotoxic and immunotoxic effects of *Fusarium* mycotoxins using a rapid colorimetric bioassay. *Mycopathologia*, 113, 181-186.
- Visconti, A. (2001). Problem associated with *Fusarium* mycotoxins in cereals. *Bulletin of the Institute for Comprehensive Agricultural Sciences, Kinki University*, 9, 39-55.
- Vlastimil, D., Jezkova, A., Daniel, J. & Kamil, K. (2008). Metabolic pathways of T-2 toxin. *Current Drug Metabolism*, 9, 77-82.
- WHO/FAO (2001) Safety evaluation of certain mycotoxins in food. Prepared by the fifty-sixth meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA). WHO food additives series 47. FAO food and nutrition paper 74. International Programme on Chemical Safety, World Health Organization, Geneva
- Yli-Mattila, T., Mach, R.L., Alekhina, I.A., Bulat, S.A. & Koskinen, S. (2004). Phylogenetic relationship of *Fusarium langsethiae* to *Fusarium poae* and *Fusarium sporotrichioides* as

442 inferred by IGS, ITS, beta-tubulin sequences and UP-PCR hybridization analysis. *International*
443 *Journal of Food Microbiology*, 95, 267-285.

FIGURE CAPTIONS.

Figure 1. Effect of two water activity conditions (a_w ; 0.995 and 0.95) on the production of T-2 and HT-2 toxins by *F. langsethiae* strains after 10-days of incubation at 25 °C on an agar oat-based medium.

Figure 2. Effect of temperature (20 and 10°C) on the production of T-2 and HT-2 toxins in *F. langsethiae* strains after 10-days incubation at 0.995 a_w on an agar oat-based medium.

Figure 3. Two dimensional contour maps of T-2 + HT-2 toxin production profiles of *F. langsethiae* from different countries of origin (A: England, B: Finland, C: Norway and D: Sweden) in relation to temperature and water activity. The numbers on the isopleths are for the same mg of toxin/Kg of agar.

Figure 4. Two dimensional contour graph showing *F. langsethiae* mean boundary conditions for growth (growth 0.1 mm radius/day) vs. mean boundary conditions for T-2 + HT-2 production (0.5 mg of toxin/Kg). The graph has been obtained using average values obtained from all strains.

Figure 5. Effect of temperature (20 and 15°C) and a_w (0.995 and 0.98) on the HT-2/T-2 ratio (R) in *F. langsethiae* cultures after 10-days incubation on an agar oat-based medium.

Research highlights


Ms. Ref. No.: FM-D-10-00483


Title: Water availability and temperature affects production of T-2 and HT-2 by *Fusarium langsethiae* strains from north European countries.


Fusarium langsethiae is an important mycotoxigenic species and has become very important in northern Europe in a range of small grains that appeared to be contaminated with high amounts of these toxins. Thus it was very important to understand the ecology and how control of these toxins can be achieved.


- In the present study we have detailed, for the first time, information on the influence of interacting ecophysiological factors on T-2 and HT-2 toxin by strains of *F. langsethiae*.
 - Water availability appears to be very important in determining contamination levels with these toxins. In contrast, they are produced over a wide temperature range.
- Two dimensional profiles for optimum and marginal toxin production conditions have been built for two strains from each of four northern European countries (UK, Norway, Sweden and Finland) on an oat-based medium.
- It is the first time that, for this new species, contour map of the growth/no growth and toxin/no toxin boundaries have been built and will be published.
 - These maps will be useful for determining whether a high risk of contamination might occur post-harvest, and perhaps also in the field during ripening, in small grain cereals generally, especially oats.
- For the first time the ratio T-2/HT-2 ratio has been studied for *F. langsethiae*. We have described the effect that environmental conditions have on this ratio increasing the production of HT-2 under stress regimes.
 - This is the first time that the production of T-2 toxin as main toxins and the different possibilities for its transformation to HT-2 have been discussed.


Thus this study provides useful base line data on the conditions which represent a high and low risk for contamination by these mycotoxins which are becoming of growing importance in Europe. Also we provide interesting information that will enable further research to be carried out on this species.




























DSPACE

<https://dspace.org/>

Temperature and water activity effects on production of T-2 and HT-2 by *Fusarium langsethiae* strains from north European countries

Medina, Angel; Magan, Naresh

2011-05-06

Elsevier

<https://doi.org/10.1016/j.fm.2010.09.012>

A. Medina, N. Magan. Temperature and water activity effects on production of T-2 and HT-2 by *Fusarium langsethiae* strains from north European countries. *Food Microbiology*, Volume 28, Issue 3, May 2011, pp. 392-398