

Journal Pre-proof

Comparison of growth and aflatoxin B₁ production profiles of *Aspergillus flavus* strains on conventional and isogenic GM-maize-based nutritional matrices

Alessandra M. Gasperini, Angel Medina, Naresh Magan



PII: S1878-6146(21)00137-9

DOI: <https://doi.org/10.1016/j.funbio.2021.10.006>

Reference: FUNBIO 1280

To appear in: *Fungal Biology*

Received Date: 7 September 2021

Revised Date: 25 October 2021

Accepted Date: 27 October 2021

Please cite this article as: Gasperini, A.M., Medina, A., Magan, N., Comparison of growth and aflatoxin B₁ production profiles of *Aspergillus flavus* strains on conventional and isogenic GM-maize-based nutritional matrices, *Fungal Biology* (2021), doi: <https://doi.org/10.1016/j.funbio.2021.10.006>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd on behalf of British Mycological Society.

**Comparison of growth and aflatoxin B₁ production profiles of *Aspergillus flavus* strains
on conventional and isogenic GM-maize-based nutritional matrices**

Alessandra M. Gasperini*, Angel Medina and Naresh Magan

*Applied Mycology Group, Environment and AgriFood Theme, Cranfield University,
Cranfield, Beds., MK43 AL5, UK.*

**Present address: Department of Food Technology and Science, Federal University of Santa
Maria, Roraima Avenue 1000, Camobi, Santa Maria, Postal Code 97105-900, RS, Brazil*

Corresponding author: Prof Naresh Magan, Applied Mycology Group, Environment and
AgriFood Theme, Cranfield University, Cranfield, Beds. MK430AL, U.K. E-mail:
n.magan@cranfield.ac.uk.

ABSTRACT

Maize grown in both North and South America are now predominantly genetically modified (GM) cultivars with some resistance to herbicide, pesticide, or both. There is little information on the relative colonisation and aflatoxin B₁ (AFB₁) production with maize meal-based nutritional matrices based on kernels of non-GM maize and isogenic GM-ones by strains of *Aspergillus flavus*. The objectives were to examine the effect of interacting conditions of temperature (25-35 °C) and water availability (0.99-0.90 water activity, a_w) on (a) mycelial growth, (b) AFB₁ production and (c) develop contour maps of optimum and marginal conditions of these parameters for four strains of *A. flavus* on three different non-GM and isogenic GM-maize based nutritional media. The growth of the four strains of *A. flavus* (three aflatoxigenic; one non-aflatoxigenic) was relatively similar in relation to the temperature $\times$ a_w conditions examined on both non-GM and GM-based matrices. Optimum growth overall was at 30-35 °C and 0.99 a_w for all four strains. Under water stress (0.90 a_w) growth was optimum at 35 °C. Statistically: non-GM, GM cultivars, temperature and a_w all significantly affected growth rates. For AFB₁ production, all single and interacting factors were statistically significant except for non-GM $\times$ GM cultivar. In conclusion, colonisation of GM- and non-GM nutritional sources was similar for the different *A. flavus* strains examined. The contour maps will be very useful for understanding the ecological niches for both toxigenic and non-toxigenic strains in the context of the competitive exclusion of those producing aflatoxins.

Keywords: genetically modified (GM), non-GM, nutritional media, temperature, water activity, growth, mycotoxin, contour maps

1. Introduction

Aspergillus flavus is an important opportunistic fungal pathogen able to infect developing maize cobs during silking, especially if the plants have been weakened by drought stress (Dolezal et al., 2014), or damaged by insects (Chulze, 2010). Once *A. flavus* becomes established in the ripening cobs, it produces aflatoxins (AFs). At harvest, the moisture content (m.c.) of the maize cobs can be 18-20% which allows continued colonisation by *A. flavus* unless they are effectively dried to around 15% m.c. promptly (Gasperini et al., 2021). Often a delay in drying will lead to increased post-harvest AFs contamination, especially in tropical and sub-tropical environments. Due to its xerophilic nature (Araújo et al., 2020), *A. flavus* and related species have less competition during this process and are able to aggressively dominate the maize cob ecosystem and cause characteristic storage rots and concomitant AFs contamination (Battilani et al., 2016; Bhatnagar et al., 2006a; Scheidegger and Payne, 2003). The prevailing environmental conditions play a critical role in disease development. Interactions between intrinsic and extrinsic factors have been shown to have a significant influence on growth and mycotoxin production by such mycotoxigenic fungi in stored commodities (Magan et al., 2004).

A. flavus is capable of surviving and overwintering in plant residues as mycelium or sclerotia (Battilani et al., 2012). Crop residue is consequently a primary source of the fungus to allow conidia to be transmitted by air or insects to serve as new inoculum for the host plants to initiate a new infection cycle (Abbas et al., 2009). Infection and subsequent colonization is most aggressive on maize plants that have been subjected to heat or drought stress (Scheidegger and Payne, 2003). *A. flavus* development is favoured by hot dry conditions. It is considered to have a relatively wide temperature tolerance range (19–35 °C) with about 28 °C being considered optimum for growth and 28–30 °C for AFs production in maize. This contrasts with some other commodities, e.g., pistachio nuts, where optimum

conditions for growth were around 35 °C and for AFB₁ production 30-35 °C (Baazeem et al., 2021). Some strains of *A. flavus* have been shown to grow in very dry environments (e.g. 0.73 a_w) and produce aflatoxins at $>0.85 a_w$ (Sanchis and Magan, 2004). However, several other authors demonstrated that *A. flavus* requires $>0.90 a_w$ for AFB₁ production and $>0.85 a_w$ for growth (Lv et al., 2019; Nesci et al., 2003; Northolt et al., 1977; Peromingo et al., 2016). However, none of these previous studies compared growth and toxin production on genetically modified (GM) or non-GM isogenic lines of maize as nutritional sources.

There has been interest in the development of improved resistance to pests which cause significant damage and allows *A. flavus* to enter as a secondary coloniser. There has also been interest in better resistance to herbicides to maximise yield (Munkvold, 2014). This has led to the development of hybrids with thicker pericarp have also been utilised to try and increase resistance to pests and fungal infection. This resulted in the introduction of genes from *Bacillus thuringiensis* (especially related to the production of *Bt* toxin) to make the maize plants more resistant to certain pest species (Qaim, 2014). Similarly, GM crops with herbicide-tolerance (HT) have been successfully developed and are now commonly used by farmers for weed control in both North and South America.

The development and use of GM-maize has resulted in lower production costs and increased farmer incomes (Brookes and Barfoot, 2015). This has resulted in GM cultivars such as Roundup Ready[®] (tolerance to the herbicide glyphosate; Monsanto, USA), LibertyLink[™] (Bayer CropScience, USA) which provide tolerance to both glyphosate and Glufosinate ammonium. Today, most of the maize cultivars have stacked technologies to provide insect resistance (IR) and herbicide tolerance (HT) at the same time. Some studies have suggested that GM-IR has reduced fungal infections, and in some cases a reduction in mycotoxin content (Alma et al., 2005; Schnepf et al., 1998; Windham et al., 1999; Wu, 2006). However, no detailed studies have been conducted to examine the relationship

between interacting abiotic factors such as water availability $\times$ temperature on growth and AFs production by strains of *A. flavus* especially in relation to the nutritional content of conventional and isogenic GM-cultivars.

The objectives of this study were to compare the effect of (a) interacting water activity (a_w) $\times$ temperature conditions on growth profiles and (b) on aflatoxin B₁ (AFB₁) production by three toxigenic and one non-toxigenic strains of *A. flavus* on maize-based media using milled maize kernels of three conventional (non-GM) and isogenic GM cultivars as nutritional sources. Those examined included three conventional cultivars, and their associated isogenic pesticide resistant, or herbicide resistant or pesticide + herbicide stacked resistance. In addition, contour maps were developed on the optimum and marginal conditions for growth and AFB₁ production by the toxigenic and non-toxigenic strains.

2. Materials and methods

2.1. Strains used in this study

The *A. flavus* strains were isolated from maize and their identity confirmed using morphological and molecular analyses, and based on their AFs production patterns (Gasperini, 2019). Thus, a total of four strains were examined. Three were selected for these studies based on their ability to produce AFB₁ (toxigenic strains: *AfLRG-a*, *AfASC-a*, *AfP30C-a*). A fourth non-toxigenic strain (*AfPHx*) was also included as it was shown in the screening phase to not produce any AFs.

2.2. Maize-based media used in these studies

Three cultivars of non-GM maize and their respective isogenic GM lines (Table 1) were used in these studies. Kernels of each maize cultivar (250 g) were milled in a laboratory stainless steel blender (Waring® Commercial, Stamford, USA) for 60 seconds. The blender

was carefully cleaned between the milling of each cultivar using isopropanol and dried before subsequent use. The milled maize of each cultivar was kept in separate sterile Duran bottles and kept closed. These were afterwards used to prepare the nutritional media. The maize meal agar (MMA) was prepared by mixing 3% milled maize and 2% technical agar number 3 (Oxoid, Basingstoke, UK) in water: glycerol mixtures to modify the a_w from 0.99 (control) to 0.95, 0.90. Glycerol was used as the non-ionic solute due to its stability over the experimental temperature range tested. The MMA was prepared using deionised water (15 Ω m) and autoclaved at 121 °C (103 kPa) for 15 min. The molten cooled media (50 °C) were shaken well before pouring into 9 cm Petri plates. The cooled a_w treatment media were stored separately in closed plastic bags at 4 °C until being used. The accuracy of the modifications in a_w was confirmed using an Aqualab 4TE instrument (Decagon, Pullman, USA) to be within $\pm 0.002 a_w$

2.3. Inoculation and mycelial growth measurements

Spore suspensions (10^6 spores mL^{-1}) of each of the *A. flavus* strains were spread-plated on MEA (Malt Extract Agar; Oxoid, Basingstoke, UK) and incubated at 25 °C for 24 h to allow germination to occur (Gasperini, 2019). These Petri plate colonies were used to provide the inoculum for the experiments. Agar plugs (3 mm $\varnothing$) were taken from these inoculation plates with a surface sterilised cork-borer and transferred to the centre of the treatment and replicate MMA plates (9 cm $\varnothing$). The replicate plates for each a_w level was enclosed in separate polyethylene sealed boxes with a glycerol/water solution of the same a_w as the plates to keep the equilibrium relative humidity at the target abiotic conditions. The boxes were incubated at 25, 30 and 35 °C for 10 days.

The diameters of the colonies were measured in two perpendicular directions at right angles to each other daily for each maize treatment and replicate for each *A. flavus* strain.

The growth rate of each strain under the set of nutritional treatments at the different $a_w \times$ temperature conditions were calculated by plotting the temporal diametric growth rate daily. The regression lines of the linear growth phase were used for each replicate to obtain the relative growth rate (mm/day) for each strain and under the different interacting abiotic conditions for each maize nutritional medium using Microsoft Excel®. Experiments were carried out with three replicates per treatment and the whole experiment carried out twice.

2.4. Quantification of aflatoxin B_1 production by the *A. flavus* strains on the different non-GM and GM-based nutritional media

Extraction of aflatoxins from cultures: After 10 days incubation, 6 agar plugs (approx. 0.3 g) were taken from the cultures across each plate with a 5 mm-diameter sterile cork-borer in a sterile flow bench. These plugs were placed into pre-weighed 2-mL safe-lock tubes and re-weighed. Aflatoxins were extracted from the agar plugs by adding 1 mL of chloroform and shaking for 1h at room temperature in the dark. An 800 μ L aliquot from the tube was transferred to a new tube and the solvent evaporated to dryness overnight. The derivatization was then performed as detailed below.

Derivatization of aflatoxins: Into the 2-ml safe-lock tube with the residue of each sample, 200 μ L of hexane were added, vortexed for 30 sec, followed by the addition of 50 μ L trifluoroacetic acid (TFA, Fisher Scientific, UK). The mixture was vortexed for 30 sec and left for 5 min. Thereafter, 950 μ L water: acetonitrile (9:1, v/v) was added and the entire contents of the tube vortexed for 30 sec. Then the mixture was left for 10 min to allow thorough separation of the layers. The hexane (upper layer) was discarded, and the aqueous portion filtered through Nylon filters (13 mm $\times$ 0.22 μ m) using a 1 mL syringe directly into amber

silanized 2-mL vials (Agilent Technologies, USA) for HPLC analysis. All the analytical reagents used were HPLC grade.

HPLC analysis: The quantification of the production of AFs by the four strains of *A. flavus* was performed by reverse phase HPLC. The HPLC system used was an Agilent 1200 series with fluorescence detector (λ_{exc} 360 nm; λ_{em} 440 nm). A C₁₈ column (Agilent Poroshell® 120 EC-18, 4.6 x 100 mm, 2.7 μ m particle size preceded by a Phenomenex® Gemini C18 column, 3 mm, 3 μ m guard cartridge; Phenomenex, CA, USA) was used for separation. Isocratic elution was made with methanol:water:acetonitrile (30:60:10, v/v/v) and a mobile phase flow rate of 1.0 mL/min. The injection volume was 10-50 μ L depending on each set of samples. The mixed aflatoxin standard solution used was obtained from Romer Labs U.K Ltd and comprised 0.2 μ L mL⁻¹ of aflatoxin B₁ (AFB₁) and aflatoxin G₁ (AFG₁), and 0.06 μ L mL⁻¹ of aflatoxin B₂ (AFB₂) and aflatoxin G₂ (AFG₂). This was diluted, left to evaporate to dryness overnight and, thereafter, derivatized as described previously. The set of standards injected was 0.05–4 ng of aflatoxins per injection, and the curves generated by plotting the peaks area against the amounts of aflatoxins. Only AFB₁ was detected in the samples from three of the strains. The correlation coefficient value for the calibration curve was $R^2 = 0.995$.

2.5. Statistical analysis

Statistical analysis was performed using the package JMP® Pro 14 (SAS Institute Inc., USA). Shapiro-Wilk tests was used to determine normality and Levene's test to assess the equality of variance. Both data from growth rates and AFB₁ production failed the requirements for analysis of variance therefore the nonparametric (Wilcoxon /Kruskal-Wallis; $p=0.05$) were used for analyses (Chan and Walmsley, 1997). Where there was significance after the Kruskal-Wallis test, median comparisons for each pair of the different

cultivars were made using the Wilcoxon – Each Pair test ($p=0.05$). Forward stepwise regression was used to obtain factorial to degree equations for AFB₁ (ng/g) for conditions of toxin production (a_w , T °C, $a_w \times T$ °C) for each strain and for the combined strains for each nutritional medium based on the maize cultivar. Contour maps were built in JMP® Pro 14 using 5000 simulation data from the predicted formula.

3. Results

3.1. Effects of maize cultivar nutritional source and environmental factors on growth of toxigenic and non-toxigenic *A. flavus* strains

The overall growth rate for all four strains (toxigenic: *AfLRG-a*, *AfASC-a* and *AfP30C-a*; non-toxigenic strain *AfPHx*) were optimum at 30-35 °C and 0.99 a_w on both non-GM and GM maize meal-based media (Figure 1). The slowest growth rate was observed at 0.90 a_w in all the temperatures tested. However, growth rates were higher at 35 °C and 0.90 a_w for the four strains regardless of whether grown on non-GM or GM-based maize media. Statistically, there was no significant difference in the growth rate at 30 and 35 °C at 0.95 and 0.99 a_w ($p = 0.05$). Overall, there was no significant difference ($p = 0.05$) in the colonisation rates of GM $\times$ non-GM nutritional media and between the *A. flavus* strains studied, although there was a statistically significant effect ($p = 0.05$) of cultivar on the growth of the *A. flavus* strains (Table 2). The detailed growth data for each strain on non-GM and GM nutritional media are shown in Suppl. Figure 1.

A contour map for interacting conditions of temperature $\times$ a_w for a toxigenic and non-toxigenic strain of *A. flavus* is provided in Figure 2. The data for the toxigenic strains are based on the pooled data for the growth on both non-GM and GM cultivars. The mycelial growth profiles for the toxigenic and non-toxigenic strains were similar. The optimal growth

conditions for both strains were at 0.99 a_w and ≥ 30 °C with a maximum of 14 mm and 12 mm day⁻¹ (strains AfLRG-a and AfPHX, respectively) over 10 days incubation.

3.2. Effect of non-GM and GM nutritional sources and environmental factors on aflatoxin B₁ production by the *A. flavus* strains

The overall AFB₁ production by the three toxigenic *A. flavus* strains was integrated across non-GM and GM cultivars as there was no significant difference ($p = 0.05$) between them. This shows that optimum AFB₁ production was at 0.99 a_w , especially at 30 and 35 °C on both non-GM and GM nutritional media for all three strains. The other strain did not produce any AFB₁ or indeed other AFs. It was noticeable that at 0.95 and 0.90 a_w the amounts of AFB₁ produced was at least ten times less than at 0.99 a_w , especially at 30 and 35 °C (Figure 3). Statistically, all single and interacting factors were significant, except for non-GM $\times$ GM cultivars (Table 2). The detailed data on production of AFB₁ on non-GM and GM nutritional media for each of the three strains is shown in Suppl. Figure 2.

The contour maps for the two toxigenic strains which produced the highest amounts of AFB₁ are shown in Figures 4 and 5 based on the different media made from non-GM and GM maize. The strain AfLRG-a was the highest AFB₁ producer amongst all the toxigenic strains when the cultivar P30F53 CON was used as a substrate (≥ 6000 ng g⁻¹; Figure 4A). Interestingly, the toxin production on the cultivar P2530 CON was about 60-fold lower (≥ 100 ng. g⁻¹; Figure 4C). In 5 out of 6 maize cultivars used, this strain had optimal conditions for AFB₁ production at 30 °C and >0.98 a_w . However, the opposite occurred when the strain was grown on media made with cultivar P2530 Hx[®], where more toxin was produced under stress conditions: <30 °C/0.90 a_w (800 ng g⁻¹; Figure 4F).

The toxigenic strain AfP30C-a also produced more AFB₁ when cultivated on the non-GM P30F53 CON maize meal as a substrate (≥ 1000 ng/g; Figure 5A). However, lower levels were synthesised by this species on the non-GM cultivar P2530 CON (≥ 80 ng g⁻¹; Figure 5C). The optimal conditions for this strain were also modified, when grown on the GM-maize cultivar. P2530 Hx[®], with more AFB₁ produced at 35°C/0.98 a_w (≥ 300 ng g⁻¹; Figure 5F). The contour maps for the other toxigenic strain is provided in Suppl. Figure 3.

4. Discussion

Colonisation of nutritional media based on GM or non-GM isogenic maize kernels under different a_w x temperature conditions has no effect on the relative growth rates by three toxigenic strains and a non-toxigenic strain of *A. flavus*. However, AFB₁ production was significantly different for the toxigenic strains. Data was utilised to develop contour maps of both the optimum and boundary conditions for colonisation and AFB₁ production by individual and pooled data for the three toxigenic strains and for the non-toxigenic strains on both GM and non-GM matrices.

The optimal conditions for growth and AFB₁ production found in this work were distinct with slightly different optima depending on the a_w × temperature conditions. Previously, Medina et al. (2014) also found differences in growth and toxin boundary conditions depending on a_w × temperature interactions. Some studies have also suggested that the biosynthesis of secondary metabolites can sometimes be increased under environmental stress conditions, especially related to water availability, and elevated temperatures. According to Bhatnagar et al. (2006, 2018), aflatoxin biosynthesis in *A. flavus* is optimal at temperatures between 29 and 30 °C, but it is significantly decreased at temperatures <25 and >37 °C. Unlike most fungi, *A. flavus* is favoured by hot dry conditions. Sanchis and Magan

(2004) described that some strains of *A. flavus* can grow in very dry environments (e.g., 0.73 a_w) and produce aflatoxins at $\geq 0.85 a_w$. These were based on long incubation periods of up to 25-30 days. However, other studies demonstrated that *A. flavus* required $>0.90 a_w$ for AFB₁ production and $>0.85 a_w$ for growth, dependent on the nutritional medium used (Lv et al., 2019; Nesci et al., 2003; Northolt et al., 1977; Peromingo et al., 2016). In addition, these abiotic interactions have been demonstrated to influence the biosynthetic gene cluster expression and thus AFB₁ biosynthesis (Abdel-Hadi et al., 2010; Medina et al., 2015).

It was noted that the interactions between environmental effects and the nutritional status of different maize cultivars influenced the patterns of AFB₁ production. The same strain produced different amounts of AFB₁ in each maize cultivar. This suggests that the cultivars may perhaps have slightly different micro-nutritional components. This could have influenced mycotoxin production, although no differences in colonisation were discernible.

The concentration of AFB₁ was not consistently higher in a specific type of maize (non-Gm vs GM) indicating that each cultivar itself results in more or less toxin despite the genetic modifications. For example, when the cultivar P30F53 CON was used, for example, the AFB₁ accumulation was significantly higher than its isogenic GM line for 2 strains (*Af*LRG-a and *Af*P30C-a). Nevertheless, the conditions for the maximum production were modified by the interactions between the abiotic factors and cultivar. Thus, according to the contour maps for the strain *Af*LRG-a: 0.90 a_w at 25 °C were best for AFB₁ production for the GM cultivar (P2530 Hx), whereas 0.99 a_w at 25 °C were optimum for the equivalent non-GM one (P2530 CON). One possibility for these differences could be the nutritional content of each hybrid that may have influenced the toxin amount produced if the metabolism of the *A. flavus* strain was influenced, despite there being no significant impacts on colonisation patterns.

Bakan et al. (2002) suggested that the biochemical composition of GM maize (MON 810 *Bt*) and the near isogenic non-GM line could be responsible for the differences in fumonisins content found. However, it was reported that the content of proteins, soluble nitrogen, starch and sugars did not differ between GM hybrids and their near-isogenic non-GM lines (Masoero et al., 1999; Betz et al., 2000). Therefore, from a macro-nutritional assessment, the GM maize cultivars may have many similar amounts to the parental maize line (Reuter et al., 2002).

Another possibility is that there is a link between fatty acid (FAs) content in the non-GM and GM cultivars which may influence the AFB₁ production by the different strains. Levels of free FAs increase in response to various stresses and can play a pivotal role in plant–microbe interactions (Walley et al., 2013). Seed FAs composition has also been suggested to be a component of pathogen susceptibility and seed colonisation (Dall’Asta et al., 2012). As free FAs, linoleic acid levels partly regulate development, seed colonisation and mycotoxin production by *Aspergillus* spp. (Calvo et al., 1999; Scarpari et al., 2014).

Studies by Jiménez et al. (2009) found FA profiles in the non-GM and GM maize with insect-resistance (*Cry1Ab* gene) cultivars to be relatively similar. (Frank et al., 2012) suggested that most differences between non-GM and GM maize cultivars with insect or herbicide resistance (DKC78-15B, TXP 138F, DKC78-35R, respectively) were due to abiotic and not genetic factors. Thus, location, weather patterns and season predominantly influenced the FA profiles. Battilani et al. (2018) have suggested that oxylipins may be important, both in terms of defence reactions to fungal infection and potentially as virulence factors for mycotoxigenic fungi such as *Fusarium verticillioides*. However, no studies have examined comparisons of oxylipins in non-GM and cultivars and their effect on *A. flavus* development.

Manetti et al. (2004, 2006) and Piccioni et al. (2009) tried to identify the metabolomic fingerprint of non-GM and GM maize seeds using Nuclear Magnetic Resonance spectroscopy

(NMR). NMR was thus used to profile the metabolome of insect-resistant GM maize containing *cry1A(b)* gene. Amongst the 40 water-soluble metabolites identified in all samples, those which could be used for discrimination between non-GM and GM cultivars were ethanol, citric acid, trehalose, and glycine-betaine. These were all present in higher levels in the GM maize samples (Piccioni et al., 2009). However, the relationship between maize cultivar modifications, the content of these metabolites and the genetic modification has not been elucidated (Simó et al., 2014). However, none of these studies examined the effect that modification of such compounds may have on colonisation by *A. flavus* or AFs contamination.

The growth of the non-aflatoxigenic strain suggests similar growth patterns to the toxigenic strains on both non-GM and GM-based media as a nutritional source. This may be important that the resilience of such strains over the range of interacting abiotic factors over which toxigenic strains can grow and produce AFs (Medina et al., 2017; Gasperini et al., 2019). It has also been shown that the utilisation patterns of key C-sources in maize vary between toxigenic and non-toxigenic species (Mohale et al., 2013; Medina et al., 2017). It is critical for effective control of both pathogens and potential toxin contamination that the ecological niche profile of the biocontrol strain is similar to that of the toxigenic strains (Magan, 2020). The present study suggests that in both non-GM and GM cultivars the non-toxigenic strain can grow effectively over the key two-dimensional abiotic ranges similar to the toxigenic strains.

Overall, the growth profiles for the toxigenic and non-toxigenic strains were similar, regardless of whether the nutritional sources were based on non-GM or GM isogenic maize cultivar kernels. The contour maps show this clearly in relation to optimum and marginal conditions for colonisation. In addition, this also suggests that biocontrol strains may be effective on both non-GM and GM cultivars in terms of ecological windows for competition

with toxigenic strains. In terms of AFB₁ production, there were differences in terms of the amounts of toxin produced and the optimum and marginal quantities produced after a 10-day period. This does suggest that there may be some influence of cultivar type on AFB₁ production, whether they are non-GM or GM-based. More detailed studies are required of the potential impact of individual or groups of macro- and micro-nutrients which may be present in these two groups maize cultivars. The composition of the maize hybrids may have a bigger impact on AFs contamination. This is important as many countries where GM maize cultivars are grown have maximum legislative limits for food and feed chains.

Acknowledgements

AMG is grateful to CAPES Foundation, Ministry of Education of Brazil for the financial support for her PhD (Project BEX 12937/13-4).

References

- Abbas, H., Wilkinson, J., Zablotowicz, R., Accinelli, C., Abel, C., Bruns, H., Weaver, M., 2009. Ecology of *Aspergillus flavus*, regulation of aflatoxin production, and management strategies to reduce aflatoxin contamination of corn. *Toxin Rev.* 28, 142–153.
- Abdel-Hadi, A., Carter, D., Magan N. 2010. Temporal monitoring of the *nor-1* (*aflD*) gene of *Aspergillus flavus* in relation to aflatoxin B₁ production during storage of peanuts under different environmental conditions. *J. Appl. Microbiol.* 109, 1914–1922.
- Alma, A., Lessio, F., Reyneri, A., Blandino, M., 2005. (*Lepidoptera: Crambidae*) feeding activity, crop technique and mycotoxin contamination of corn kernel in northwestern Italy. *Int. J. Pest Manag.* 51, 165–173. <https://doi.org/10.1080/09670870500179698>
- Araújo, C.A.S., Ferreira, P.C., Pupin, B., Dias, L.P., Avalos, J., Edwards, J., Hallsworth, J.E., Rangel, D.E.N., 2020. Osmotolerance as a determinant of microbial ecology: A study of phylogenetically diverse fungi. *Fungal Biol.* 124, 273–288. <https://doi.org/10.1016/j.funbio.2019.09.001>

- Baazeem, A., Garcia-Cela, E., Medina, A., Magan, N., 2021. Interacting abiotic factors affect growth and aflatoxin B₁ production profiles of *Aspergillus flavus* strains on pistachio-based matrices and pistachio nuts. *Front. Microbiol.* 11. <https://doi.org/10.3389/fmicb.2020.624007>
- Bakan, B., Melcion, D., Richard-Molard, D., Cahagnier, B., 2002. Fungal growth and *Fusarium* mycotoxin content in isogenic traditional maize and genetically modified maize grown in France and Spain. *J. Agric. Food Chem.* 50, 728–731. <https://doi.org/10.1021/jf0108258>
- Battilani, P., Lanubile, A., Scala, V., Reverberi, M., Gregori, R., Falavigna, C., Dall'asta, C., Park, Y.-S., Bennett, J., Borrego, E.J., Kolomiets, M. V., 2018. Oxylipins from both pathogen and host antagonize jasmonic acid-mediated defence via the 9-lipoxygenase pathway in *Fusarium verticillioides* infection of maize. *Mol. Plant Pathol.* 19, 2162–2176. <https://doi.org/10.1111/mpp.12690>
- Battilani, P., Rossi, V., Giorni, P., Pietri, A., Gualla, A., Fels-Klerx, H.J. van der, Booi, C.J.H., Moretti, A., Logrieco, A., Miglietta, F., Toscano, P., Miraglia, M., Santis, B.D., Brera, C., 2012. Modelling, predicting and mapping the emergence of aflatoxins in cereals in the EU due to climate change. *EFSA Support. Publ.* 9, 223E. <https://doi.org/10.2903/sp.efsa.2012.EN-223>
- Battilani, P., Toscano, P., Van der Fels-Klerx, H.J., Moretti, A., Camardo Leggieri, M., Brera, C., Rortais, A., Goumperis, T., Robinson, T., 2016. Aflatoxin B₁ contamination in maize in Europe increases due to climate change. *Sci. Rep.* 6, 24328. <https://doi.org/10.1038/srep24328>
- Betz, F.S., Hammond, B.G., Fuchs, R.L., 2000. Safety and advantages of *Bacillus thuringiensis*-protected plants to control insect pests. <https://doi.org/10.1006/rtph.2000.1426>
- Bhatnagar, D., Cary, J.W., Ehrlich, K., Yu, J., Cleveland, T.E., 2006b. Understanding the genetics of regulation of aflatoxin production and *Aspergillus flavus* development. *Mycopathologia* 162, 155–166. <https://doi.org/10.1007/s11046-006-0050-9>
- Bhatnagar, D., Rajasekaran, K., Gilbert, M., Cary, J.W., Magan, N., 2018. Advances in molecular and genomic research to safeguard food and feed supply from aflatoxin contamination. *World Mycotoxin J.* 11, 47–72. <https://doi.org/10.3920/WMJ2017.2283>

- Brookes, G., Barfoot, P., 2015. Global income and production impacts of using GM crop technology 1996–2013. *GM Crops Food Biotechnol. Agric. Food Chain* 6, 13–46. <https://doi.org/10.1080/21645698.2015.1022310>
- Calvo, A.M., Hinze, L.L., Gardner, H.W., Keller, N.P., 1999. Sporogenic effect of polyunsaturated fatty acids on development of *Aspergillus* spp. *Appl. Environ. Microbiol.* 65, 3668–73.
- Chan, Y., Walmsley, R.P., 1997. Learning and understanding the Kruskal-Wallis one-way analysis-of-variance-by-ranks test for differences among three or more independent groups. *Phys. Ther.* 77, 1755–62.
- Chulze, S.N., 2010. Strategies to reduce mycotoxin levels in maize during storage: a review. *Food Addit. Contam. Part A* 27, 651–657. <https://doi.org/10.1080/19440040903573032>
- Dall'Asta, C., Falavigna, C., Galaverna, G., Battilani, P., 2012. Role of maize hybrids and their chemical composition in *Fusarium* infection and fumonisin production. *J. Agric. Food Chem.* 60, 3800–3808. <https://doi.org/10.1021/jf300250z>
- Dolezal, A.L., Shu, X., O'Brian, G.R., Nielsen, D.M., Woloshuk, C.P., Boston, R.S., Payne, G.A., 2014. *Aspergillus flavus* infection induces transcriptional and physical changes in developing maize kernels. *Front. Microbiol.* 5, 1–10. <https://doi.org/10.3389/fmicb.2014.00384>
- Frank, T., Röhligh, R.M., Davies, H. V., Barros, E., Engel, K.-H., 2012. metabolite profiling of maize kernels—genetic modification versus environmental influence. *J. Agric. Food Chem.* 60, 3005–3012. <https://doi.org/10.1021/jf204167t>
- Gasperini, A.M., 2019. Fungal diversity, pest damage and biocontrol of aflatoxins in GM and conventional Brazilian maize cultivars under existing and future climate change scenarios (Ph.D Thesis). Cranfield University, England.
- Gasperini, A.M., Garcia-Cela, E., Sulyok, M., Medina, A., Magan, N., 2021. Fungal diversity and metabolomic profiles in GM and isogenic non-GM maize cultivars from Brazil. *Mycotoxin Res.* 37, 39–48. <https://doi.org/10.1007/s12550-020-00414-8>
- Gasperini, A.M., Rodriguez-Sixtos, A., Verheecke-Vaessen, C., Garcia-Cela, E., Medina, A., Magan, N., 2019. Resilience of biocontrol for aflatoxin minimization strategies: climate change abiotic factors may affect control in non-GM and GM-maize cultivars. *Front. Microbiol.* 10, 2525. <https://doi.org/10.3389/fmicb.2019.02525>
- Jiménez, J.J., Bernal, J.L., Nozal, M.J., Toribio, L., Bernal, J., 2009. Profile and relative concentrations of fatty acids in corn and soybean seeds from transgenic and isogenic

- crops. J. Chromatogr. A 1216, 7288–7295.
<https://doi.org/10.1016/J.CHROMA.2009.08.015>
- Lv, C., Jin, J., Wang, P., Dai, X., Liu, Y., Zheng, M., Xing, F., 2019. Interaction of water activity and temperature on the growth, gene expression and aflatoxin production by *Aspergillus flavus* on paddy and polished rice. Food Chem. 293, 472–478.
<https://doi.org/10.1016/j.foodchem.2019.05.009>
- Medina, A., Schmidt-Heydt, M., Rodríguez, A., Parra, R., Geisen, R., Magan, N. 2015. Impacts of environmental stress on growth, secondary metabolite biosynthetic gene clusters and metabolite production of xerotolerant/xerophilic fungi. Curr. Genet. 61, 325–334.
- Magan, N., 2020. Importance of Ecological windows for efficacy of biocontrol agents, in: De Cal, A., Melgarejo, P., Magan, N. (Eds.), How research can stimulate the development of commercial biological control against plant diseases, progress in biological control. Springer International Publishing, Cham, pp. 1–14.
https://doi.org/10.1007/978-3-030-53238-3_1
- Magan, N., Aldred, D., Sanchis, V., 2004. The role of spoilage fungi in seed deterioration, In: Dilip K. Arora (Ed.), Fungal Biotechnology in Agricultural, Food, and Environmental Applications. CRC Press, pp. 311–323.
- Manetti, C., Bianchetti, C., Bizzarri, M., Casciani, L., Castro, C., Dascenzo, G., Delfini, M., Dicocco, M., Lagana, A., Miccheli, A., Motto, M., Conti, F., 2004. NMR-based metabonomic study of transgenic maize. Phytochem. 65, 3187–3198.
<https://doi.org/10.1016/j.phytochem.2004.10.015>
- Manetti, C., Bianchetti, C., Casciani, L., Castro, C., Di Cocco, M.E., Miccheli, A., Motto, M., Conti, F., 2006. A metabonomic study of transgenic maize (*Zea mays*) seeds revealed variations in osmolytes and branched amino acids. J. Exp. Bot. 57, 2613–2625.
<https://doi.org/10.1093/jxb/erl025>
- Masoero, F., Moschini, M., Rossi, F., Prandini, A., Pietri, A., 1999. Nutritive value, mycotoxin contamination and *in vitro* rumen fermentation of normal and genetically modified corn (CRY 1A(B)) grown in northern Italy. Maydica 44, 205–209.
- Medina, A., Gilbert, M.K., Mack, B.M., OBrian, G.R., Rodríguez, A., Bhatnagar, D., Payne, G., Magan, N., 2017. Interactions between water activity and temperature on the *Aspergillus flavus* transcriptome and aflatoxin B₁ production. Int. J. Food Microbiol. 256, 36–44. <https://doi.org/10.1016/J.IJFOODMICRO.2017.05.020>

- 478 Medina, A., Rodríguez, A., Magan, N., Rodriguez, A., Magan, N., 2014. Effect of climate
479 change on *Aspergillus flavus* and aflatoxin B₁ production. Front. Microbiol. 5, 1–7.
480 <https://doi.org/10.3389/fmicb.2014.00348>
- 481 Mohale, S., Magan, N., Medina, A., 2013. Comparison of growth, nutritional utilisation
482 patterns, and niche overlap indices of toxigenic and atoxigenic *Aspergillus flavus*
483 strains. Fungal Biol. 117, 650–659. <https://doi.org/10.1016/j.funbio.2013.07.002>
- 484 Munkvold, G., 2014. Crop management practices to minimize the risk of mycotoxins
485 contamination in temperate-zone maize, In: Leslie, J.F., Logrieco, A. (Eds.),
486 Mycotoxin Reduction in Grain Chains. John Wiley & Sons, Chichester, UK, pp. 59–
487 77. <https://doi.org/10.1002/9781118832790.ch5>
- 488 Nesci, A., Rodriguez, M., Etcheverry, M., 2003. Control of *Aspergillus* growth and aflatoxin
489 production using antioxidants at different conditions of water activity and pH. J. Appl.
490 Microbiol. 95, 279–287. <https://doi.org/10.1046/j.1365-2672.2003.01973.x>
- 491 Northolt, M.D., van Egmond, H.P., Paulsch, W.E., 1977. Differences between *Aspergillus*
492 *flavus* strains in growth and aflatoxin b₁ production in relation to water activity and
493 temperature. J. Food Prot. 40, 778–781. <https://doi.org/10.4315/0362-028X-40.11.778>
- 494 Peromingo, B., Rodríguez, A., Bernáldez, V., Delgado, J., Rodríguez, M., 2016. Effect of
495 temperature and water activity on growth and aflatoxin production by *Aspergillus*
496 *flavus* and *Aspergillus parasiticus* on cured meat model systems. Meat Sci. 122, 76–
497 83. <https://doi.org/10.1016/j.meatsci.2016.07.024>
- 498 Piccioni, F., Capitani, D., Zolla, L., Mannina, L., 2009. NMR Metabolic Profiling of
499 Transgenic Maize with the *CryIA(b)* Gene. J. Agric. Food Chem. 57, 6041–6049.
500 <https://doi.org/10.1021/jf900811u>
- 501 Qaim, M., 2014. Genetically modified crops and development, In: Desai, V., Potter, R.
502 (Eds.), The Companion to Development Studies. Routledge, pp. 249–254.
- 503 Reuter, T., Aulrich, K., Berk, A., Flachowsky, G., 2002. Investigations on genetically
504 modified maize (Bt-maize) in pig nutrition: chemical composition and nutritional
505 evaluation. Arch. Tierernahr. 56, 23–31.
- 506 Sanchis, V., Magan, N., 2004. Environmental conditions affecting mycotoxins, in: Magan,
507 N., Olsen, M. (Eds.), Mycotoxins in Food. Elsevier, pp. 174–189.
508 <https://doi.org/10.1533/9781855739086.2.174>
- 509 Scarpari, M., Punelli, M., Scala, V., Zaccaria, M., Nobili, C., Ludovici, M., Camera, E.,
510 Fabbri, A.A., Reverberi, M., Fanelli, C., 2014. Lipids in *Aspergillus flavus*-maize
511 interaction 5, 74. <https://doi.org/10.3389/fmicb.2014.00074>

- Scheidegger, K. a., Payne, G. a., 2003. Unlocking the secrets behind secondary metabolism: A review of *Aspergillus flavus* from pathogenicity to functional genomics. *Toxin Rev.* 22, 423–459. <https://doi.org/Doi.10.1081/Txr-120024100>
- Schnepf, E., Crickmore, N., Van Rie, J., Lereclus, D., Baum, J., Feitelson, J., Zeigler, D.R., Dean, D.H., 1998. *Bacillus thuringiensis* and its pesticidal crystal proteins. *Microbiol. Mol. Biol. Rev. MMBR* 62, 775–806.
- Simó, C., Ibáñez, C., Valdés, A., Cifuentes, A., García-Cañas, V., 2014. Metabolomics of genetically modified crops. *Int. J. Mol. Sci.* 15, 18941–66. <https://doi.org/10.3390/ijms151018941>
- Walley, J.W., Kliebenstein, D.J., Bostock, R.M., Dehesh, K., 2013. Fatty acids and early detection of pathogens. *Curr. Opin. Plant Biol.* 16, 520–526. <https://doi.org/10.1016/J.PBI.2013.06.011>
- Windham, G.L., Williams, W.P., Davis, F.M., 1999. Effects of the southwestern corn borer on *Aspergillus flavus* kernel infection and aflatoxin accumulation in maize hybrids. *Plant Dis.* 83, 535–540. <https://doi.org/10.1094/PDIS.1999.83.6.535>
- Wu, F., 2006. Mycotoxin reduction in Bt corn: Potential economic, health, and regulatory impacts. *Transgenic Res.* 15, 277–289. <https://doi.org/10.1007/s11248-005-5237-1>

Figure legends

Figure 1. Comparison of the growth rate (mm day^{-1}) of the three toxigenic and one non-toxigenic strains of *A. flavus* on maize media based on non-GM and GM grain in relation to water availability ($0.99-0.90 a_w$) and temperature ($25-35^\circ\text{C}$). Bars indicate Average $\pm$ Standard Error.

Figure 2. Contour maps of the growth rate (mm day^{-1}) of *Aspergillus flavus* with pooled data from all the maize cultivars used as substrate (GM + non-GM) for (A) toxigenic strain (AfLRG-a) and (B) non-toxigenic strain (AfPHx) over 10 days incubation. The colour scale (lighter > darker tones) indicates higher growth rate (mm day^{-1}) in relation to the interacting abiotic factors.

Figure 3. Effect of colonisation of the maize-based media using non-GM and GM maize kernels on aflatoxin B₁ production (ng g^{-1}) under different water activity conditions at $25-35^\circ\text{C}$. Bars indicate Average $\pm$ Standard Error. *Not the same scale.

Figure 4. Contour plots for aflatoxin B₁ (ng g^{-1}) production by a toxigenic strain of *Aspergillus flavus* (AfP30C) on milled maize agar using the non-GM maize cultivars (A) AS 1555 CON, (B) P30F53 CON and (C) P2530 CON and, their respective isogenic GM lines (D) AS 1555 PRO®, (E) P30F53 H® and (F) P2530 Hx® as substrates. The colour scale (lighter > darker tones) indicates higher concentration of AFB₁ (ng g^{-1}) in relation to the interacting abiotic factors.

Figure 5. Contour plots for aflatoxin B₁ (ng g^{-1}) production by the toxigenic strain of *Aspergillus flavus* (AfLRG-a) on milled maize agar using the non-GM maize cultivars (A) AS 1555 CON, (B) P30F53 CON and (C) P2530 CON and, their respective isogenic GM lines (D) AS 1555 PRO®, (E) P30F53 H® and (F) P2530 Hx® as substrates. The colour scale (lighter > darker tones) indicates higher concentration of AFB₁ (ng g^{-1}) in relation to the interacting abiotic factors.

Supplementary Figures

Suppl. Figure 1. Growth rate (mm day^{-1}) for the *A. flavus* strains AfLRG-a, AfASC-a,

AfP30C-a and AfPHx on milled maize agar using the non-GM maize cultivars (AS 1555 CON, P30F53 CON and P2530 CON) and their respective isogenic GM lines (AS 1555 PRO®, P30F53 H® and P2530 Hx®) as a substrate. Bars represent AV±SE.

Suppl. Figure 2. Aflatoxin production (AFB1 ng g⁻¹) for the *A. flavus* strains, AfASC-a, AfLRG-a and AfP30C-a on milled maize agar using the non-GM maize cultivars (AS 1555 CON, P30F53 CON and P2530 CON) and their respective isogenic GM lines (AS 1555 PRO®, P30F53 H® and P2530 Hx®) as a substrate. Bars represent AV±SE.

Suppl. Figure 3. Contour plots for aflatoxin B₁ (ng g⁻¹) production by a toxigenic strain of *Aspergillus flavus* (AfASC-a) on milled maize agar using the non-GM maize cultivars (A) AS 1555 CON, (B) P30F53 CON and (C) P2530 CON and, their respective isogenic GM lines (D) AS 1555 PRO®, (E) P30F53 H® and (F) P2530 Hx® as substrates. The colour scale (lighter > darker tones) indicates higher concentration of AFB1 (ng g⁻¹) over the variables.

Table 1. Maize cultivars (non-GM and GM) selected for the ecophysiological studies as substrate for *A. flavus* growth. Key: IR- insect resistance; HT – herbicide tolerance.

Conventional cultivar (non-GM)	Isogenic line (GM)	Tolerance traits present in the GM cvs
AS 1555 CON	AS 1555 PRO [®]	IR - Lepidopteran Antibiotic resistance
P30F53 CON	P30F53 H [®]	HT - Glufosinate ammonium and glyphosate IR - Lepidopteran
P2530 CON	P2530 Hx [®]	IR - Lepidopteran

Table 2. Statistical effect of abiotic and biotic factors on growth and aflatoxin B₁ production by *A. flavus* strains.

Effects	<i>p</i> -value (Prob>ChiSq)	
	Growth (mm/day)	AFB ₁ (ng/g) *
Fungal strains	0.4661	0.0013
Maize cvs	<.0001	0.0020
T°C	<.0001	0.0002
a _w	<.0001	<.0001
GM × non-GM	0.0616	0.6002
Maize cv × T°C	0.0683	0.0020
Maize cv × a _w	0.0683	<.0001
Maize cv × T°C × a _w	0.0683	0.0002

$P \geq 0.05$ was considered to be not statistically significant. Non-parametric Wilcoxon/Kruskal-Wallis Tests with 1-Way Test Chi-Square Approximation.

The *p*-value (Prob>ChiSq) was calculated from the values of ChiSquare and DF (degrees of freedom) of each effect.

* The effects on AFB₁ excluded the non-toxigenic *A. flavus* strain AfPHx.




Figure 1.




Figure 2.




Figure 3.




Figure 4.






Figure 5.




Supplementary Figure 1.

Growth rate (mm day⁻¹) for the *A. flavus* strains *AfLRG-a*, *AfASC-a*, *AfP30C-a* and *AfPHx* on milled maize agar using the non-GM maize cultivars (AS 1555 CON, P30F53 CON and P2530 CON) and their respective isogenic GM lines (AS 1555 PRO[®], P30F53 H[®] and P2530 Hx[®]) as substrate. Bars represent AV±SE.



Supplementary Figure 2.

Aflatoxin production (AFB₁ ng g⁻¹) for the *A. flavus* strains, AfASC-a, AfLRG-a and AfP30C-a on milled maize agar using the non-GM maize cultivars (AS 1555 CON, P30F53 CON and P2530 CON) and their respective isogenic GM lines (AS 1555 PRO[®], P30F53 H[®] and P2530 Hx[®]) as substrate. Bars represent AV±SE.



Supplementary Figure 3.

Contour plots for aflatoxin B₁ (ng g⁻¹) production by a toxigenic strain of *Aspergillus flavus* (AfASC-a) on milled maize agar using the non-GM maize cultivars (A) AS 1555 CON, (B) P30F53 CON and (C) P2530 CON and, their respective isogenic GM lines (D) AS 1555 PRO®, (E) P30F53 H® and (F) P2530 Hx® as substrates. The colour scale (lighter > darker tones) indicates higher concentration of AFB₁(ng g⁻¹) over the variables.

Highlights

- *A. flavus* strains grew similarly on GM/non-GM nutritional media under interacting abiotic factors
- Contour maps of optimum and marginal conditions for growth were developed for 4 strains of *A. flavus*
- Aflatoxin B₁ production was optimal at 0.99 water activity and 30 °C, regardless of strain.
- Optimum and marginal toxin production contour maps were determined for toxigenic strains



Vincent Building
Cranfield, Bedfordshire,
MK43 0AL, England
Tel +44 (0) 1234 7538308
www.cranfield.ac.uk/sas

8th September, 2021
Senior Editor,
Fungal Biology

Re: Conflict of Interests Statement

We can confirm that there are no conflict of interests for the authors of this manuscript submitted to the Journal.

This was part of a PhD thesis which the Senior Author completed at Cranfield under the 'Science without Borders' scholarship scheme.

With best wishes,
Yours sincerely,

A handwritten signature in blue ink that reads 'N. Magan'.

Prof. N. Magan DSc
Head, Applied Mycology Group,
Cranfield Soil and AgriFood Institute,
Environment and AgriFood Theme
(n.magan@cranfield.ac.uk)



THE QUEEN'S
ANNIVERSARY PRIZES
FOR HIGHER AND FURTHER EDUCATION
2015 & 2017



DSPACE

<https://dspace.org/>

Comparison of growth and aflatoxin B1 production profiles of *Aspergillus flavus* strains on conventional and isogenic GM-maize-based nutritional matrices

Gasparini, Alessandra M.; Medina, Angel; Magan, Naresh

2021-10-30

Elsevier

<https://doi.org/10.1016/j.funbio.2021.10.006>

Gasparini AM, Medina A, Magan N. (2022) Comparison of growth and aflatoxin B1 production profiles of *Aspergillus flavus* strains on conventional and isogenic GM-maize-based nutritional matrices. *Fungal Biology*, Volume 126, Issue 1, January 2022, pp. 82-90