

Pre-print version of the paper: Rivas Casado, M., Mead, A., Burgess P.J., Howard, D.C., Butler, S. (2014). Predicting the impact of land-based renewable energy on farmland birds at a landscape scale. *Science of the Total Environment* 476-477: 7-19.

<http://dx.doi.org/10.1016/j.scitotenv.2013.12.080>

Predicting the impacts of bioenergy production on farmland birds

Monica Rivas Casado^{a*}, Andrew Mead^b, Paul J Burgess^a, David C Howard, Simon J Butler^d

^aDepartment of Environmental Science and Technology, Cranfield University, Cranfield, MK43 0AL, UK

^bSchool of Life Sciences, University of Warwick, Wellesbourne, Warwick, CV35 9EF, UK

^cCentre for Ecology and Hydrology, Lancaster Environment Centre, Bailrigg, Lancaster, LA1 4AP, UK.

^dSchool of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

* Corresponding author. Contact details for corresponding author: m.rivas-casado@cranfield.ac.uk, Tel ++44 (0)1234750111 ext 2706.

Highlights

- First GIS model to predict spatially the “local” impact of bioenergy policies
- Systematic functional space models covering 19 farmland bird species.
- A 16,000 ha case study shows a synergy between bioenergy and farmland bird populations.
- Renewable energy production strategy affects impact on farmland bird populations.
- The model provides a method to determine the effects of policy driven land use change on biodiversity.

Abstract

Meeting European renewable energy production targets is expected to cause significant changes in land use patterns. With an EU target of obtaining 20% of energy consumption from renewable sources by 2020, national and local policy makers need guidance on the impact of potential delivery strategies on ecosystem goods and services to ensure the targets are met in a sustainable manner. Within agroecosystems, models are available to explore consequences of such policy decisions for food, fuel and fibre production but few can describe the effect on biodiversity. This paper describes the integration and application of a farmland bird population model within a geographical information system (GIS) to explore the consequences of land use changes arising from differing strategies to meet renewable energy production targets. Within a 16,000 ha arable dominated case study area in England, the population growth rates of 19 farmland bird species were predicted under baseline land cover, a scenario maximising wheat production for bioethanol, and a scenario focused on mix of bioenergy sources. Both scenarios delivered renewable energy production targets for the region (>12 kWh per person per day) but, despite differences in resultant landscape composition, the response of the farmland bird community as a whole to each scenario was small and broadly similar. However, this similarity in overall response masked significant intra- and inter-specific variation across the study area and between scenarios suggesting contrasting mechanisms of impact and highlighting the need for context dependent,

species-level assessment of land use change impacts. This framework provides one of the first systematic attempts to spatially model the effect of policy driven land use change on the population dynamics of a suite of farmland birds. The GIS framework also facilitates its integration with other ecosystem service models to explore wider synergies and trade offs arising from national or local policy interventions.

Keywords: farmland birds, ecosystem services, Geographical Information System, land use, renewable energy

1. Introduction

Finite fossil fuel resources and the need to reduce greenhouse gas emissions have led to a global focus on increasing energy supplies from renewable sources. The European Union has set a target of obtaining 20% of energy consumption from renewable sources by 2020 (EC, 2009). The target set for the UK is 15%, which would be equivalent to renewable energy providing the equivalent of 4.6 kWh of electricity, 3.4 kWh of transport fuel and 3.7 kWh of heat per person per day (Burgess *et al.*, 2012). In 2011, the proportion of gross energy consumption from renewable sources was 13.4% within the EU27 but only 3.8% in the UK (EurObserv'ER, 2013). Realizing the 2020 targets will require a significant change in land use patterns at local, national, European (Rounsevell *et al.*, 2003) and even global scales. The recent revision of EU renewable energy policy (European Commission, 2012) in light of concerns over its impact on food production means that the long term implications for land use are unclear but in Britain, this may initially be an expansion or redirection of arable crops such as wheat and oilseed rape as first generation transport fuel production (Gallagher, 2008) and/or an expansion in the area under biomass crops, such as perennial grasses (e.g. miscanthus *Miscanthus giganteus*) and short rotation coppice (Burgess *et al.*, 2012; Committee on Climate Change, 2011).

Large scale, often policy driven, land use changes have the potential to cause unexpected and significant detrimental environmental impacts. In Europe, for example, this is perhaps best evidenced by significant declines in farmland biodiversity and deteriorations in soil, air and water quality over recent decades associated with agricultural intensification and land abandonment and driven to a great extent by the Common Agricultural Policy (Stoate *et al.*, 2001). There is also already evidence of unforeseen detrimental environmental impacts resulting from renewable energy policies. Rapidly increasing demand for biofuels, driven in part at least by EU policy (European Commission, 2006), have caused significant damage to biodiversity and ecosystem service provision through both direct and indirect land use change with impact reported in parts of South America and south east Asia in particular (e.g. Fargione *et al.*, 2008; Fitzherbert *et al.*, 2008). In implementing EU renewable energy policy it is crucial that we learn from these past mistakes and manage the delivery of renewable energy production targets in a sustainable manner (Petersen *et al.*, 2007). In particular this requires that renewable energy policies are integrated with other policies designed to manage issues such as food production and biodiversity conservation policies so that trade offs made between these potentially conflicting demands for finite land resources are sustainable (Murphy *et al.*, 2011). A key component of this is developing the capability to predict any potential detrimental environmental impacts of

proposed land use and management changes so that appropriate prevention or mitigation actions can be identified and implemented where necessary.

Here we focus on the effects of policy driven renewable energy options on farmland biodiversity, using the impact on birds as a proxy for the consequences for wider biodiversity. Both the UK and other European governments have identified birds as indicators of biodiversity health and have adopted indices of population trends as headline indicators of sustainable development. More broadly, bird population trends have also been used as an indicator of continued biodiversity losses at a global scale (Butchart *et al.*, 2010). Hence the objective of this paper is to use a recently published modelling framework (Butler and Norris, 2013), integrated into a geographical information system (GIS), to predict the response of farmland bird populations to land use change scenarios associated with delivering renewable energy production targets for a landscape in the UK.

2. Method

The modelling framework uses the concept of functional cover types to link land use to the population trends of farmland birds. In brief, structural land covers (e.g. wheat, grassland, woodland) are classified into functional land covers (e.g. foraging and nesting sites) according to their capacity to provide key resources. This approach provides a more mechanistic link between land use and population growth than more traditional habitat association models, it helps to reduce content specificity, and it facilitates the incorporation of novel land uses (Butler and Norris, 2013). The quantity, in terms of area, and quality, in terms of resource provision, of each functional cover type in a landscape effectively delimits the functional space available to a species. In brief, Butler and Norris (2013) modelled the population trends of each of the 19 species included in the UK Farmland Bird Index at a 1km square level as a function of the availability of six functional space components at the same spatial scale: high and low quality summer foraging space (SHQ and SLQ respectively), high and low quality breeding space (BHQ and BLQ respectively) and high and low quality winter foraging space (WHQ and WLQ respectively). Parameter estimates for each functional space component, reflecting the strength and direction of its relationship with local population trends, were derived using a model averaging approach underpinned by bird abundance and habitat data collected from more than 600 1 km squares covered by both the Breeding Bird Survey (BBS) (Risely *et al.*, 2011) and Winter Farmland Bird Survey (WFBS) (Gillings *et al.*, 2008) (see Butler and Norris, 2013 for full details).

The first stage of our automated process used a GIS platform (ArcGIS version 9.3; ESRI Inc) to generate habitat data in the same format as used in the BBS and WFBS habitat surveys. In the second stage these habitat data were classified into functional space and used to predict farmland bird population trends. Full details of this process are provided below. A toolbox named “BirdMod” was developed to undertake these analyses, which can be installed and run on a standard computer. The script was developed using ModelBuilder and runs in Visual Basic or Python.

2.1. Case study area

The Marston Vale extends over about 16,000 ha in Bedfordshire in lowland England (Fig. 1a). Once currently consented urban developments are in place, the population density (3.1 ha^{-1}) and proportion of area allocated to agriculture (69%) and woodland

(8%) will broadly reflect national values. However the area under crops and fallow (52%) is higher and the area under grassland (17%) is lower than the equivalent national means (30% and 37% respectively). The work presented here is part of a wider project exploring the interactions between renewable energy demand and supply, land use and the stocks and flows of ecosystem services and goods in the area (Burgess *et al.*, 2012; Howard *et al.*, 2013).

Land use across the Marston Vale was digitised using aerial photography from 2005 with polygons generated for each field, woodland, major road, watercourse, urban and commercial area (Table 1). This landscape configuration is hereafter referred to as *BASELINE*.




Fig. 1. Land use maps of the Marston Vale (2009): (a) location of the Marston Vale (b) *BASELINE*, (c) *MAXIMIZE* and (d) *RESILIENCE* scenarios.

Aerial images from Google EarthTM were used to assess the relative proportion of specific boundary types. Within ten randomly selected 1 km squares, all field boundaries were classified as either a) hedgerow with trees, b) hedgerow without trees, c) tree line with no hedge or d) no vertical structure; these classifications match those used to describe boundary features in BBS. The relative proportions of each boundary type across the ten squares were calculated as 0.22 (± 0.06), 0.30 (± 0.06), 0.03 (± 0.01) and 0.45 (± 0.05) respectively and these values were used for the whole of Marston Vale in subsequent landscape structure assessments (but see sensitivity analyses below).

Table 1. Assumed area for each land use type in Marston Vale, under BASELINE conditions (assuming consented development takes place), a MAXIMIZE scenario (focused on meeting renewable energy targets from wheat), and a RESILIENCE scenario (focused on meeting renewable energy targets from a range of crops)

Land use	Area (ha)		
	<u>BASELINE</u>	<u>MAXIMIZE</u>	<u>RESILIENCE</u>
Wheat	4150	10745	4150
Grass	2596	0	2596
Winter oilseed rape	1209	0	1752
Fallow	984	0	315
Other spring crop	693	0	693
Barley	455	0	455
Crop	392	0	392
Spring oilseed rape	263	0	0
Bare soil	3	0	3
Miscanthus	0	0	193
Urban	1844	1844	1844
Commercial areas	372	372	372
Transport	279	279	279
Landfill	235	235	235
Woodland	1232	1232	1232
Woodland screening	186	186	186
Short rotation coppice	0	0	196
Water body	351	351	351
Other	853	853	853
Total	16097	16097	16097

2.2. *Future landscape scenarios*

Two alternative scenarios representing different approaches to increasing land based renewable energy production within Marston Vale were constructed for this study. These were principally defined to illustrate the application of BirdMod for exploring contrasting energy production scenarios and therefore represent plausible rather than optimal land use configurations. In each scenario, polygons classified as woodland, urban, commercial, transport, water or landfill stayed the same as in BASELINE. The first, hereafter referred to as MAXIMIZE, represents an extreme example, with all arable and grassland areas planted with wheat. Through the harvest of grain for bioethanol production and straw for heat, this offers the greatest gross energy output (Burgess *et al.*, 2012) (Fig. 1c). In the second scenario, hereafter referred to as RESILIENCE, the objective was to maximise renewable energy targets without an undue reliance on any individual renewable energy source (Grubb *et al.*, 2006). Similar areas of land were

allocated to wheat, grass and barley as in *BASELINE* but a greater area was allocated to winter oilseed rape and small areas of miscanthus and short rotation coppice were introduced. As a consequence, the area of fallow land decreased and spring oilseed rape was lost from rotations (Fig. 1d). The land use allocation under *BASELINE*, *MAXIMIZE* and *RESILIENCE* is summarised in Table 1. Using an existing framework for exploring trade offs between land use, renewable energy, food, feed and wood production (Burgess *et al.*, 2012), we calculated the capacity of *BASELINE*, *MAXIMIZE* and *RESILIENCE* landscapes to meet a range of energy demand types within Marston Vale. Currently, the level of food production is greater than the local demand within Marston Vale (see results and Table 2). We therefore also estimated energy output capacity for *BASELINE* under a scenario where, once local food demand is met, “surplus” wheat and oilseed rape are used for bioethanol and biodiesel production and arable straw and the non-timber biomass of woodlands used for heating.

2.3. Predicting farmland bird trends from functional space availability

To mirror the BBS and WFBS habitat recording methodologies, calculations within the BirdMod toolbox were based on 1 km (100 ha) British Ordnance Survey grid squares overlain on the land use map. All squares containing less than 50 ha farmland, whether due to the extensive presence of other land use types (e.g. woodland or urban) or because the boundary of the Marston Vale bisected them, were excluded in accordance with original model parameterisation rules (Butler and Norris, 2013) (Fig. 2). Summer and winter habitat within the remaining squares under *BASELINE*, *MAXIMIZE* and *RESILIENCE* were then quantified as follows:

2.3.1. Summer foraging and breeding habitat

Two transects (1000 m x 50 m), each subdivided into 200 m x 50 m sections, were overlain on each grid square (Figs. 2b and 2c). If a square overlapped the boundary of the Marston Vale but was retained in our analyses because the section falling inside contained more than 50 ha farmland (see above), the transects stopped at the boundary and the total number of complete 200 m sections may have been less than 10. The area of each land use type encompassed by each transect section was quantified, as was the proportion of each classified as “disturbed” or “undisturbed”; on the basis of BBS habitat coding methodology, “disturbed” areas were defined as land within 50 m of an urban settlement or road (Figs. 2d and 2e). The length of any boundary features falling within each 200 m x 50 m section was also calculated. If this was greater than 50 m, boundary characteristics were included in the classification of habitat features for that section. A set of habitat allocation algorithms (see Appendix A, Figs. A.1-A.3 in Supporting Information) was then applied to these data to assign primary and secondary BBS habitat classifications to each transect section. Boundary characteristics and polygon-specific spring or autumn sowing date for cereals, were assigned using probability based number generators, underpinned by direct observation across Marston Vale and Defra Agricultural and Horticultural Census data for Bedfordshire (2005-2008 data: www.defra.gov.uk) respectively.




Fig. 2. The different steps run within BirdMod to quantify habitat availability prior to reclassification into functional space: (a) location of the 1 km x 1 km squares within the Marston Vale, (b) example of the location of the 200 m x 50 m transect sections (red lines) within each square, (c) distance between parallel transects, (d) detail of the different land uses identified on one of the transects showing the disturbed (red dots) areas and (e) table summarising the land uses for the selected transect as estimated by BirdMod. Grey cells show squares that have been excluded from the analysis (e.g. area of farmland is <50 ha).

2.3.2. Winter foraging habitat

The digitized land use maps described above were built from spring and summer land use data. An additional habitat allocation algorithm (Appendix A, Fig. A.4) was therefore used to backcast from these data to predict the WFBS habitat code for each

polygon in the preceding winter. Polygon-specific allocation of spring or autumn sowing date for cereals and stubble weediness, assigned using probability based number generators, was underpinned where necessary by and Defra Agricultural and Horticultural Census data for Bedfordshire and WFBS habitat data respectively.

2.3.3. *Quantifying functional space availability*

Butler and Norris (2013) identified the BBS and WFBS codes contributing to each of the six functional space components identified (SHQ, SLQ, BHQ, BLQ, WHQ and WLQ) of each Farmland Bird Index species. We used the same classifications and methodology to quantify functional space for each species in each square under BASELINE, MAXIMIZE and RESILIENCE based on the BBS and WFBS classifications generated by BirdMod. For each species, the number of transect sections providing BHQ and BLQ within each square, weighted by whether it was provided by the primary and/or secondary habitats, was divided by the total number of transect sections in that square and multiplied by 100 to estimate the total area (ha) of BHQ and BLQ available. This process was repeated to quantify the area of SHQ and SLQ available for each species in each square. Finally, the summed areas of polygons with WFBS habitats classified as providing WHQ or WLQ were calculated for each species in each square. Two energy crops, miscanthus and short rotation coppice, which are not currently present in Marston Vale, were introduced into the landscape in the RESILIENCE scenario. They were assigned BBS and WFBS codes for broadly equivalent structural cover types and their contribution to the six functional space components for each species was assessed accordingly. For summer foraging and breeding cover, short rotation coppice was equated to a young woodland plantation with moderate shrub and field layer and to a farm scrub patch for winter foraging cover. Equivalent structural cover types in the current landscape were less apparent for miscanthus. For summer foraging and breeding cover, it was coded as an arable crop, but restrictions to its contribution to functional space were applied in line with the expected influence of the much taller, denser structure on food availability and perceived/actual predation risk for each species (Butler *et al.*, 2005; Whittingham and Devereux, 2008). Similarly, for winter foraging cover, miscanthus was broadly equated to a tall cereal crop but the structure of miscanthus crops over winter and its impact on resource availability were again taken into account when defining the quality of functional space provided (Sage *et al.*, 2006, 2010).

For each farmland bird species, high and low quality classifications of each functional cover type were mutually exclusive for any given polygon so the total area (i.e. high plus low quality) of breeding, summer foraging and winter foraging functional cover within a 1 km square could not exceed 100 ha. However, a polygon could potentially contribute to more than one functional cover type for each species so the area of functional space (i.e. breeding plus summer foraging plus winter foraging functional cover) within a square could exceed 100 ha.

Butler and Norris (2013) also showed that conspecific abundance in the surrounding landscape influences both population trends and the relationship between functional space and population trends. To account for this, they included a measure of conspecific abundance in the surrounding landscape, calculated as the distance weighted average of observed counts over a three year period for that species in all BBS squares, in their functional space models. To calculate the equivalent metric, we first calculated the average BBS count of each species in each BBS/WFBS square based on the three years

immediately prior to the year the digital photographs (i.e. 2002, 2003 and 2004) were taken; if a square was not surveyed in one or more of these years, records from the closest three years were used. We then calculated a weighted average of these counts for each species and each square based on the Euclidean distance between that square and each BBS/WFBS square. Parameter estimates for each species' functional space response (Butler and Norris, 2013) were then applied to the functional space area and conspecific abundance data to calculate annual population growth rate (*pgr*) in each square. It is important to note that elements of the automation process described above are stochastic because random number generators underpin the assignment of particular habitat characteristics, such as spring or autumn sown cereals or boundary type, to each polygon when relative availability is dictated by set probabilities (Appendix A). We therefore repeated this process ten times and used the average *pgr* predicted for each species in each square in subsequent analyses.

To explore the sensitivity of the model to specified thresholds of key model parameters associated with the proportions of spring sown cereal and weedy stubbles present and the assignment of boundary feature characteristics, a series of sensitivity analyses were also undertaken using the BASELINE scenario: 1) Based on Defra Agricultural and Horticultural Census data for Bedfordshire, the probability of any polygon characterised as "Wheat" being identified as autumn sown was set at 0.9. We explored the impact on the predicted *pgr* of each species if this was increased to 1 or decreased to 0.8; 2) Based on WFBS habitat data, the probability of any polygon characterised as "Stubble" during the winter being characterised as "Weedy" (as opposed to "Clean") was set at 0.5. We explored the impact on the predicted *pgr* of each species if this was reduced to 0.25 or increased to 0.75; 3) As detailed above, the probabilities of a particular boundary being classified as either hedgerow with trees, hedgerow without trees, tree line with no hedge or no vertical structure were set at 0.22, 0.30, 0.03 and 0.45 respectively. Three alternative proportional classifications (i: 0.25, 0.25, 0.25, 0.25; ii: 0.5, 0.5, 0, 0; iii: 0, 0, 0.5, 0.5 respectively) were tested to assess the impact on the predicted *pgr* of each species. Sensitivity analyses for each parameter adjustment were carried out in isolation, with each configuration iterated ten times as described above. The response of each species to land use changes under MAXIMIZE and RESILIENCE was also calculated under each set of parameter values.

The impact of the land use changes associated with each scenario on the *pgr* of individual species and the community as a whole (i.e. *pgr* averaged across all 19 species) was assessed using paired *t* tests, with each 1 km square under BASELINE paired with the corresponding square under MAXIMIZE and RESILIENCE. The average *pgr* across all species effectively represents the expected extent and direction of the annual change in the Farmland Bird Index for the study site under each scenario. The paired *t* test works under the assumption that the paired differences are independent and identically normally distributed; this second assumption was broken in the cases of Turtle dove and Woodpigeon and Wilcoxon's signed ranks test, a nonparametric method analogous to the paired *t* test, was used for these species instead.

3. Results

3.1 Bioenergy production

The daily energy demand per person within Marston Vale equates to about 80 kWh. Under BASELINE land cover patterns and prioritisation of food production, the output of heat and transport energy is assumed to be zero. If surplus food products were reallocated to energy production, it was estimated that BASELINE energy output could be increased to 11.3 kWh p⁻¹ d⁻¹, comprising 4.9 kWh p⁻¹ d⁻¹ for transport fuel and 6.4 kWh p⁻¹ d⁻¹ for heating (Table 2). The combined value is similar value to the 2020 renewable targets, but it still only represents about 15% of the total energy requirement. Under the MAXIMIZE scenario, conversion of all arable and grassland areas to wheat was calculated to increase potential production levels to 11.4 kWh p⁻¹ d⁻¹ of transport fuel and 9.6 kWh p⁻¹ d⁻¹ for heating (Table 2). The output of animal feed was also predicted to increase because of the formation of distillers grains in bioethanol production. Under the RESILIENCE scenario, the transport fuel availability was marginally greater than under BASELINE, because of the greater area of oilseed rape, and the area of miscanthus and short rotation coppice contributed to an increase in the available energy for heating.

Table 2. Equivalent per capita demand in the UK for energy, food, feed and timber, the renewable energy targets for 2020, the capacity for the current land use in the Marston Vale (BASELINE) to meet those demands assuming prioritisation of use for food or energy, and the corresponding outputs for a scenario maximising the wheat area and bioethanol production (MAXIMIZE), and a RESILIENCE scenario. The output is expressed in terms of equivalent energy per person per day (kWh p⁻¹ d⁻¹). The methodology for determining the values is described by Burgess *et al.* (2012)

Form of demand	Current demand	Renewable target (2020)	Output capacity			
			<u>BASELINE</u> <i>prioritise food</i>	<u>BASELINE</u> <i>prioritise energy</i>	<u>MAXIMIZE</u>	<u>RESILIENCE</u>
Electricity	15.0	4.6	0.0	0.0	0.0	0.0
Transport fuel	34.0	3.4	0.0	4.9	11.4	5.1
Heat	31.0	3.7	0.0	6.4	9.6	7.2
Energy subtotal	80.0	11.7	0.0	11.3	21.0	12.3
Food	1.9		9.4	1.9	1.9	1.9
Animal feed	5.6		4.2	7.3	9.9	7.3
Timber	4.4		0.9	0.9	0.9	0.9

3.2 Community-level response

The mean predicted annual *pgr* across all 19 species for BASELINE was -0.0091 ± 0.006 (Table 3). This represents an annual decline in farmland bird populations of 0.91%. The MAXIMIZE scenario was predicted to result in a significantly greater mean rate of decline across the 19 species (-0.0106 ± 0.0034 ; paired *t* test: $t = 3.91$, $n = 142$, $p < 0.01$). Similarly, changing from BASELINE to the RESILIENCE scenario was also

predicted to lead to a significantly greater rate of decline (-0.0103 ± 0.0053 ; paired t test: $t = 2.62$, $n = 142$, $p < 0.01$). This suggests that the Farmland Bird Index would continue declining under each scenario but the rate of decline would be greatest under MAXIMIZE (Table 3). The above values are the mean predicted *pgr* values across all 142 1 km squares; some individual squares showed positive values, and some showed much larger negative values (Fig. 3).




Fig. 3. Mean annual *pgr* across the 19 bird species predicted for the (a) BASELINE, (b) MAXIMIZE and (c) RESILIENCE scenarios. Positive and negative values were coded using a blue and red coloured scale, respectively. The range and the breaks for each of the scales were determined to enhance visualisation.

3.3. Species-level response

When averaged across all squares, eleven species were predicted to have a negative *pqr* under each scenario whilst eight species were predicted to have a positive *pqr* under all three landscape configurations (Table 3, Appendix B Fig. B.1). Under MAXIMIZE, eight species were predicted to have significantly lower *pqr* than that predicted for BASELINE ($p < 0.05$ in all cases) but this did not involve an overall change in the direction of population trajectory for any. Of these, six are species that were predicted to be declining under BASELINE. Ten species were predicted to have significantly higher *pqr* under MAXIMIZE than under BASELINE ($p < 0.05$ in all cases) but again this did not result in a change in the direction of population trajectory for any. Of these, four were predicted to be declining under BASELINE. There was no significant change in the predicted *pqr* of Kestrel *Falco tinnunculus* between BASELINE and MAXIMIZE.

Changing from BASELINE to RESILIENCE led to significant declines in the predicted *pqr* of twelve species ($p < 0.05$ in all cases) and significant increases in the predicted *pqr* of just two (Yellow wagtail *Motacilla flava* and Woodpigeon *Columba palumbus*, $p < 0.05$ in both cases). Of those predicted to show significant decline in predicted *pqr*, seven are predicted to be declining under BASELINE. For no species did the change in land use result in a switch in the overall direction of predicted population trajectory.

Table 3. Mean and standard deviation (s.d.) of the predicted square level *pqr* under each land use scenario for each of the 19 species considered and the community as a whole (i.e. *pqr* averaged across the 19 species). Species are listed according to predicted mean *pqr* for BASELINE. Results for the square level paired t test or Wilcoxon's signed ranks test in the case of Turtle dove and Woodpigeon, are also indicated (* $p < 0.05$ and ** $p < 0.01$). The paired mean difference and associated s.d. together with the test statistic for each of the comparisons is given in Appendix B, Table B.1. All values are in *pqr* * 10^3 to reduce the number of decimals being reported.

Common name	Scientific name	<u>BASELINE</u>		<u>MAXIMIZE</u>		<u>RESILIENCE</u>	
		$\bar{X}$	s.d.	$\bar{X}$	s.d.	$\bar{X}$	s.d.
Turtle dove	<i>Streptopelia turtur</i>	-72.0	63.10	-134.6*	3.27	-81.0**	56.74
Yellow wagtail	<i>Motacilla flava</i>	-62.4	2.13	-60.3**	1.23	-61.8**	1.80
Starling	<i>Sturnus vulgaris</i>	-55.9	7.86	-57.6**	7.43	-55.5	7.76
Corn bunting	<i>Miliaria calandra</i>	-51.0	5.04	-53.6**	1.44	-52.7**	4.19
Linnet	<i>Carduelis cannabina</i>	-38.5	3.74	-40.1**	2.34	-39.6**	3.28
Yellowhammer	<i>Emberiza citrinella</i>	-32.1	10.43	-27.7**	9.32	-33.5**	10.05
Stock dove	<i>Columba oenas</i>	-24.2	10.28	-22.4**	8.43	-24.6	10.48
Rook	<i>Corvus frugilegus</i>	-24.1	6.11	-22.4**	5.91	-24.2	6.12
Skylark	<i>Alauda arvensis</i>	-19.5	6.62	-22.0**	1.43	-21.9**	5.31
Kestrel	<i>Falco tinnunculus</i>	-4.7	7.97	-4.5	12.07	-5.3*	8.18
Greenfinch	<i>Carduelis chloris</i>	-1.8	14.06	-8.5**	11.85	-3.6**	13.54
Lapwing	<i>Vanellus vanellus</i>	4.0	11.85	15.2**	12.88	3.4**	11.93
Whitethroat	<i>Sylvia communis</i>	14.1	5.29	14.8**	5.32	13.5**	5.24
Reed bunting	<i>Emberiza schoeniclus</i>	14.4	1.53	14.9**	1.02	14.1**	1.55
Goldfinch	<i>Carduelis carduelis</i>	15.6	2.04	15.1**	1.50	15.0**	1.71
Woodpigeon	<i>Columba palumbus</i>	24.1	5.07	30.7*	2.74	24.5*	5.07
Jackdaw	<i>Corvus monedula</i>	33.3	33.01	41.7**	37.47	34.0	33.82
Grey partridge	<i>Perdix perdix</i>	36.1	13.55	56.2**	13.57	36.1	13.50
Tree sparrow	<i>Passer montanus</i>	72.5	14.55	64.4**	9.89	68.2**	14.20
All species		-9.1	6.02	-10.6**	3.39	-10.3**	5.34

Maps of spatial patterns in *pgr* across Marston Vale under each scenario for three exemplar species are presented in Fig. 4. Equivalent maps for the remaining 16 species are available in Fig. B.2. Again, it is evident from Table 3 and Fig. 4 that there is considerable spatial variation at the 1 km scale in predicted *pgr* for individual species, with the extent varying between species and landscape configurations.




Fig. 4. Spatial variation in predicted annual *pgr* across Marston Vale for three exemplar species. Positive and negative values were coded using a blue and red coloured scale, respectively. The range and the breaks for each of the scales were determined to enhance visualisation. See Fig. B.2 for equivalent maps for the other 16 species.

3.4 Sensitivity analyses

Species responded predictably to changes in parameter estimates associated with the proportions of autumn or spring sown wheat, clean or weedy stubbles and different boundary types in the landscape. As expected, the mean predicted *pgr* was lower if the availability of spring sown cereal declined and *vice versa*. Similarly, the mean predicted *pgr* was lower if the specified proportion of stubble polygons characterised as weedy

was lower and *vice versa* (Table B.2). The mean absolute difference in predicted *pgr* when the proportion of autumn sown cereal was increased to 1 or reduced to 0.8, compared to that calculated using default parameter settings, was 0.0006 ± 0.00012 in both cases. The mean absolute difference in predicted *pgr* when the proportion of weedy stubbles was increased to 0.75 or reduced to 0.25 was 0.001 ± 0.0004 in both cases. These values are substantially lower than equivalent metric when switching from BASELINE to MAXIMIZE (0.0076 ± 0.0033) and between approximately 45% and 70% of that when switching between BASELINE and RESILIENCE (0.0014 ± 0.0005). Altering boundary feature characteristics had a greater impact on predicted *pgr* (Table B.2), with the mean absolute difference in predicted *pgr* of the three different configurations falling between 0.0024 ± 0.0008 and 0.0027 ± 0.0009 when compared to those calculated using the default parameter settings. Patterns of change in predicted *pgr* associated with each set of boundary characteristic parameters applied varied more between species, in line with the specific contribution of each boundary type to their functional space (Table B.2). For all but two species, the direction of change in predicted *pgr* when moving from BASELINE to MAXIMIZE or BASELINE to RESILIENCE remained consistent across all combinations of parameter estimates tested. For Kestrel, the predicted *pgr* decreased under MAXIMIZE if the proportion of spring sown wheat was set to 0% but increased under all other parameter combinations. The direction of change in predicted *pgr* of Grey partridge on moving to RESILIENCE varied across parameter combinations but these were small differences around an initial prediction of zero impact (Tables B.3 and B.4). Overall, changes in parameter estimates did not affect the results obtained when detecting differences between BASELINE, MAXIMIZE and RESILIENCE scenarios except for few species and parameter combinations (Tables B.3 and B.4); consistent results were obtained for all the parameter combinations when considering the combined *pgr* value for the 19 species (Tables B.3 and B.4).

4. Discussion

Our results indicate that the strategy adopted to deliver the UK's land based renewable energy targets can affect both gross bioenergy production and farmland bird population trends. This integration of biodiversity and bioenergy production assessments for a common set of scenarios for a defined area, alongside assessments of the effect of the same land use changes on the level of food, animal feed and fibre as more fully reported by Burgess *et al.* (2012), serves as a prototype of the model integration framework that is needed to allow policy makers to predict the economic and environmental impacts of different land use policies and to assess potential mitigation measures.

Each of the three bioenergy production scenarios examined here could, in the context of the Marston Vale, deliver the 2020 renewable energy production targets for transport fuel and heat (Table 2), albeit at a cost to food production. Predicted gross energy levels were higher under MAXIMIZE ($21 \text{ kWh p}^{-1} \text{ d}^{-1}$) than RESILIENCE ($12.3 \text{ kWh p}^{-1} \text{ d}^{-1}$) which itself was marginally greater than that for BASELINE ($11.3 \text{ kWh p}^{-1} \text{ d}^{-1}$). Note that whilst the predicted gross energy levels were highest under MAXIMIZE, it creates a potentially volatile portfolio of a single renewable energy production type, where failure of the wheat crop (e.g. through disease) could result in near total collapse of overall bioenergy and food production. Both MAXIMIZE and RESILIENCE were predicted to

have significant impacts on population trends of most farmland bird species but, despite substantial differences in resultant landscape composition, the response of the community as a whole to each scenario was small and broadly similar (Table 3). Interestingly, the reallocation of post harvest products to energy production once food demand had been met under BASELINE was predicted to deliver broadly equivalent levels of energy output to RESILIENCE, without the added detrimental impacts on farmland birds. It is important to note that our calculations for this post-production reallocation scenario did not take into account factors such as the likely reduction in soil carbon and nutrient levels, and hence long term crop yields, associated with annual removal of straw. Although they are therefore likely an oversimplification of long term effects, those calculations serve to highlight the potential contribution of alternative strategies, beyond direct changes in land use, for meeting renewable energy production targets.

Primarily, our results highlight the need for context dependent, species-level assessment of land use change impacts at a range of spatial scales including field, farm and landscape (Ekroos and Kuusaari, 2011; Robinson *et al.*, 2001; Schweiger *et al.*, 2005) and provide such a framework. It is clear that the relatively small and broadly equivalent detrimental impacts of MAXIMIZE and RESILIENCE on overall farmland bird trends mask substantial variations within and between individual species' responses to each scenario. Under MAXIMIZE, the small decline in mean *pqr* across species is a composite of declines across approximately half the species and increases across the others whilst that under RESILIENCE is underpinned by small declines across the majority of species, with few demonstrating a predicted positive response (Table 3). This suggests that the mechanism of impact of the two scenarios on farmland bird population trends may well be different (see below). Furthermore, many species exhibited positive predicted annual *pqr* in some squares even if their population trend across Marston Vale was predicted to be declining overall under a given scenario and *vice versa* (Fig. B.1). Given the context dependence of species' responses identified, it is important to note that the results presented here relate specifically to the implementation of each scenario in the current landscape of Marston Vale. Although there are substantial areas of lowland England with similar wheat and oilseed rape dominated agricultural landscapes to which our results are likely to be broadly applicable, the response of the farmland bird community to these land use scenarios in other regions should be assessed on a case-by-case basis. Furthermore, our analyses assume that the management of crops for bioenergy is the same as for food. If, for example, it becomes evident that crop management practices, such as rates of agrochemical application or sowing and harvesting dates, change as a result of switching from management for food to management for renewable energy, the habitat allocation algorithms used to quantify functional space would need revision.

Whilst providing a detailed discussion of the response of individual species to each scenario is not the main focus of this paper, the broadly similar overall response to the two contrasting land use change scenarios demands further examination. Variations in species' predicted responses to land use change can be attributed to the type and number of habitat types that contribute to each species' functional space, and the absolute and relative abundance of those habitats in each square in the current landscape and under each bioenergy production scenario. For example, a species that relies on a limited number of habitats could show low spatial variation if that habitat type is very dominant or very rare in the landscape but high spatial variation if that habitat is more patchily

distributed across the landscape. Under MAXIMIZE, all arable and grassland areas were planted to wheat. Whilst this led to an increase in the overall arable area, it greatly reduced landscape heterogeneity and the area of spring cropping (Table 1). This simplification of the landscape is reflected in the generally reduced levels of spatial variation in predicted *pgr* across species under MAXIMIZE (Table 3). Broadly speaking, this arable-dominated landscape homogenisation epitomises continued agricultural intensification, representing an extension of many of the land use changes that have occurred in UK lowland agricultural landscapes over recent decades (Stoate *et al.*, 2001, Benton *et al.*, 2003). As a result, the majority of species predicted to respond detrimentally to MAXIMIZE, such as Skylark (*Alauda arvensis*), Corn bunting (*Miliaria calandra*) and Linnet (*Carduelis carduelis*), are those currently declining as a result of past land use changes associated with agricultural intensification (Table 3; Newton, 2004). Indeed, the greatest predicted detrimental impact of MAXIMIZE was on Turtle dove (*Streptopelia turtur*), a species previously identified as showing a strong preference for non-cereal areas in lowland England (Browne and Aebischer, 2003). Our results support and extend findings by Engel *et al.* (2012) who used a spatial explicit, single species model to demonstrate the potential of bioenergy production to detrimentally impact Skylark populations through reduced habitat heterogeneity.

Under RESILIENCE, the areas assigned to each crop did not change substantially from BASELINE but there were reductions in both total arable area and area of spring cropping as two novel crops, miscanthus and short rotation coppice, replaced areas of fallow and spring sown oilseed rape (Table 1). Due to the structural characteristics of these two novel crops, they are predicted to contribute little to the functional space of many of the farmland specialists included in the Farmland Bird Index (Anderson *et al.*, 2004; Sage *et al.*, 2006, 2010). Given our focus on farmland bird species specifically, the predicted detrimental impact of RESILIENCE is likely driven largely through reductions in functional space associated with a net reduction in total arable area, akin to the detrimental impacts of land abandonment of farmland bird populations (e.g. Butler *et al.*, 2010). This conclusion is supported by the fact that both currently declining and increasing species were predicted to respond negatively to the land use changes associated with RESILIENCE (Table 3). In line with this, it is important to emphasise that our assessment of the biodiversity impacts of each scenario is based on the predicted response of the Farmland Bird Index species, with any inferences of the effects on wider farmland biodiversity based on the broadly accepted assumption that bird population trends are indicative of wider biodiversity health (Gregory *et al.*, 2003). Whilst a decrease in the cropped area of an agricultural landscape may decrease the functional space for farmland species it may also increase the opportunities for more generalist species or those specialised to other ecosystems. For example, although short rotation coppice is expected to reduce the functional space for farmland specialists such as Skylark and Lapwing *Vanellus vanellus*, which require more open vegetation, it can provide functional space for species associated with scrubland and early succession forests (Sage *et al.*, 2006). Such observations suggest that a full assessment of the biodiversity impacts of land use change needs more than a focus on solely farmland species whilst, where relevant, taking into account both local and national conservation priorities.

Our calculations of functional space in each 1 km square are also dependent on a number of assumptions. These include the categorization of boundary features based on a subsample of squares, the use of regional agricultural census data to infer winter crop

cover types, and the use of the national WFBS data to assign proportions of weedy and non-weedy stubbles. In any modelling exercise, assumptions are needed and we believe that the assumptions we have made are broadly representative and that there is no directional bias, particularly as we undertake relative comparisons between scenarios. This is supported by the results of our sensitivity analyses which showed a) that the predicted response of farmland birds to changes in land use associated with each scenario were greater than those associated with changing the proportion of spring sown crop, weedy stubble or boundary characteristics in the landscape and b) that the direction of change in species' predicted *pqr* between scenarios was the same for all combinations of parameters tested in all but a few cases. Unsurprisingly, the relative difference between species' responses to scenario-dictated land use change and parameter-dictated land use change was much greater for MAXIMIZE than RESILIENCE. As discussed above, MAXIMIZE led to much greater changes in landscape composition than RESILIENCE so the resultant land use changes induced by altering parameter estimates in the sensitivity analyses were closer in scale to those which occurred under RESILIENCE and the consequent impact on predicted *pqr* more similar. If more site-specific data were available for the above characteristics, the habitat allocation algorithms could be readily adapted to accommodate them. Note that assumptions relating specifically to the development of the functional space models are discussed in detail elsewhere (Butler and Norris, 2013) but one key difference between their approach and that reported here deserves mention. Here, transects are spatially positioned independently with respect to land use but for BBS this is usually not the case as observers tend to use paths or field boundaries when undertaking surveys. Thus, whilst delivering a more realistic assessment of the availability of functional space associated with boundary features, the quantity calculated using the GIS based approach reported here is likely to be lower than that if a real BBS habitat assessment had been undertaken. This will influence the predicted *pqr* presented for each species according to the strength and direction of any relationship between local population trend and functional space provided by boundary features. However, given that this difference in methodology is reflected in functional space and predicted *pqr* calculations across all scenarios, that the sensitivity analyses revealed a relatively limited influence of changes in boundary feature availability on predicted *pqr* and that our conclusions are based primarily on relative comparisons in predicted *pqr*, these differences are unlikely to have unduly biased our results.

The application of our approach is not limited to renewable energy based land use change and developing a GIS based framework facilitates the integration of BirdMod with other land use based models for a range of ecosystem services (e.g. Burgess *et al.*, 2012; Carver *et al.*, 2011; Smith *et al.*, 2010; Kareiva *et al.*, 2011). Furthermore, this approach is not solely restricted to predicting the impacts of land use changes. It could, for example, also be used to explore alternative mitigation strategies designed to offset the detrimental impacts of any land use change identified or to assess the likely benefits of proposed conservation management. However, we recognize that there are still limitations that we intend to address in the future. Importantly, BirdMod requires a digitized version of the land uses within the study area as input data. Digitization of all the parcels within the area where the model is to be applied can be time consuming and may prove impractical for managers and researchers. Moreover, there is always an intrinsic error in the identification of structural land use types from aerial photography assessment. One option is to modify BirdMod and the underlying functional space

responses to use input data from a national data source such as the Land Cover Map derived from the UK Countryside Survey (Morton *et al.*, 2011) to describe structural parcels and the boundary characteristics. However this in turn creates new inaccuracies and uncertainties because of the way in which land cover maps are developed and their spatial and temporal resolution is likely to limit the quantification of the functional space delivered by, for example, linear features. Furthermore, functional space models have so far only been developed for farmland bird species but previous work (Butler *et al.*, 2009) suggests that it should be possible to quantify functional spaces for other taxonomic groups and ecosystems and to develop the equivalent models.

5. Conclusions

A variety of contrasting land use strategies could be employed to meet UK and European renewable energy production targets and the UK Government (2011) seeks to promote “an integrated approach to managing the natural environment, particularly at the landscape scale” when deciding on the most appropriate approach. To support this, land planners and managers need access to tools and models that can predict the impact of alternative strategies on the stocks and flow of ecosystem goods and services, including biodiversity. Whilst there are an increasing number of tools to describe the interactions between land use and food, feed, fibre and fuel production, it has proved more difficult to develop tools to describe the effects on biodiversity; developing the capability to model context dependent biodiversity responses to land use change is therefore fundamental to the development of the evidence base needed to guide policy implementation decisions. We believe BirdMod, and the wider conceptual framework that underpins it, offers that capability and, given the opportunity it provides to explore potential mitigation approaches for each policy implementation strategy, propose that it could play a key role in ensuring renewable energy and other land use change driving policies are delivered in a sustainable manner.

Acknowledgements

The research described was funded by a Natural Environment Research Council (NERC) grant (reference NE/H010432/1). The group wishes to express their thanks to those who have commented and made positive suggestions about the work and all the stakeholders, colleagues and students who have assisted in the process.

Appendix A. Algorithms applied for primary and secondary habitat classifications

BBS classifies primary and secondary habitats in each transect section. The algorithms used to replicate this classification are summarized in Figs. A.1 and A.2 for the primary habitat and Fig. A.4 for the secondary habitat. These algorithms were implemented in an Excel platform and followed the guidelines for the UK Defra Agricultural and Horticultural Census data 2005-2008 (www.defra.gov.uk).

The primary habitat is defined by four levels named P-L1, P-L2, P-L3 and P-L4. P-L1 is classified into WOODLAND, FARMLAND, HUMAN or WATER based on the dominant land use identified in the digitized polygons from the aerial photography as described in the methodology section. Each of the P-L1 classes follows a different set of habitat allocation algorithms to identify P-L2, P-L3 and P-L4 (Figs. A.1 and A.2).

Similarly, the secondary habitat also has four levels coded (S-L1, S-L2, S-L3 and S-L4), with classification based on the sequence of habitat allocation algorithm in Fig. A.3. The stochastic component in the model is introduced by the RAND() variable, where RAND represents a randomly generated numbers between 0 and 1, independently identified for each tier of habitat classification.

Fig. A.4 summarizes the algorithm applied to each winter land use classification of each polygon larger than 0.3 ha. Winter habitat is defined by 3 levels (i.e. W-L1, W-L2 and W-L3). Polygons with any other summer land use type classification than these included in Fig. A.4 were not assigned a winter habitat code as only farmland habitats were recorded in Winter Farmland Bird Survey.




Fig. A.1. Primary habitat allocation algorithm applied when the first tier of classification (P-L1) was FARMLAND. B is the total length of hedges within the 1 km square; D is the total area of disturbed habitat within the transect section; RAND() is a randomly generated number between 0 and 1, with the associated subscript number identifying the tier within the four level hierarchical BBS habitat code structure; P-L1, P-L2, P-L3 and P-L4 represent the four primary habitat levels; and WHT, OSR, CRP and FLW are the land uses coded as specified in Table 1.




Fig. A.2. Primary habitat allocation algorithm applied when the first tier of classification (P-L1) was WOODLAND, WATER or HUMAN. B is the total length of hedges within the 1 km square; D is the total area of disturbed habitat within the transect section; RAND() is a randomly generated number between 0 and 1, with the associated subscript number identifying the tier within the four level hierarchical BBS habitat code structure; P-L1, P-L2, P-L3 and P-L4 represent the four primary habitat levels; and TR, CM, LD, OTH and URB are the land uses coded as specified in Table 1.




Fig. A.3. Secondary habitat allocation algorithm applied. B is the total length of hedges within the 1 km square; D is the total area of disturbed habitat within the transect section; RAND() is a randomly generated number between 0 and 1, with the associated subscript number identifying the tier within the four level hierarchical BBS habitat code structure; P-L1, P-L2, P-L3 and P-L4 represent the four primary habitat levels; S-L1, S-L2, S-L3 and S-L4 represent the four secondary habitat levels; and WHT, OSR, CRP and FLW are the land uses coded as specified in Table 1.




Fig. A.4. Winter habitat allocation algorithm applied. GRS, WDL, WSC, OSR, CRP, BRL, WHT are the land use (LU) classes as coded in Table 1; RAND() is a randomly generated number between 0 and 1, with the associated subscript number identifying the tier within the three level hierarchical WFBS habitat code structure; W-L1, W-L2 and W-L3 represent the three winter habitat levels.

Appendix B. Response of individual species to the three land use configurations

Table B.1. Results from the paired t test analysis. $\bar{X}$ refers to the mean paired difference between the BASELINE and MAXIMIZE (M) scenarios or the BASELINE and RESILIENCE (R) scenarios, with positive values indicating a higher predicted annual *pqr* and negative values a lower predicted annual *pqr*. “s.d.” and “test results” stand for the standard deviation of the mean paired difference and the t values from the t test (n=142, df=141). For the case of turtle dove and woodpigeon, the “test results” show the outputs of the Wilcoxon’s signed ranks test (n = number of signed ranks and z = z-ratio). All the mean and standard deviation are in *pqr* * 10³ to reduce the number of decimals being reported.

Common name	Scientific name	Paired $\bar{X}$ (M)	s.d. (M)	Test results (M)	Paired $\bar{X}$ (R)	s.d. (R)	Test results (R)
Turtle dove	<i>Streptopelia turtur</i>	62.56	64.11	n=140, z=10.11	8.97	27.17	3.93
Yellow wagtail	<i>Motacilla flava</i>	-2.10	2.28		-0.59	1.29	-5.45
Starling	<i>Sturnus vulgaris</i>	1.71	3.75	5.43	-0.38	3.49	-1.30
Corn bunting	<i>Miliaria calandra</i>	2.56	4.64	6.56	1.73	2.91	7.08
Linnet	<i>Carduelis cannabina</i>	1.60	2.39	7.95	1.09	1.99	6.55
Yellowhammer	<i>Emberiza citrinella</i>	-4.42	5.32	-9.89	1.40	3.33	5.00
Stock dove	<i>Columba oenas</i>	-1.82	6.16	-3.51	0.38	3.67	1.23
Rook	<i>Corvus frugilegus</i>	-1.74	1.65	-12.61	0.07	0.71	1.22
Skylark	<i>Alauda arvensis</i>	2.51	6.56	4.56	2.46	4.85	6.04
Kestrel	<i>Falco tinnunculus</i>	-0.19	10.97	-0.20	0.67	3.54	2.26
Greenfinch	<i>Carduelis chloris</i>	6.71	7.02	11.39	1.81	3.57	6.04
Lapwing	<i>Vanellus vanellus</i>	-11.22	7.58	-17.63	0.58	2.11	3.26
Whitethroat	<i>Sylvia communis</i>	-0.76	1.52	-5.99	0.55	1.75	3.77
Reed bunting	<i>Emberiza schoeniclus</i>	-0.55	1.40	-4.66	0.26	0.58	5.31
Goldfinch	<i>Carduelis carduelis</i>	0.51	1.18	5.12	0.64	1.41	5.39
Woodpigeon	<i>Columba palumbus</i>	-6.52	4.88	n=142, z=10.16	-0.38	1.91	-2.39
Jackdaw	<i>Corvus monedula</i>	-8.49	18.37		-0.79	5.09	-1.84
Grey partridge	<i>Perdix perdix</i>	-20.10	11.74	-20.40	-0.03	2.72	-0.14
Tree sparrow	<i>Passer montanus</i>	8.14	12.04	8.05	4.34	9.93	5.20
All species		1.49	4.55	3.91	1.20	2.62	5.44

Table B.2. Mean predicted annual *pgr*, across 10 iterations, for all species in the Farmland Bird Index for *BASELINE* land use under the default parameter settings (*Default*) and the difference between this and the predicted annual *pgr* for the same land use under each parameter combination tested in the sensitivity analyses. All values are in *pgr* * 10³ to reduce the number of decimals being reported. Boundary types – 1: Hedgerow with trees; 2: Hedgerow without trees; 3: Treeline; 4: No vertical structure. Within this context, the boundary characteristics have the following values (a) 25%, 25%, 25%, 25%, (b) 50%, 50%, 0%, 0% and (c) 0%, 0%, 50%, 50%.

Common name	Scientific name	Default	Sowing date		Stubble condition		Boundary characteristics		
			Autumn 100%	Autumn 80%	Weedy 25%	Weedy 75%	(a)	(b)	(c)
Turtle dove	<i>Streptopelia turtur</i>	-72.0	0.2	0.1	0.3	0.3	0.3	3.7	-3.5
Yellow wagtail	<i>Motacilla flava</i>	-62.4	0.6	-0.5	0.1	0.0	0.7	1.6	-0.1
Starling	<i>Sturnus vulgaris</i>	-55.9	-0.3	0.1	-0.3	0.1	-7.5	-14.1	-0.5
Corn bunting	<i>Miliaria calandra</i>	-51.0	-1.0	0.9	-0.5	0.6	-0.1	-0.1	0.0
Linnet	<i>Carduelis cannabina</i>	-38.5	-0.5	0.4	-0.3	0.4	-0.1	0.6	-0.7
Yellowhammer	<i>Emberiza citrinella</i>	-32.1	-1.0	1.0	-0.8	0.9	-3.0	0.9	-6.6
Stock dove	<i>Columba oenas</i>	-24.2	-0.4	0.8	-1.0	1.5	7.6	7.8	7.8
Rook	<i>Corvus frugilegus</i>	-24.1	0.1	-0.1	0.3	-0.3	-1.0	-1.0	-1.1
Skylark	<i>Alauda arvensis</i>	-19.5	-1.6	1.4	-1.1	1.4	-1.6	-1.4	-1.4
Kestrel	<i>Falco tinnunculus</i>	-4.7	-0.5	0.4	0.0	0.0	0.0	0.0	0.0
Greenfinch	<i>Carduelis chloris</i>	-1.8	-0.8	0.9	-1.5	1.8	0.7	-0.1	2.0
Lapwing	<i>Vanellus vanellus</i>	4.0	0.8	-0.8	-0.1	0.0	0.0	0.0	0.1
Whitethroat	<i>Sylvia communis</i>	14.1	0.0	0.0	0.0	0.0	-2.8	0.4	-6.1
Reed bunting	<i>Emberiza schoeniclus</i>	14.4	-0.2	0.2	-0.3	0.3	-0.2	0.0	-0.3
Goldfinch	<i>Carduelis carduelis</i>	15.6	0.0	0.0	-0.1	0.1	1.0	0.5	1.4
Woodpigeon	<i>Columba palumbus</i>	24.1	0.4	-0.2	0.7	-0.8	-2.3	-5.5	0.7
Jackdaw	<i>Corvus monedula</i>	33.3	0.0	-0.2	-0.7	0.6	-2.6	-3.0	-3.1
Grey partridge	<i>Perdix perdix</i>	36.1	1.6	-1.6	2.9	-3.2	0.8	0.6	0.4
Tree sparrow	<i>Passer montanus</i>	72.5	-1.6	2.0	-7.7	8.5	-12.9	-9.0	-14.7
All species		-9.1	-0.2	0.3	-0.5	0.6	-1.2	-0.9	-1.4

Table B.3. Predicted annual *pgr* for *MAXIMIZE* for each parameter combination tested, where Default - default parameter settings; C1 – Percentage autumn sown wheat 100%; C2 – Percentage autumn sown wheat 80%; C3 – Percentage weedy stubbles 25%; C4 – Percentage weedy stubbles 75%; C5, C6, C7 – Percentage of boundary features classified as hedgerow with trees, hedgerow without trees, tree line with no hedge or no vertical structure set at 25%, 25%, 25%, 25%; 50%, 50%, 0%, 0%; 0%, 0%, 50%, 50% respectively. For comparison, the predicted *pgr* for *BASELINE* under default parameter settings is also shown for each species. All values are in *pgr* * 10³ to reduce the number of decimals being reported. Results for the square level paired t test or Wilcoxon's signed rank test in the case of turtle dove and woodpigeon are also indicated (* p<0.05 and ** p<0.01).

Common name	Scientific name	<i>BASELINE</i>	<i>MAXIMIZE</i>							
		Default	Default	C1	C2	C3	C4	C5	C6	C7
Turtle dove	<i>Streptopelia turtur</i>	-72.0	-134.6*	-134.6**	-134.5**	-134.7**	-134.5**	-134.7**	-131.0**	-138.5**
Yellow wagtail	<i>Motacilla flava</i>	-62.4	-60.3**	-59.2**	-61.4**	-60.3**	-60.3**	-60.0**	-59.6**	-60.3**
Starling	<i>Sturnus vulgaris</i>	-55.9	-57.6**	-57.4**	-57.0*	-57.1**	-57.4**	-64.9**	-72.2**	-57.9**
Corn bunting	<i>Miliaria calandra</i>	-51.0	-53.6**	-56.0**	-51.0*	-53.7**	-53.3**	-53.5**	-53.5**	-53.5**
Linnet	<i>Carduelis cannabina</i>	-38.5	-40.1**	-41.3**	-38.9**	-40.2**	-40.0**	-40.1**	-39.4**	-40.8**
Yellowhammer	<i>Emberiza citrinella</i>	-32.1	-27.7**	-30.2**	-25.2**	-28.1**	-27.4**	-31.6**	-26.4**	-36.5**
Stock dove	<i>Columba oenas</i>	-24.2	-22.4**	-23.6**	-21.4	-23.1**	-22.1	-14.3**	-14.4**	-14.4**
Rook	<i>Corvus frugilegus</i>	-24.1	-22.4**	-22.2**	-22.7**	-22.3**	-22.5**	-23.5**	-23.6**	-23.6**
Skylark	<i>Alauda arvensis</i>	-19.5	-22.0**	-25.6**	-18.2	-22.5**	-21.5**	-22.4**	-22.4**	-22.4**
Kestrel	<i>Falco tinnunculus</i>	-4.7	-4.5	-5.7	-3.1	-4.3	-4.4	-4.3	-4.3	-4.3
Greenfinch	<i>Carduelis chloris</i>	-1.8	-8.5**	-10.5**	-6.4**	-9.2**	-7.8**	-7.5**	-8.5**	-6.4**
Lapwing	<i>Vanellus vanellus</i>	4.0	15.2**	17.1**	13.2**	15.1**	15.2**	15.2**	15.1**	15.1**
Whitethroat	<i>Sylvia communis</i>	14.1	14.8**	14.9**	14.8**	14.8**	14.8**	11.7**	15.3**	8.2
Reed bunting	<i>Emberiza schoeniclus</i>	14.4	14.9**	14.4	15.5**	14.8**	15.0*	14.5*	15.0**	14.0
Goldfinch	<i>Carduelis carduelis</i>	15.6	15.1**	15.1**	15.1**	15.1**	15.2**	16.1**	15.7**	16.5**
Woodpigeon	<i>Columba palumbus</i>	24.1	30.7*	31.4**	30.0**	31.0**	30.4**	27.9**	24.7**	31.3**
Jackdaw	<i>Corvus monedula</i>	33.3	41.7**	41.3**	42.2**	41.5**	42.0**	38.8**	38.9**	38.9**
Grey partridge	<i>Perdix perdix</i>	36.1	56.2**	59.9**	52.1**	57.5**	54.9**	56.4**	56.3**	56.3**
Tree sparrow	<i>Passer montanus</i>	72.5	64.4**	60.7**	69.3**	61.3**	68.3**	52.3**	54.8**	50.7**
All species		-9.1	-10.6**	-11.1**	-9.9**	-10.8**	-10.3**	-11.8**	-11.5**	-12.0**

Table B.4. Predicted annual *pgr* for *RESILIENCE* for each parameter combination tested, where Default - default parameter settings; C1 – Percentage autumn sown wheat 100%; C2 – Percentage autumn sown wheat 80%; C3 – Percentage weedy stubbles 25%; C4 – Percentage weedy stubbles 75%; C5, C6, C7 – Percentage of boundary features classified as hedgerow with trees, hedgerow without trees, tree line with no hedge or no vertical structure set at 25%, 25%, 25%, 25%; 50%, 50%, 0%, 0%; 0%, 0%, 50%, 50% respectively. For comparison, the predicted *pgr* for *BASELINE* under default parameter settings is also shown for each species. All values are in *pgr* * 10³ to reduce the number of decimals being reported. Results for the square level paired t test or Wilcoxon's signed rank test in the case of turtle dove and woodpigeon are also indicated (* p<0.05 and ** p<0.01).

Common name	Scientific name	<i>BASELINE</i>	<i>RESILIENCE</i>							
		Default	Default	C1	C2	C3	C4	C5	C6	C7
Turtle dove	<i>Streptopelia turtur</i>	-72.0	-81.0**	-81.0**	-81.1**	-81.0**	-80.9**	-81.3**	-77.8**	-84.4**
Yellow wagtail	<i>Motacilla flava</i>	-62.4	-61.8**	-61.3**	-62.2**	-61.8**	-61.8**	-61.3	-60.8	-61.9**
Starling	<i>Sturnus vulgaris</i>	-55.9	-55.5	-55.5*	-54.9*	-55.4*	-55.5	-62.3**	-68.9**	-55.6*
Corn bunting	<i>Miliaria calandra</i>	-51.0	-52.7**	-53.6**	-51.7**	-53.1**	-52.3**	-52.7**	-52.7**	-52.7**
Linnet	<i>Carduelis cannabina</i>	-38.5	-39.6**	-40.0**	-39.1**	-39.9**	-39.3**	-39.6**	-38.9**	-40.3**
Yellowhammer	<i>Emberiza citrinella</i>	-32.1	-33.5**	-34.5**	-32.5**	-34.0**	-32.9**	-36.4**	-32.6**	-39.9**
Stock dove	<i>Columba oenas</i>	-24.2	-24.6	-24.9	-23.9	-25.5	-23.6*	-17.3	-17.5**	-17.3*
Rook	<i>Corvus frugilegus</i>	-24.1	-24.2	-24.1	-24.3*	-24.0*	-24.4	-25.3	-25.3	-25.2
Skylark	<i>Alauda arvensis</i>	-19.5	-21.9**	-23.7**	-20.3**	-22.9**	-21.0**	-23.0**	-23.0**	-22.9**
Kestrel	<i>Falco tinnunculus</i>	-4.7	-5.3*	-5.7	-4.7	-5.3	-5.2*	-5.2	-5.2	-5.2
Greenfinch	<i>Carduelis chloris</i>	-1.8	-3.6**	-4.4**	-2.8**	-4.6**	-2.3**	-2.8**	-3.7**	-1.7**
Lapwing	<i>Vanellus vanellus</i>	4.0	3.4**	4.1**	2.5**	3.4**	3.3**	3.3**	3.3**	3.3**
Whitethroat	<i>Sylvia communis</i>	14.1	13.5**	13.5**	13.5**	13.5**	13.5**	10.6**	13.9**	7.5**
Reed bunting	<i>Emberiza schoeniclus</i>	14.4	14.1**	13.9**	14.3**	13.9**	14.3**	13.9**	14.1**	13.8**
Goldfinch	<i>Carduelis carduelis</i>	15.6	15.0**	14.9**	15.0**	14.9**	15.0**	15.8**	15.5**	16.3**
Woodpigeon	<i>Columba palumbus</i>	24.1	24.5*	24.8	24.2	24.9	23.9**	22.1**	19.1**	25.0
Jackdaw	<i>Corvus monedula</i>	33.3	34.0	33.7	34.3**	33.8**	34.7*	31.9*	31.4*	31.4*
Grey partridge	<i>Perdix perdix</i>	36.1	36.1	37.7	34.5	38.0**	33.9**	36.4*	36.3	36.2
Tree sparrow	<i>Passer montanus</i>	72.5	68.2**	66.2**	70.1**	63.1**	73.7**	56.3**	58.8**	55.1**
All species		-9.1	-10.3**	-10.5**	-10.0**	-10.6**	-9.8**	-11.4**	-11.3**	-11.5**




Fig. B.1. Box plot (median, quartiles and non outlier range) of the predicted annual *pgr* for each species and scenario. Species are coded as: CB - Corn bunting; GF - Goldfinch; GE - Greenfinch; GP - Grey partridge; JD - Jackdaw; KT - Kestrel; LW - Lapwing; LN - Linnet; RB - Reed bunting; RK - Rook; SK - Skylark; SG - Starling; SD - Stock dove; TS - Tree sparrow; TD - Turtle dove; WT - Whitethroat; WP - Woodpigeon; YW - Yellow wagtail; YH - Yellowhammer.




Fig. B.2. Spatial variation in predicted annual *pgr* across Marston Vale for 16 species not included in Fig. 4. Positive and negative values were coded using a blue and red coloured scale, respectively. The range and the breaks of the scale were determined to enhance visualization.

References

- Anderson GQA, Haskins LR, Nelson SH. The effects of bioenergy crops on farmland birds in the United Kingdom: A review of current knowledge and future predictions. In: Parris K, Poincet T, editors. Biomass and Agriculture: sustainability, markets and policies. OECD, Paris; 2004. p. 199-218.
- Benton, TG, Vickery, JA, Wilson, JD. Farmland biodiversity: is habitat heterogeneity the key? *Trends Ecol Evol* 2003;18:182-188.
- Browne SJ, Aebischer NJ. Habitat use, foraging ecology and diet of Turtle Doves *Streptopelia turtur* in Britain. *Ibis* 2003;145:572-582.
- Burgess PJ, Rivas Casado M, Gavu J, Mead M, Cockerill T, Lord R, van der Horst D, Howard DC. A framework for reviewing the trade offs between renewable energy, food, feed and wood production at a local level. *Renew Sust Energ Rev* 2012;16:129-142.
- Butchart SHM, Walpole M, Collen B, van Strien A, Scharlemann JPW, Almond REA, Baillie JEM, Bomhard B, Brown C, Bruno J, Carpenter KE, Carr GM, Chanson J, Chenery AM, Csirke J, Davidson NC, Dentener F, Foster M, Galli A, Galloway JN, Genovesi P, Gregory RD, Hockings M, Kapos V, Lamarque JF, Leverington F, Loh J, McGeoch MA, McRae L, Minasyan A, Morcillo MH, Oldfield TEE, Pauly D, Quader S, Revenga C, Sauer JR, Skolnik B, Spear D, Stanwell-Smith D, Stuart SN, Symes A, Tierney M, Tyrrell TD, Vie JC, Watson R. Global Biodiversity: Indicators of Recent Declines. *Science* 2010;328:1164-1168.
- Butler SJ, Boccaccio L, Gregory RD, Vorisek P, Norris K. Quantifying the impacts of land use change to European farmland bird populations. *Agric Eco Env* 2010;137:348-357.
- Butler SJ, Bradbury RB, Whittingham MJ. Stubble height affects the use of stubble fields by farmland birds. *J Appl Ecol* 2005;42:469-476.
- Butler SJ, Brooks DR, Feber RE, Storkey J, Vickery JA, Norris K. A crosstaxonomic index for quantifying the health of farmland biodiversity. *J Appl Ecol* 2009;46:1154-1162.
- Butler SJ, Norris K. Functional space and the population dynamics of birds in agroecosystems. *Agr Ecosyst Environ* 2013;164:200-208.
- Carver S, Comber A, McMorran R, Nutter S. A GIS model for mapping spatial patterns and distribution of wild land in Scotland. *Landscape urban plan* 2011;104:395-409.
- Committee on Climate Change. Bioenergy Review. Committee on Climate Change. London; 2011.
- Ekroos J, Kuusaari M. Landscape context affects the relationship between local and landscape species richness of butterflies in seminatural habitats. *Ecography* 2011;35:232-239.
- Engel J, Huth A, Frank K. Bioenergy production and Skylark (*Alauda arvensis*) population abundance – a modelling approach for the analysis of land use change impacts and conservation options. *GCB Bioenergy* 2012;4:713–727.
- European Commission. An EU Strategy for Biofuels. Commission of the European Communities; 2006.
- European Commission. Directive 2009/28/EC of the European parliament and the Council; 2009.
- European Commission. Proposal for a Directive amending Directives 98/70/EC and 2009/28/EC. COM(2012) 595 final 2012/0288 (COD); 2012.

- EurObserv'ER. The State of Renewable Energies in Europe. 12th EurObser'ER report. Paris; 2013.
- Fargione F, Hill J, Tilman P, Polasky S, Hawthorne P. Land clearing and the biofuel carbon debt. *Science* 2008;319:1235-1238
- Fitzherbert EB, Struebig MJ, Morel A, Danielsen F, Brühl CA, Donald PF, Phalan B. How will oil palm expansion affect biodiversity? *Trends Ecol Evol* 2008;23:538-545.
- Gallagher E. The Gallagher Review of the indirect effects of biofuels production. Renewable Fuels Agency. St Leonards on Sea; 2008.
- Gillings S, Wilson AM, Conway GJ, Vickery JA, Fuller RJ. Distribution and abundance of birds and their habitats within the lowland farmland of Britain in winter. *Bird Study* 2008;55:8-22.
- Gregory RD, Noble D, Field RH, Marchant JH, Raven M, Gibbons DW. Using birds as indicators of biodiversity. *Ornis Hungarica* 2003;12–13:11–24.
- Grubb M, Butler L, Twomey P. Diversity and security in UK electricity generation: the influence of low carbon objectives. *Energ policy* 2006;34(18): 4050-4062.
- Howard DC, Burgess PJ, Butler SJ, Carver SJ, Cockerill T, Coleby AM, Guohui G, Goodier CI, van der Horst D, Hubacek K, Lord R, Mead A, Rivas Casado M, Wadsworth RA, Scholefield, P. Energyscapes: linking the energy system and ecosystem services in real landscape. *Biomass Bioenerg* 2013;55:17-26.
- Kareiva P, Tallis H, Ricketts TH, Daily GC, Polasky S. *Natural Capital: Theory and Practice of Mapping Ecosystem Services*. Oxford University Press; 2011.
- Morton D, Rowland C, Wood C, Meek L, Marston C, Smith G, Wadsworth R, Simpson IC. Final Report for LCM2007 - the new UK land cover map, Countryside Survey Technical Report No 11/07 NERC. Centre for Ecology and Hydrology; 2011.
- Murphy R, Woods J, Black M, McManus M. Global developments in the competition for land from biofuels. *Food Policy* 2011;36:S52-S61.
- Newton I. The recent declines of farmland bird populations in Britain: an appraisal of causal factors and conservation actions. *Ibis* 2004;146:579-560.
- Petersen JE, Elbersen B, Wiesenthal T, Feehan J, Eppler U. Estimating the environmentally compatible bioenergy potential from agriculture. European Environment Agency. Copenhagen; 2007.
- Risely K, Renwick AR, Dadam D, Eaton MA, Johnston A, Baillie SR, Musgrove AJ, Noble DG. The Breeding Bird Survey 2010. BTO Research Report 597. British Trust for Ornithology. Thetford; 2011.
- Robinson RA, Wilson JD, Crick HQP. The importance of arable habitat for farmland birds in grassland landscapes. *J Appl Ecol* 2001;38:1059-1069.
- Rounsevell MDA, Annetts JE, Audsley E, Mayr T, Reginster L. Modelling the spatial distribution of agricultural land use at the regional scale. *Agric Ecosyst Environ* 2003;95:465-479.
- Sage R, Cunningham M, Boatman N. Birds in willow short rotation coppice compared to other arable crops in central England and a review of bird census data from energy crops in the UK. *Ibis* 2006;148:184–197.
- Sage R, Cunningham M, Haughton AJ, Mallott MD, Bohan DA, Riche A, Karr A. The environmental impacts of biomass crops: use by birds of miscanthus in summer and winter in southwestern England. *Ibis* 2010;152:487-499.
- Schweiger O, Maelfit JP, van Wingerden W, Hendrickx F, Billeter R, Speelmans M, Augenstein I, Aukema B, Aviron S, Bailey D, Bukacek R, Burel F, Diekötter F,

- Dirksen J, Frenzel M, Herzog F, Liira J, Roubalova M, Butger R. Quantifying the impact of environmental factors on arthropod communities in agricultural landscapes across organizational levels and spatial scales. *J Appl Ecol* 2005;42:1129-1139.
- Smith J, Gottschalk P, Bellarby J, Chapman S, Lilly A, Towers W, Bell J, Coleman K, Nayak D, Richards M, Hillier J, Flynn H, Wattenbach M, Aitkenhead M, Yeluripati J, Farmer J, Milne R, Thomson A, Evans C, Whitmore A, Falloon P, Smith P. Estimating changes in Scottish soil carbon stocks using ECOSSE. I. Model description and uncertainties. *Climate Res* 2010;45:179-192.
- Stoate C, Boatman ND, Borralho RJ, Rio Carvalho C, de Snoo GR, Eden P. Ecological impacts of agricultural intensification in Europe. *J Environ manage* 2001;63:337-365.
- UK Government. The Natural Choice: Securing the Value of Nature. 2011; London: The Stationary Office. <http://www.official-documents.gov.uk/document/cm80/8082/8082.pdf>. Accessed 3 December 2013.
- Whittingham MJ, Devereux CL. Changing grass height alters foraging site selection by wintering farmland birds. *Basic Appl Ecol* 2008;9:779-788.



DSPACE

<https://dspace.org/>

Predicting the impacts of bioenergy production on farmland birds

Rivas Casado, Monica; Mead, Andrew; Burgess, Paul J.; Howard, David C.; Butler, Simon J.

2014-01-21

Elsevier

<http://dx.doi.org/10.1016/j.scitotenv.2013.12.080>.

Rivas Casado M, Mead A, Burgess PJ, et al., (2014) Predicting the impacts of bioenergy production on farmland birds. Science of The Total Environment, Volume 476-477, April 2014, pp. 7-19