

Relative to the NPK plot, the yield-scaled N2O+NO emission was reduced by
32% and 44% in the organic and bio-organic plots, respectively
Organic Bio-organic
EFN2O = 1.34; 1.12
EFNO = 0.09; 0.03
NPK
EFN2O = 2.08
EFNO = 0.98
Cattle manure
Urea
N2O NO
N2O NO
N2
NO3-
NH4+
Denitrification
Nitrification
Organic N
CO(NH2)2

Geoderma, Volume 383, February 2021, Article number 114730
DOI:10.1016/j.geoderma.2020.114730

Type of paper: Regular papers

Number of tables: 2

Number of figures: 4

Number of words: 9248

Number of text pages: 39

Soil N-oxide emissions decrease from intensive greenhouse vegetable fields by substituting synthetic N fertilizer with organic and bio-organic fertilizers

Yajun Geng[a], Jinyang Wang[a,b*], Zhirong Sun[a], Cheng Ji[a], Mengyuan Huang[a], Yihe Zhang[a], Pinshang Xu[a], Shuqing Li[a,b*], Mark Pawlett[c], Jianwen Zou[a,b]

[a]*Jiangsu Key Laboratory of Low Carbon Agriculture and GHGs Mitigation, College of Resources and Environmental Sciences, Nanjing Agricultural University, Nanjing 210095, China*

[b]*Jiangsu Key Lab and Engineering Center for Solid Organic Waste Utilization, Jiangsu Collaborative Innovation Center for Solid Organic Waste Resource Utilization, Nanjing Agricultural University, Nanjing 210095, China*

[c] *School of Energy, Environment and Agrifood, Cranfield University, MK43 0AL, UK*

[*]**Corresponding author:**

Weigang 1, Nanjing Agricultural University, Nanjing 210095, Jiangsu, China

E-mail: jywang@njau.edu.cn (J. Wang), shuqingli@njau.edu.cn (S. Li); Tel.: 86-025-84395210; Fax: 86-025-84395210

Abstract

In order to reduce soil and environmental quality degradation associated with the use of synthetic nitrogen (N), substituting chemical fertilizer with organic or bio-organic fertilizer has become an increasingly popular option. However, components of this fertilizer strategy related to mitigation of soil N-oxide emissions and maintenance of crop yield remain uncertain. Here, we evaluated the effects of three different fertilizer strategies, with equal amounts of N, on nitrous oxide (N_2O) and nitric oxide (NO) emissions, vegetable yield, and yield-scaled N_2O and NO emissions under three consecutive cucumber growing seasons. The three treatments were chemical fertilizer (NPK, urea), organic fertilizer (O, composted cattle manure), and bio-organic fertilizer (O+T, O combined with *Trichoderma*.spp). Results showed that the NPK plot had the highest area-scaled emissions of N_2O (13.1±0.48 kg N ha^{-1} yr^{-1}) and NO (5.01±0.34 kg N ha^{-1} yr^{-1}), which were 1.3–1.4 and 3.1–3.7 times greater than the O and O+T plots, respectively. The annual direct emission factors for N_2O and NO were 2.08% and 0.92% for the NPK plot, which declined to 1.34% and 0.09% in the O plot, and 1.12% and 0.03% in the O+T plot, respectively. The annual vegetable yield was 117±2.9 t ha^{-1} for NPK plot and 122±2.0 t ha^{-1} for O + T plot, which was higher than 111±1.7 t ha^{-1} for O plot. The yield-scaled N_2O+NO emissions differed significantly with fertilization treatment, with the lowest value observed in the O+T plot. We attributed the lower soil N-oxide emissions following organic fertilizer application to the slow release of available N and enhanced denitrification caused by the increase of soil dissolved organic carbon and pH. Compared with the use of organic fertilizer alone, the addition of *Trichoderma*.spp significantly increased the potential denitrification rate but decreased N_2O emissions, which

Published by Elsevier. This is the Author Accepted Manuscript issued with: Creative Commons Attribution Non-Commercial No Derivatives License (CC:BY:NC:ND 4.0). The final published version (version of record) is available online at DOI:10.1016/j.geoderma.2020.114730. Please refer to any applicable publisher terms of use.

may have promoted the reduction of N_2O to N_2. Therefore, our results suggest that adopting composted organic fertilizer mixtures with microbial inoculants could be a win-win practice to mitigate gaseous N losses and simultaneously improve crop yield in intensively managed vegetable cropping systems.

Keywords: Nitrous oxide; Nitric oxide; Static chamber technique; *Trichoderma*; Emission factor; Drip irrigation

1. Introduction

Nitrous oxide (N_2O) and nitric oxide (NO) are two important atmosphere trace gases, directly or indirectly contributing to global climate and environmental changes (IPCC, 2013). N_2O is a potent greenhouse gas and the dominant stratospheric ozone-depleting substance (Ravishankara et al., 2009). NO is a key precursor of tropospheric ozone (O_3) and contributes to the formation of acid rain (Pilegaard, 2013). Agricultural soils are significant sources of N_2O and NO, primarily due to the increased use of chemical or organic nitrogen (N) fertilizers, emitting about 80% and 10% of the total anthropogenic emissions of N_2O and NO, respectively (Davidson, 2009; IPCC, 2013; UNEP, 2019). Emissions of N_2O and NO are predicted to increase in the future, as the global use of N fertilizers is forecast to increase threefold by 2050 to meet the doubling of global food demand (Alexandratos and Bruinsma, 2012; Mueller et al., 2012).

Vegetable cultivation under greenhouse conditions can lead to substantial emissions of soil N-oxides. Nitrogen fertilizer application rates in greenhouse vegetable fields are often several times higher than in other cereal grain cropping systems (Liu et al., 2013; Rashti et al., 2015). Because of frequent irrigation and high temperatures, greenhouse vegetable cropping systems are highly susceptible to N losses, with annual soil N_2O and NO emissions are as high as 60.5 and 10.8 kg N ha^{-1}, respectively (Yao et al., 2019a; Zhang et al., 2016). In agricultural soils, N_2O and NO emissions are mainly produced as by-products through the biotic processes of nitrification, denitrification, and nitrifier denitrification (Firestone and Davidson, 1989; Pilegaard, 2013; Wrage-Mönnig et al., 2018), while their emissions are influenced by the same factors in different ways (Loick et al., 2017). There has been a much

75 stronger focus on N_2O than on NO or combined N_2O+NO emissions from vegetable fields
76 (e.g., De Rosa et al., 2018; Zhang et al., 2018). At the regional or country scale, N_2O or NO
77 estimates generally described using emission factors (EF) to quantify the amount of N_2O-N or
78 NO-N lost as a function of the N inputs, excluding background N_2O or NO emissions
79 (Hergoualc'h et al., 2019). However, studies have found that the N_2O emission factors for
80 upland crops may differ between the crop-growing season (EF_{gs}) and the whole year (EF_{wy}),
81 especially for vegetables (Shang et al., 2020). Different water management practices in
82 agricultural systems can also result in high variability in EFs (Cayuela et al., 2017). Besides,
83 5.5-20.6% of the annual NO emissions from greenhouse vegetables occurred in the non-
84 growing stage (Yao et al., 2019a), suggesting the importance of annual measurement of NO
85 emissions. Nevertheless, the EF_{gs} has been commonly used to determine national N_2O or NO
86 inventories for vegetable fields (Rashti et al., 2015; Wang et al., 2011). Exploring the EFs of
87 N_2O and NO over an annual period would contribute to a more accurate estimate of N_2O and
88 NO emissions from intensively managed greenhouse vegetable cropping systems.

89 In the past decade, shifts in fertilization strategies have received increasing attention due
90 to the potential for greenhouse gas mitigation. For example, to address the challenges
91 associated with the use of chemical N fertilizer (e.g., soil quality degradation, soil
92 acidification, and groundwater pollution), substitution of synthetic fertilizer with organic
93 fertilizer is promoted in intensively managed cropping systems (Zhang et al., 2020; Sanz-
94 Cobena et al., 2017). However, the side effects of this fertilization strategy on N_2O and NO
95 emissions are uncertain. Organic fertilizers play multiple roles in microbial-mediated N_2O
96 production, leading to stimulatory or inhibitory effects. For instance, the easily mineralizable

97 carbon (C) supply from poultry manure can stimulate heterotrophic denitrification, resulting
98 in larger N_2O emissions than occur with chemical N fertilization (Hayakawa et al., 2009). In
99 contrast, a higher denitrification rate as a result of organic fertilizer application may promote
100 the production of N_2, with no significant difference in N_2O emissions between organically
101 farmed soils and conventionally farmed soils (Kramer et al., 2006). Several studies have
102 reported that substituting organic for chemical fertilizer can contribute to decreased NO
103 emissions (Akiyama and Tsuruta, 2003a, b; Meijide et al., 2007; Vallejo et al., 2006).
104 Additionally, compared with liquid organic fertilizers, solid and composted organic fertilizers
105 have a low N_2O and NO emission potential due to reduced mineral N concentrations,
106 especially the NH_4^+ content (Aguilera et al., 2013; Bertora et al., 2008; Meijide et al., 2007).
107 The combination of drip irrigation with organic fertilizer reduced N_2O emissions by 28% but
108 had no significant effect on NO emissions when compared with furrow irrigation (Sanchez-
109 Martin et al., 2010). These scenarios could be highly dependent on climate, soil properties,
110 site-specific management practices, as well as the biochemical quality of the organic
111 materials. Nevertheless, emerging evidence suggests that the differences in the rate of N
112 release from synthetic and organic fertilizers can have the potential to significantly influence
113 soil N-oxide emissions (Prosser et al., 2020). This is mainly due to niche preference of
114 ammonia-oxidizing bacteria (AOB) or archaea (AOA) (Hink et al., 2017; 2018; Stein, 2019).

115 Substituting chemical fertilizer with organic fertilizer may have a neutral or even
116 negative effect on vegetable yield, especially when the replacement rate of synthetic N with
117 organic N exceeds 75% (Xia et al., 2017). The release of available N from composted organic
118 fertilizer is slow and may not be able to meet the high N demand of fast-growing and high-

yield vegetables (Berry et al., 2002). Thus, a promising alternative strategy is the use of bio-organic fertilizers (a mixture of organic material with beneficial soil microorganisms). *Trichoderma.* spp, a biological control and plant growth-promoting agent, is commonly used in bio-organic fertilizers (Harman et al., 2004). Colonization of *Trichoderma* on the root surface was shown to promote the development of plant roots and improve N use efficiency (Shoresh et al., 2010). The use of *Trichoderma* enriched bio-organic fertilizer has been shown to enhance plant uptake of soil nutrients and stimulate plant growth when compared with organic or chemical fertilizer (Pang et al., 2017). However, the effect of bio-organic fertilizer on soil N-oxide emissions remains unclear. Meanwhile, the benefits or trade-offs from substituting synthetic fertilizer with organic or bio-organic fertilizers are not well-known in vegetable fields. Therefore, it is necessary to evaluate the yield-scaled N_2O+NO emission (expressed as N_2O+NO produced per unit of crop yield) for balancing soil N losses and food security in agricultural ecosystems (Linquist et al., 2012).

In this study, we conducted an in situ field measurement over a 16-month period to quantify the annual N_2O and NO fluxes under various fertilizer-N regimes and to quantify crop yield benefits on a cucumber monoculture under a drip irrigation system in the greenhouse. We hypothesized that i) the substitution of synthetic N by organic or bio-organic fertilizer would mitigate N-oxide emissions due to the slow N release from added organic fertilizers; and ii) that bio-organic fertilizer would decrease N_2O and NO emissions and increase crop yields, thus resulting in lower yield-scaled N_2O and NO emissions. Specifically, the objectives of our study were to i) quantify the seasonal and annual emission intensity of N_2O and NO and their direct EFs under different fertilizer-N regimes; ii) evaluate the effects

of synthetic and organic N management on yield-scaled emissions of N_2O and NO; and iii) clarify the environmental driving factors that regulate N-oxides emissions.

2. Materials and Methods

2.1 *Site description and experimental design*

Field experiments were conducted from August 2015 to December 2016 at a research site (31°95'N, 118°83'E) of Nanjing Agricultural University in suburban Nanjing, Jiangsu province, China. The experimental site has been in continuous vegetable cultivation (e.g., Chinese cabbage, cabbage and cucumber) for >5 years. The climate is characterized by a subtropical monsoon, with hot-rainy summers and mild-less rainy winters. From 2015 to 2016, the annual average temperature is 16.5°C, and summer precipitation accounts for about 52% of the total average annual precipitation of 1786 mm (Fig. S1). From July 1, 2016 to August 12, 2016 (the second fallow stage), the plastic greenhouse was refurbished and open to the air at this time. Rainfall during this period reached 537 mm, accounting for 30% of total annual rainfall. The soil was classified as a Eutric Planosol (FAO, 1981) with 31.1% sand, 17.7% silt, 50.2% clay, and a soil bulk density of 1.33 g cm^{-3}. The surface soil (0-15 cm) contained an organic C content of 13.0 g C kg^{-1}, a total N content of 1.74 g N kg^{-1}, and a pH (1:2.5, soil/water) of 5.63.

Four treatments with different fertilizer-N regimes were arranged in a randomized block design with three replicates (each plot with a size of 2.7 m × 2 m) within a plastic greenhouse (30 m × 6 m). A 0.3 m wide buffer row separated the adjacent plots. The four treatments referred to the unfertilized control (control), urea (NPK), organic fertilizer (O), and bio-

organic fertilizer (O+T). The composted organic fertilizer was derived from a mixture of cattle manure and mushroom residue with a mass ratio of 3:1, containing organic matter 401 g kg^{-1}, total N 16.4 g N kg^{-1}, phosphorus 12.3 g P_2O_5 kg^{-1}, and potassium 12.9 g K_2O kg^{-1}. The bio-organic fertilizer was organic fertilizer plus *Trichoderma guizhouense* NJAU 4742 (10^8 CFU g^{-1} dry weight (DW), an antagonist of *Fusarium oxysporum*) containing organic matter 445 g kg^{-1}, total N 14.5 g N kg^{-1}, phosphorus 10.5 g P_2O_5 kg^{-1} and potassium 14.5 g K_2O kg^{-1}. From August 2015 to December 2016, cucumber seedlings were transplanted at the two-leaf stage on August 19, 2015, March 26, and August 12, 2016, respectively. The final harvest dates were on October 17, 2015, June 13 and October 14, 2016, respectively. Each cucumber cropping cycle was divided into the cucumber growing stage and subsequent fallow stage (Table S1). In each cucumber growing stage, each treatment received equal amounts of 180 kg N ha^{-1}, 135 kg P_2O_5 ha^{-1}, and 180 kg K_2O ha^{-1}. The missing P and K in each treatment were supplemented with superphosphate and potassium sulfate, respectively. According to the local practice, basal fertilizers and topdressing were broadcasted on the soil surface, plowed into the soil at a depth of 3-4 cm, and then followed by irrigation. The drip irrigation system was applied in this study. The cucumber planting density is 5 plants m^{-2}. The details of cultivation and fertilization management for each treatment are shown in Table S1.

2.2 *Measurements of N_2O and NO fluxes*

Flux measurements were taken over the periods of August 17, 2015 to December 19, 2016. Fluxes of N_2O and NO were measured using a static opaque chamber technique comprising PVC chambers (50 cm × 50 cm × 50 cm) (Zhang et al., 2016; Zou et al., 2005). Chamber bases (area 0.25 m^2, height 15 cm, ~10 cm inserted into the soil) were installed on each plot

184 before transplanting. Within each chamber base, one cucumber was grown throughout each
185 growing season. Each cucumber plant in the chamber base was equipped with a rope, which
186 was hung on a bamboo frame. During sampling, the rope was taken off and the cucumber
187 plant was carefully hovered in the chamber. After the gas collection, the cucumber plants
188 were hung on the bamboo frame again. Gas sampling generally occurred once or twice a week
189 except following fertilizer application when it occurred every day or every two days during
190 the cucumber growing stages, and every 7 or 10 days during the fallow periods. Gas sampling
191 events occurred from 08:00 through 10:00 local standard time on each sampling day. On the
192 sampling day, five samples were collected from the headspace with a 1.5-L gas sampling bag
193 at 0, 5, 10, 15, and 20 min after chamber closure.

194 The N_2O concentrations in the samples were analyzed with a gas chromatograph
195 (Agilent 7890A, Santa Clara, USA) equipped with an electron capture detector (ECD). The
196 carrier gas was a gas mixture of argon-methane (5%:95%) at a flow rate of 40 mL min^{-1}. The
197 temperatures of the ECD detector and column were 300°C and 40°C, respectively. The N_2O
198 calibration standard gas concentration is 0.32×10^{-6} mol mol^{-1}. The gas sample analysis for NO
199 concentrations was performed using a model 42i chemiluminescence NO-NO_2-NOx analyzer
200 (Thermo Environmental Instruments Inc., Franklin, MA, USA), which was calibrated using
201 the TE-146*i* dilution-titration instrument (Zhang et al., 2016). The calibration standard gases
202 for N_2O and NO were provided by the National Center of Standard Matters (Beijing, China).
203 The fluxes (F, μg N m^{-2} h^{-1}) of N_2O and NO were calculated using the following equation:

204 $$F = \frac{V}{A} \times \frac{M \times P}{R \times (237 + T)} \times \frac{dC}{dt}$$

205 where, V is the volume of gas sampling chamber (m^3), A is the cover area of gas sample

chamber (m^2), M is the molar mass of N_2O (44 g mol^{-1}) or NO (28 g mol^{-1}), P is the air pressure inside the gas sample chamber (1.013 × 105 Pa), R is the universal gas constant (8.314 Pa m^3 mol^{-1} K^{-1}), T is the temperature inside the chamber(°C), dC/dt is calculated by the linear increases of N_2O or NO concentration with time. Note that the volume of the sampling chamber was not corrected for the volume occupied by the cucumber plant.

The average flux of N_2O or NO over each cucumber cropping cycle (growing stage + fallow stage) was the average of all measured fluxes weighted by the intervals between two adjacent measurements. The total seasonal N_2O or NO emissions were approximated by applying the trapezoid rule on time intervals between measured fluxes, assuming constant fluxes per day.

2.3 Measurements of soil physicochemical and biological properties

Parallel to gas sampling, alcohol thermometers were used to record the air temperature and soil temperature (5 cm depth) in the greenhouse. Soil volumetric water content (0-5 cm) was measured using an MP406 Moisture probe with an MPM 160 Moisture Probe Meter (ICT International Pty Ltd, Armidale, Australia). The ratio of measured soil volumetric water content to total porosity was expressed as soil water-filled pore space (WFPS). The total soil porosity was calculated as [1-(soil bulk density (g cm^{-3})/2.65)], assuming a soil particle density of 2.65 (g cm^{-3}). Soil samples were collected from surface layers (0-15 cm) at intervals of 10-15 days during the cucumber growing stages and at intervals of 20-30 days during the fallow stages. Soil samples collected from the three points on the diagonal of each plot were mixed and passed through a 2 mm sieve to remove stones and impurities. The

227 homogenized soil samples were stored at 4℃ for analysis of soil physicochemical properties
228 (Lu, 2000). Soil pH and electrical conductivity (EC) were measured using a pH probe (PHS-
229 3C, Shanghai, China) and a conductivity meter (FE-30, Shanghai, China) at the soil to water
230 ratio of 1:2.5 (*w*/*v*), respectively. Soil mineral N (NH_4^+ and NO_3^-) was determined by
231 extracting soil with 2 M KCl (soil: water ratio of 1:10) after shaking for 1 h on a rotary
232 shaker. Soil NH_4^+ and NO_3^- concentrations were measured using the indophenol blue method
233 and two-wavelength approach using a UV-VIS spectrophotometer, respectively (U-2900,
234 Hitachi, Tokyo, Japan). Soil dissolved organic carbon (DOC) was determined using a UV-
235 Persulfate TOC analyzer (Teledyne-Tekmar Phoenix 8000, Mason, Ohio, USA).

236 The soil potential denitrification rates were performed on soil samples on the final
237 harvest date of each cucumber season (i.e., October 17, 2015, June 13, 2016, and October 14,
238 2016) using the acetylene inhibition method (Tiedje, 1982). In brief, potassium nitrate (KNO_3
239 1 mM), chloramphenicol (1 g L^{-1}), and dextrose (1 mM) were added to 5 g of fresh soil in a
240 50-mL air-tight flask, kept anaerobic by repeated flushing with N_2. Acetylene (10%, *v*/*v*) was
241 injected to inhibit the reduction of N_2O to N_2, and samples were incubated for 3 h at 25°C. A
242 5 mL headspace sample was collected using a gas-tight syringe for the determination of N_2O
243 concentration.

244 ***2.4 Vegetable yield and area/yield-scaled N-oxide emissions***

245 Fresh cucumber yields (g N t^{-1} yield) were weighed in each plot. Due to the staggered
246 maturation of cucumber, the yields of multiple harvests were added to calculate the total
247 yields. The annual cucumber yield across the three cucumber cropping cycles was used to

calculate the annual N-oxide emission below.

The annual N-oxide emission (kg N ha^{-1}) was calculated as total emissions of N-oxide over the experimental period (16 months) multiplied by 12/16. The yield-scaled N-oxide emission (g N t^{-1} yield) was calculated as follows (Linquist et al., 2012):

Yield-scaled N_2O or NO emissions=Area-scaled N_2O or NO emissions/Fresh cucumber yield (1)

The N fertilizer-induced direct EF (%) of N_2O and NO were calculated as:

$$EF_{wy} = (ER_N - ER_0)/N\ applied \times 100 \quad (2)$$

$$EF_{gs} = (ER_N - ER_0)/N\ applied \times 100 \quad (3)$$

$$\Delta EF = EF_{wy} - EF_{gs} \quad (4)$$

where, *wy* and *gs* represent the whole year and cucumber growing season periods, respectively. ER_N (kg N ha^{-1}) is the total N_2O or NO emissions from the fertilizer N-applied (NPK, O, or O+T) plots; ER_0 (kg N ha^{-1}) is the total N_2O or NO emissions from the control.

2.5 ***Statistical analyses***

Data were tested for normality (using Shapiro-Wilk's test) and equality of variance (using Levene's test), and parameters with non-normal distributions or unequal variances were either logarithmical or square-root transformed prior to analysis. Differences in seasonal or annual N-oxide emissions, EFs, and vegetable yield between treatments were determined by the LSD test ($P<0.05$). The effects of fertilization, season, and their interactions on the fluxes of N_2O, NO and N_2O+NO, and soil properties were examined using a repeated-measures analysis of

variance (MANOVA) with the R package MANOVA.RM (Friedrich et al., 2019). A linear model was fitted with the logarithmically transformed values of N_2O+NO emission and soil parameters. To assess the relative importance of predictors in the linear model, we used the R package *relaimpo* to calculate the importance of each soil parameter (Gromping, 2006). The Kruskal–Wallis one-way analysis of variance (ANOVA) was used to investigate differences in the potential denitrification rate between treatments of each sampling date and in the N_2O/NO ratio (log-transformed) during the study period. A nonparametric multiple comparison test (Dunn's test) was performed to compare the difference in the sum of ranks between groups with the expected average difference. All statistical analyses were conducted using R 3.6.3 (R Core Team, 2020).

3. Results

3.1 Environmental variables

Soil temperature during the three cropping cycles was comparable, ranging from 10.7 to 33.5°C (mean: 22.4°C), with the highest value occurring in July 2016 (Fig. 1a). Soil WFPS ranged from 20.5 to 80.6% (mean: 40.6%) in the second cropping cycle and was higher than the average of first and third cropping cycles, ranging from 23.6 to 57.3% (mean: 28.4%). The main reason for this difference was that the field was subjected to heavy rainfall during the second fallow stage due to the refurbishment of the greenhouse. Since there was no significant difference in soil temperature and WFPS among all treatments, the average values are shown in Fig. 1a.

Nitrogen fertilization significantly increased soil NH_4^+-N and NO_3^--N contents, with the

289 highest soil NH_4^+-N and NO_3^--N contents in the NPK plot (both P<0.001; Fig. 1b and c, Table
290 S2). Urea had a much stronger effect on soil mineral N content than organic or bio-organic
291 fertilizers. During the whole experimental period, soil NH_4^+ contents averaged 10.0±0.1,
292 51.0±1.3, 15.6±0.7, and 15.5±1.1 mg N kg^{-1} for the control, NPK, O and O+T plots,
293 respectively. Soil NO_3^- contents ranged from 24.3 to 190.5 mg N kg^{-1} over the entire
294 observation period, averaging 52.5±2.1, 131.7±3.5, 87.7±4.4 and 80.8±1.5 mg N kg^{-1} for the
295 control, NPK, O and O+T plots, respectively.

296 Organic fertilizer application resulted in an increase in soil pH and DOC content
297 (P<0.001; Fig. 1d and f, Table S2). The average pH of the control was 5.5±0.0, which
298 increased significantly to 6.0±0.1 and 6.1±0.1 in the O and O+T plots, respectively, but
299 decreased to 5.1±0.0 in the NPK plot. The mean DOC content in the control plot was
300 122.2±0.3 mg C kg^{-1}, which increased by 19.4% and 16.0% in the O and O+T plots,
301 respectively, but declined by 11.5% in the NPK plot. The EC value in the NPK plot was
302 greater than the other three treatments, where no statistical differences occurred (Fig. 1e,
303 Table S2).

304 ***3.2 N_2O fluxes***

305 The seasonal patterns of N_2O fluxes were similar among the fertilization treatments, with
306 frequent N_2O peaks generally driven by fertilizer application and/or irrigation (and/or rainfall)
307 events (Fig. 2a). Throughout the entire experimental period, N_2O fluxes were lower in the
308 control plot than in fertilized treatments, although several relatively small peaks were
309 observed following irrigation or rainfall events. Seasonal N_2O fluxes averaged 61.2±3.5,

310 173.8±8.1, 120.9±6.9 and 132.8±11.2μg N m^{-2} h^{-1} for the control, NPK, O and O+T plots,
311 respectively (Table S3). Generally, substantial N_2O emissions occurred during the cucumber-
312 growing stages, while N_2O fluxes were relatively low during the fallow stages, except for the
313 second fallow stage. Across all treatments, total N_2O emissions were 0.33–0.56 and 0.23–0.41
314 kg N ha^{-1} during the first and third fallow stages, accounting for only 15.6–23.0% and 11.5–
315 18.9% of the first and third cucumber cropping cycle, respectively. While for the second
316 fallow stage, total N_2O emissions across all treatments were 2.07–5.91 kg N ha^{-1}, accounting
317 for 55–61% of the second cucumber cropping cycle (Fig. 2a, Table S4). Overall, relative to
318 the NPK plot, seasonal N_2O emissions were significantly reduced by 13.2–34.3% and 17.4–
319 36.6% under O and O+T plots (P<0.01), respectively, but there was no significant difference
320 between the O and O+T plots (Table S5).

321 The N fertilizer-induced EF of N_2O over the whole year period (EF_{wy}) ranged from
322 1.12% to 2.08% (Table 1), with an EF_{gs} of 0.61–0.89%, 0.82–1.81% and 0.52–1.19% for the
323 corresponding cropping cycle (Table S4). Both EF_{wy} and EF_{gs} of N_2O for the three fertilizer
324 plots followed the order NPK > O ≈ O+T, such that EF_{wy} under O and O+T plots were
325 significantly reduced by 35.6% and 46.0% relative to the NPK plot, respectively.

326 *3.3* ***NO fluxes***

327 The seasonal pattern of the soil NO flux was distinct from that of the soil N_2O flux and
328 differed significantly among N fertilized plots (Fig. 1b). During the entire experimental
329 period, NO fluxes under the NPK plot ranged from 11.7 to 216.0 μg N m^{-2} h^{-1}, with
330 pronounced peaks captured following urea application events. In contrast, the NO fluxes in

331 the O and O+T plots remained unaffected. Seasonal mean NO fluxes were 16.0±1.7,
332 64.2±4.4, 21.3±1.6, and 17.4±1.6 μg N m^{-2} h^{-1} for the control, NPK, O and O+T plots,
333 respectively (Table S3).

334 Across all treatments, annual NO emissions ranged from 1.28 to 5.01 kg N ha^{-1}, with the
335 corresponding range of 0.51–1.76, 0.68–3.00, and 0.48–1.92 kg N ha^{-1} for the three cucumber
336 cropping cycles (Table1, Table S4). About 56.7% of NO emissions occurred during the fallow
337 periods. Across all treatments, the mean total NO emission during the second fallow period
338 was 0.62±0.03 kg N ha^{-1}, which was slightly higher than that of 0.55±0.04 kg N ha^{-1} (P >0.05)
339 in the first fallow period and significantly higher than that of 0.43±0.03 kg N ha^{-1} (P <0.05) in
340 the third fallow period. Relative to the NPK plot, seasonal NO emissions were significantly
341 reduced by 60.3-70.8% and 69.3-76.2% in the O and O+T plots, respectively, while there was
342 no clear difference between the O and O+T plots. The N fertilizer-induced EF for NO (EF_{gs})
343 was -0.02–0.28%, -0.01–0.84%, and 0.03–0.50% for the first, second, and third cropping
344 cycle, respectively (Table S4). The EF_{wy}s of NO in the O and O+T plots were significantly
345 decreased by 90.3% and 97.4% relative to the NPK plot, respectively (Table 1).

346 *3.4* ***Combined N_2O+NO emissions and yields***

347 Annual N_2O+NO emissions totaled 5.98, 18.2, 11.8, and 10.6 kg N ha^{-1} for the control, NPK,
348 O, and O+T plots, respectively (Table 1). The application of N fertilizers enhanced vegetable
349 yields by 6.5–35.6%. Vegetable yields decreased by 2.9–5.8% in the O plot, but increased by
350 2.8-9.0% in the O+T plot, when compared to the NPK plot. The annual area- and yield-scaled
351 N_2O+NO emissions were significantly higher in the NPK plot than in the O and O+T plots

(Table 1). Compared with the O plot, annual yield-scaled N_2O+NO emissions were significantly reduced by 17.7% in the O+T plot. The N fertilizer-induced emission factor of combined N_2O+NO (EF_{wy}) was 3.00%, 1.45%, and 1.15% for the NPK, O, and O+T plots over the annual scale, respectively (Table 1).

3.5 ***The potential denitrification rate and variable importance analysis***

Soil potential denitrification rates were measured at the end of each cucumber growing season and differed significantly between treatments. Nitrogen fertilization generally increased soil potential denitrification rate relative to the control (Fig. 3). A greater increase in the potential denitrification rate was observed in the O (110.7–222.4%) and O+T plots (219.7–287.5%) compared with the NPK plot (22.5–53.4%), and the O+T plot always had the highest potential denitrification rate.

Results of linear regression analysis showed that the selected variables explained 71.4% to 81.4% of the observed variability in the combined N_2O+NO emissions (Fig. 4). Across all treatments, soil temperature and WFPS were the first two important variables in explaining the variance of N_2O+NO emissions. Soil NH_4^+ content played a greater role in the NPK plot, whereas soil NO_3^- in the O and O+T plots was more important than the remaining variables.

4. Discussion

4.1 ***Responses of N-oxide emissions to synthetic and organic N fertilizer applications***

Our results demonstrated that applying organic fertilizers instead of synthetic N fertilizer can significantly decrease soil N_2O and NO emissions in greenhouse cucumber monocultures with

372 a drip irrigation system. For comparison, we compiled data from previous studies as shown in
373 Table 2 and found that the mitigation effects of organic substitution on soil N-oxide emissions
374 also occurred in cereal (Meijide et al., 2007; Yan et al., 2015; Yao et al., 2019b) and other
375 vegetable fields (Akiyama and Tsuruta, 2003a; Vallejo et al., 2006). Several reasons below
376 may contribute to explain lower N-oxide emissions following organic N fertilizer application.
377 First, relative to the organic N fertilization treatments, significantly higher mineral N
378 concentrations in the NPK plot may have contributed to greater N-oxide emissions, which is
379 the major factor driving microbial formation of N-oxide (Davidson et al., 2000). This is
380 consistent with the expectations of the conceptual hole-in-the-pipe model, which hypothesizes
381 that the total N-oxide gas flux is proportional to the rates of microbial N transformation
382 processes (Firestone and Davidson, 1989). The lower WFPS in this study (usually <60%)
383 suggested that the nitrifying communities may play an important role in soil N_2O and NO
384 production (Bateman and Baggs, 2005; Russow et al., 2008). Hence, the second possible
385 reason is that niche specialization of AOB and AOA associated with NH_4^+ supply may result
386 in differences in N_2O emission between synthetic and organic fertilization treatments (Hink et
387 al., 2017; 2018). As suggested by Hink et al. (2018), low NH_4^+ supply from the organic
388 fertilization treatments may favor AOA growth with low N_2O yield (N_2O-N produced per
389 NO_2^--N generated from ammonia oxidation), while high NH_4^+ supply after urea application
390 can lead to growth of AOA and AOB, thereby resulting in greater N_2O production because of
391 high N_2O yield by AOB. Furthermore, we probably attributed the reduced emission of N-
392 oxide following organic fertilizers application to the enhanced soil denitrification and the
393 promoted reduction of N_2O to N_2. Support for this assumption comes from recent evidence

394 suggesting that organic fertilizer application can promote the consumption of N_2O to N_2 in
395 global agricultural soils, despite increased soil denitrification following synthetic or organic
396 fertilizer application (Wang et al., 2018). The rationale for this is that organic C addition can
397 stimulate heterotrophic microbial activity and enhance soil O_2 consumption, which thereby
398 favors the development of anaerobic conditions for denitrification (Dambreville et al., 2006;
399 Kramer et al., 2006).

400 Nevertheless, organic fertilizer application has divergent impacts on soil N_2O emissions
401 (Table 2). For example, some studies have shown that organic fertilizer increased N_2O
402 emissions 6-fold compared to synthetic N fertilizer (Akiyama and Tsuruta, 2003b; Hayakawa
403 et al., 2009). Differential responses of N_2O emissions to organic fertilizer application are
404 associated with their decomposability, that is, organic matter with a relatively low C/N ratio
405 can be important source, but organic matter with a relatively high C/N ratio is less likely to
406 emit N_2O (Akiyama and Tsuruta 2003a). A recent meta-analysis indicated that the threshold
407 of C/N ratios for organic fertilizers was 8.6 (95% confidence interval: 4.5–22.3) in terms of
408 the positive or negative responses of N_2O emissions from acid soils (He et al., 2019). In our
409 study, organic fertilizers with a C/N ratio of ~15 contributed to lower N_2O emissions relative
410 to the chemical N application.

411 Regarding NO, it is an essential metabolite in AOA that is produced and immediately
412 consumed with tight control, whereas AOB tend to produce and release it (Kozlowski et al.,
413 2016; Stein, 2019). Support for this explanation comes from a global meta-analysis where it
414 was found that organic or mixed N fertilizer application is less effective at stimulating soil
415 NO emissions when compared to synthetic N fertilizer application (Liu et al., 2017). Unlike

N_2O, the consistent and negative response of soil NO emission to organic fertilizer applications as shown in Table 2 is likely due to i) as discussed above, the dominant AOA emit less NO when organic fertilizer is applied (Kozlowski et al., 2016; Stein, 2019); and ii) the addition of organic matter can promote some heterotrophic bacteria to oxidize NO through aerobic co-oxidation, rather than reducing NO to N_2O by denitrifying bacteria (Dunfield and Knowles, 1998).

4.2 Responses of yield-scaled N-oxide emissions to synthetic and organic N fertilizer applications

When linking N-oxide emissions to vegetable yields, our results show that organic and bio-organic fertilizer applications significantly decreased yield-scaled N_2O+NO emissions when compared to the synthetic N fertilizer. However, there was a yield penalty when organic fertilizer was used, but the use of bio-organic fertilizer slightly increased vegetable yields. In agreement with our findings, plant growth and crop yield were better in the *Trichoderma* enriched bio-organic treatment than in the organic fertilizer treatment (Pang et al., 2017). *Trichoderma*.spp were reported to be able to produce auxin-like phytohormones and several sparingly soluble nutrients, thereby promoting the development of the plant root system and uptake of soil nutrients (Cai et al., 2013). Enriched bio-organic fertilizer can also maintain a more diverse and stable soil microbiome for plant growth (Pang et al., 2017). Additionally, a meta-analysis of 143 studies showed that the full substitution of chemical N by manure in short-or medium-term (<10 years) will decrease crop yields, but the negative impact on yields will disappear over time (≥10 years), probably due to the various positive effects of manure in soil remediation (Zhang et al., 2020). Thus, our results demonstrate that the combination of

organic fertilizers with microbial inoculants present a potentially pragmatic option for maintaining crop yield and decreasing yield-scaled N-oxide emissions.

4.3 ***Emission factors of N_2O and NO***

The N fertilizer-induced EF of N_2O and NO may be underestimated without considering their emissions during the fallow period. Our results showed that EF_{gs} of N_2O and NO were, on average, 41% and 58% lower, respectively than those calculated for the whole cycle. The underestimation of NO by more than half indicates that the fallow period is an important period for NO emissions in this vegetable system. It has been well acknowledged that N-oxide emissions from agricultural land are spatially and temporally variable (Smith, 2017). In this study, the EFs of N_2O and NO varied seasonally because of changes in environmental conditions (Table S4). The greater N-oxide emissions were captured during the second cucumber cropping cycle when soil moisture ranged from 40 to 80% WFPS and soil temperature became gradually elevated. Thus, there is a need to measure soil N-oxide emissions over the whole year if we are to obtain the reliable EFs for their emission estimates using the Tier 2 method (Shang et al., 2020; Hergoualc'h et al., 2019).

Across all fertilized treatments, the mean value of EF_{wy} of N_2O is 1.51%, which is very close to the mean EF of N_2O (1.41%, 95% confidence interval: 1.19–1.64%) for vegetable cropping systems (Yang et al., 2019). The EF_{wy} of N_2O also falls within the IPCC reported default range of 0.1–1.8% (Hergoualc'h et al., 2019). However, our results are notably higher than the earlier estimates of 0.49–0.55% for Chinese vegetable fields (Wang et al., 2011), 0.51% for drip irrgation systems in Spain (Cayuela et al., 2017), and 0.94% for global

vegetable fields (Rashti et al., 2015). For NO, the EF_{wy} of 0.92% in the NPK plot was largely higher than 0.03% and 0.09% for O and O+T plots. These values were markedly lower than the reported mean value of 1.71% for global vegetable fields (Liu et al., 2017). Similarly, the EF_{wy} of N_2O+NO was estimated to be 1.15–3.00%, which was lower than the global average of 4.13% for vegetable fields (Liu et al., 2017). Collectively, these results suggest that there is great uncertainty in estimating N-oxide emissions from specific cropping system using default EFs.

4.4 Implications

This study reveals several key implications for soil N-oxide emissions within intensively managed greenhouse vegetable cropping systems. First, substantial increases in the magnitude of soil N-oxide emissions can be expected following N application from this cropping system under drip irrigation, independent of the forms of N fertilizer. In order to avoid significant N-oxide emissions, it is necessary to avoid maintaining plastic greenhouses in the hot-rainy summers in the subtropical regions. Second, the shift from synthetic N fertilizers to organic fertilizers can mitigate soil N-oxide emissions under greenhouse vegetable cultivation conditions. Our results demonstrate that organic fertilizers not only contribute greatly to reduced N-oxide emissions but also result in a very low emission intensity for NO. Third, from an agronomic perspective, bio-organic fertilizer is a promising option for fertilizer management in the current cropping system because of the significantly lower yield-scaled N-oxide emissions. We found that the fresh vegetable yield in bio-organic fertilizer plot was comparable to and slightly higher than that of the synthetic fertilization plot. Finally, our

findings highlight the importance of the year-round measurement of soil N-oxide emissions toward a reliable estimate of their emissions. Taken together, combining soil beneficial microbial inoculants with organic fertilizers is a promising strategy to reduce N losses and ensure yield in intensive greenhouse vegetable cropping systems. Further research is required to explore the effectiveness of microbial strains inoculate with organic fertilizer under a wider range of soil and environmental conditions.

Acknowledgments

This work was supported by the National Key Research and Development Projects of China (2017YFD0800201), the Fundamental Research Funds for the Central Universities (KYXK202002), and the National Natural Science Foundation of China (41877093).

Declaration of interests

 The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

491 **References**

492 Aguilera, E., Lassaletta, L., Sanz-Cobena, A., Garnier, J., Vallejo, A., 2013. The potential of
493 organic fertilizers and water management to reduce N_2O emissions in Mediterranean
494 climate cropping systems. A review. Agric. Ecosyst. Environ. 164, 32-52.
495 https://doi.org/10.1016/j.agee.2012.09.006.

496 Akiyama, H., Tsuruta, H., 2003a. Effect of organic matter application on N_2O, NO, and NO_2fluxes
497 from an Andisol field. Glob. Biogeochem. Cycle 17(4).
498 https://doi.org/10.1029/2002GB002016.

499 Akiyama, H., Tsuruta, H., 2003b. Nitrous oxide, nitric oxide, and nitrogen dioxide fluxes from
500 soils after manure and urea application. J. Environ. Qual. 32(2), 423-431.
501 https://doi.org/10.2134/jeq2003.4230.

502 Alexandratos, N., Bruinsma, J., 2012. World agriculture towards 2030/2050: the 2012 revision.

503 Bateman, E.J., Baggs, E.M., 2005. Contributions of nitrification and denitrification to N_2O
504 emissions from soils at different water-filled pore space. Biol. Fertil. Soils 41(6), 379-
505 388. https://doi.org/10.1007/s00374-005-0858-3.

506 Berry, P., Sylvester-Bradley, R., Philipps, L., Hatch, D.J., Cuttle, S.P., Rayns, F., Gosling, P.,
507 2002. Is the productivity of organic farms restricted by the supply of available nitrogen?
508 Soil Use Manage. 18, 248-255. https://doi.org/10.1111/j.1475-2743.2002.tb00266.x.

509 Bertora, C., Alluvione, F., Zavattaro, L., van Groenigen, J.W., Velthof, G., Grignani, C., 2008. Pig
510 slurry treatment modifies slurry composition, N_2O, and CO_2 emissions after soil

511 incorporation. Soil Biol. Biochem. 40(8), 1999-2006.
512 https://doi.org/10.1016/j.soilbio.2008.03.021.

513 Cai, F., Yu, G., Wang, P., Wei, Z., Fu, L., Shen, Q., Chen, W., 2013. Harzianolide, a novel plant
514 growth regulator and systemic resistance elicitor from *Trichoderma harzianum*. Plant
515 Physiol. Bioch. 73, 106-113. https://doi.org/10.1016/j.plaphy.2013.08.011.

516 Cayuela, M.L., Aguilera, E., Sanz-Cobena, A., Adams, D.C., Abalos, D., Barton, L., Ryals, R.,
517 Silver, W.L., Alfaro, M.A., Pappa, V.A., Smith, P., Garnier, J., Billen, G., Bouwman, L.,
518 Bondeau, A., Lassaletta, L., 2017. Direct nitrous oxide emissions in Mediterranean
519 climate cropping systems: Emission factors based on a meta-analysis of available
520 measurement data. Agric. Ecosyst. Environ. 238, 25-35.
521 https://doi.org/10.1016/j.agee.2016.10.006.

522 Dambreville, C., Hallet, S., Nguyen, C., Morvan, T., Germon, J. C., Philippot, L., 2006. Structure
523 and activity of the denitrifying community in a maize-cropped field fertilized with
524 composted pig manure or ammonium nitrate. Fems. Microbiol. Ecol. 56(1), 119-131.
525 https://doi.org/10.1111/j.1574-6941.2006.00064.x.

526 Davidson, E. A., Keller, M., Erickson, H.E., Verchot, L. V., Veldkamp, E., 2000. Testing a
527 conceptual model of soil emissions of nitrous and nitric oxides. BioScience 50, 667.
528 doi:10.1641/0006-3568(2000)050[0667:TACMOS]2.0.CO;2

529 Davidson, E.A., 2009. The contribution of manure and fertilizer nitrogen to atmospheric nitrous
530 oxide since 1860. Nat. Geosci. 2(9), 659-662. https://doi.org/10.1038/ngeo608.

531 De Rosa, D., Rowlings, D.W., Biala, J., Scheer, C., Basso, B., Grace, P.R., 2018. N_2O and CO_2

emissions following repeated application of organic and mineral N fertiliser from a vegetable crop rotation. Sci. Total Environ. 637, 813-824. https://doi.org/10.1016/j.scitotenv.2018.05.046.

Dunfield, P.F., Knowles, R., 1998. Organic matter, heterotrophic activity, and NO consumption in soils. Glob. Change Biol. 4(2), 199-207. https://doi.org/10.1046/j.1365-2486.1998.00123.x.

FAO, 1981. FAO-UNESCO Soil Map of the World 1:5000000 V Europe, UNESCO, Paris, France.

Firestone, M.K., Davidson, E.A., 1989. Microbiological basis of NO and N_2O production and consumption in soil. Exchange of trace gases between terrestrial ecosystems and the atmosphere 47, 7-21.

Friedrich, S., Konietschke, F., Pauly, M., 2019. Resampling-based analysis of multivariate data and repeated measures designs with the R package MANOVA. RM. R package version 0.3.4. https://CRAN.R-project.org/package=MANOVA.RM

Gromping, U., 2006. Relative importance for linear regression in R: The package relaimpo. J. Stat. Softw. 17(1), 1-27.

Guardia, G., Cangani, M.T., Sanz-Cobena, A., Lucas Junior, J., Vallejo, A., 2017. Management of pig manure to mitigate NO and yield-scaled N_2O emissions in an irrigated Mediterranean crop. Agric. Ecosyst. Environ. 238, 55-66. https://doi.org/10.1016/j.agee.2016.09.022.

Harman, G.E., Howell, C.R., Viterbo, A., Chet, I., Lorito, M., 2004. *Trichoderma* species-opportunistic, avirulent plant symbionts. Nat. Rev. Mic. 2(1), 43-56.

https://doi.org/10.1038/nrmicro797.

Hayakawa, A., Akiyama, H., Sudo, S., Yagi, K., 2009. N_2O and NO emissions from an Andisol field as influenced by pelleted poultry manure. Soil Biol. Biochem. 41(3), 521-529. https://doi.org/10.1016/j.soilbio.2008.12.011.

He, T., Yuan, J., Luo, J., Wang, W., Fan, J., Liu, D., Ding, W., 2019. Organic fertilizers have divergent effects on soil N_2O emissions. Biol. Fertil. Soils 55(7): 685-699. https://doi.org/10.1007/s00374-019-01385-4.

Hergoualc'h, K., Akiyama, H., Bernoux, M., Chirinda, N., Prado, A. del, Kasimir, Å., MacDonald, J.D., Ogle, S.M., Regina, K., van der Weerden, T.J., Liang, C., Noble, A., 2019. N_2O Emissions From Managed Soils, and CO_2 Emissions From Lime and Urea Application, in: Calvo Buendia, E., Tanabe, K., Kranjc, A., Baasansuren, J., Fukuda, M., Ngarize, S., Osako, A., Pyrozhenko, Y., Shermanau, P., Federici, S. (Eds.), 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. IPCC, Switzerland, pp. 1–48.

Hink, L., Nicol, G.W., Prosser, J.I., 2017. Archaea produce lower yields of N_2O than bacteria during aerobic ammonia oxidation in soil. Environmental Microbiology 19, 4829–4837. doi:10.1111/1462-2920.13282

Hink, L., Gubry-Rangin, C., Nicol, G.W., Prosser, J.I., 2018. The consequences of niche and physiological differentiation of archaeal and bacterial ammonia oxidisers for nitrous oxide emissions. ISME J. 12(4), 1084-1093. https://doi.org/10.1038/s41396-017-0025-5.

IPCC, 2013. Summary for policymakers. In: Climate Change 2013: The Physical science basis.

574 contribution of working group I to the fifth assessment report of the intergovernmental
575 panel on climate change. Cambridge University Press, Cambridge.
576 https://doi.org/10.1017/CBO9781107415324.

577 Kramer, S.B., Reganold, J.P., Glover, J.D., Bohannan, B.J., Mooney, H.A., 2006. Reduced nitrate
578 leaching and enhanced denitrifier activity and efficiency in organically fertilized soils. P.
579 Nati. Acad. Sci. USA. 103(12), 4522-4527. https://doi.org/10.1073/pnas.0600359103.

580 Kozlowski, J.A., Stieglmeier, M., Schleper, C., Klotz, M.G., Stein, L.Y., 2016. Pathways and key
581 intermediates required for obligate aerobic ammonia-dependent chemolithotrophy in
582 bacteria and *Thaumarchaeota*. ISME Journal 10, 1836–1845.
583 https://doi.org/10.1038/ismej.2016.2

584 Linquist, B., Groenigen, KJ., Adviento-Borbe, MA., Pittelkow, C., Kessel, C., 2012. An
585 agronomic assessment of greenhouse gas emissions from major cereal crops. Glob. Change
586 Biol. 18, 194-209. https://doi.org/10.1111/j.1365-2486.2011.02502.x.

587 https://doi.org/10.1016/j.geoderma.2017.06.007.

588 Liu, Q.H., Qin, Y.M., Zou, J.W., Guo, Y.Q., Gao, Z.L., 2013. Annual nitrous oxide emissions
589 from open-air and greenhouse vegetable cropping systems in China. Plant Soil 370(1-2),
590 223-233. https://doi.org/10.1007/s11104-013-1622-3.

591 Liu, S., Lin, F., Wu, S., Ji, C., Sun, Y., Jin, Y., Li, S., Li, Z., Zou, J., 2017. A meta-analysis of
592 fertilizer-induced soil NO and combined NO+N_2O emissions. Glob. Change Biol. 23(6):
593 2520-2532. https://doi.org/10.1111/gcb.13485.

594 Loick, N., Dixon, E., Abalos, D., Vallejo, A., Matthews, P., Mcgeough, K., Watson, C., Baggs,

595 E.M., Cardenas, L.M., 2017. “ Hot spots ” of N and C impact nitric oxide, nitrous oxide
596 and nitrogen gas emissions from a UK grassland soil. Geoderma 305, 336-345.

597 Lu, R., 2000. Methods of Soil and Agro-chemical Analysis, China Agric. Sci. Tech., Beijing,
598 China.

599 Meijide, A., Díez, J.A., Sánchez-Martín, L., López-Fernández, S., Vallejo, A., 2007. Nitrogen
600 oxide emissions from an irrigated maize crop amended with treated pig slurries and
601 composts in a Mediterranean climate. Agric. Ecosyst. Environ. 121(4), 383-394.
602 https://doi.org/10.1016/j.agee.2006.11.020.

603 Mueller, N.D., Gerber, J.S., Johnston, M., Ray, D.K., Ramankutty, N., Foley, J.A., 2012. Closing
604 yield gaps through nutrient and water management. Nature 490(7419), 254-257.
605 https://doi.org/10.1038/nature11420.

606 Pang, G., Cai, F., Li, R., Zhao, Z., Li, R., Gu, X., Shen, Q., Chen, W., 2017. *Trichoderma*-
607 enriched organic fertilizer can mitigate microbiome degeneration of monocropped soil to
608 maintain better plant growth. Plant Soil 416(1-2), 181-192.
609 https://doi.org/10.1007/s11104-017-3178-0.

610 Pilegaard, K., 2013. Processes regulating nitric oxide emissions from soils. Philos. T. R. Soc. B
611 368(1621) : 20130126. https://doi.org/10.1098/rstb.2013.0126.

612 Prosser, J.I., Hink, L., Gubry-Rangin, C., Nicol, G.W., 2020. Nitrous oxide production by
613 ammonia oxidizers: Physiological diversity, niche differentiation and potential mitigation
614 strategies. Glob. Chang Biol. 26(1), 103-118. https://doi.org/10.1111/gcb.14877.

615 R Core Team (2020). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Ravishankara, A.R., Daniel, J.S., Portmann, R.W., 2009. Nitrous Oxide (N_2O): The dominant ozone-depleting substance emitted in the 21st century. Science 326(5949), 123-125. https://doi.org/10.1126/science.1176985.

Rashti, M.R., Wang, W., Moody, P., Chen, C., Ghadiri, H., 2015. Fertiliser-induced nitrous oxide emissions from vegetable production in the world and the regulating factors: A review. Atmos. Environ.112, 225-233. https://doi.org/10.1016/j.atmosenv.2015.04.036

Russow, R., Spott, O., Stange, C.F., 2008. Evaluation of nitrate and ammonium as sources of NO and N_2O emissions from black earth soils (Haplic Chernozem) based on ^{15}N field experiments. Soil Biol. Biochem. 40(2), 380-391. https://doi.org/10.1016/j.soilbio.2007.08.020.

Sanchez-Martin, L., Meijide, A., Garcia-Torres, L., Vallejo, A., 2010. Combination of drip irrigation and organic fertilizer for mitigating emissions of nitrogen oxides in semiarid climate. Agr. Ecosyst. Environ. 137(1-2), 99-107. https://doi.org/10.1016/j.agee.2010.01.006.

Sanz-Cobena, A., Lassaletta, L., Aguilera, E., Prado, A. del, Garnier, J., Billen, G., Iglesias, A., Sánchez, B., Guardia, G., Abalos, D., Plaza-Bonilla, D., Puigdueta-Bartolomé, I., Moral, R., Galán, E., Arriaga, H., Merino, P., Infante-Amate, J., Meijide, A., Pardo, G., Álvaro-Fuentes, J., Gilsanz, C., Báez, D., Doltra, J., González-Ubierna, S., Cayuela, M.L., Menéndez, S., Díaz-Pinés, E., Le-Noë, J., Quemada, M., Estellés, F., Calvet, S., van Grinsven, H.J.M., Westhoek, H., Sanz, M.J., Gimeno, B.S., Vallejo, A., Smith, P., 2017.

637 Strategies for greenhouse gas emissions mitigation in Mediterranean agriculture: A
638 review. Agric. Ecosyst. Environ. 238, 5-24. https://doi.org/10.1016/j.agee.2016.09.038.

639 Shang, Z., Abdalla, M., Kuhnert, M., Albanito, F., Zhou, F., Xia, L., Smith, P., 2020.
640 Measurement of N_2O emissions over the whole year is necessary for estimating reliable
641 emission factors. Environ. Pollut. 259, 113864.
642 https://doi.org/10.1016/j.envpol.2019.113864.

643 Shoresh, M., Harman, G.E., Mastouri, F., 2010. Induced systemic resistance and plant responses
644 to fungal biocontrol agents. Annu. Rev. Phytopathol. 48, 21-43.
645 https://doi.org/10.1146/annurev-phyto-073009-114450.

646 Smith, K.A., 2017. Changing views of nitrous oxide emissions from agricultural soil: key
647 controlling processes and assessment at different spatial scales. Eur. J. Soil Sci. 68, 137–
648 155. doi:10.1111/ejss.12409. https://doi.org/10.1111/ejss.12409

649 Stein, L.Y., 2019. Insights into the physiology of ammonia-oxidizing microorganisms. Curr. Opin.
650 Chem. Biol. 49, 9-15. https://doi.org/10.1016/j.cbpa.2018.09.003.

651 Tiedje, J.M., 1982. Denitrificatin, In: A.L., Miller, R.H., Keeney, D.R. (Eds.), Methods of Soil
652 Analysis. Part 2. Chemical and Microbiological Properties. Madison, Wisconsin USA,
653 Madison, WI 53711, USA, pp. 1011–1026.
654 https://doi.org/10.2134/agronmonogr9.2.2ed.c47.

655 United Nations Environment Programme (UNEP), 2019. Fronties 2018/19 emerging issues of
656 environmental concern. Nairobi. http://hdl.handle.net/20.500.11822/27538.

657 Vallejo, A., Skiba, U., Garciatorres, L., Arce, A., Lopezfernandez, S., Sanchezmartin, L., 2006.

Nitrogen oxides emission from soils bearing a potato crop as influenced by fertilization with treated pig slurries and composts. Soil Biol. Biochem. 38(9), 2782-2793. https://doi.org/10.1016/j.soilbio.2006.04.040.

Wang, J., Chadwick, D.R., Cheng, Y., Yan, X., 2018. Global analysis of agricultural soil denitrification in response to fertilizer nitrogen. Sci. Total Environ.616-617, 908-917. https://doi.org/10.1016/j.scitotenv.2017.10.229.

Wang, J., Xiong, Z., Yan, X., 2011. Fertilizer-induced emission factors and background emissions of N_2O from vegetable fields in China. Atmos. Environ. 45(38), 6923-6929. https://doi.org/10.1016/j.atmosenv.2011.09.045.

Wrage-Mönnig, N., Horn, M.A., Well, R., Müller, C., Velthof, G., Oenema, O., 2018. The role of nitrifier denitrification in the production of nitrous oxide revisited. Soil Biol. Biochem. 123, A3-A16. https://doi.org/10.1016/j.soilbio.2018.03.020.

Xia, L., Lam, S.K., Yan, X., Chen, D., 2017. How does recycling of livestock manure in agroecosystems affect crop productivity, reactive nitrogen losses, and soil carbon balance? Environ. Sci. Technol. 51(13), 7450-7457. https://doi.org/10.1021/acs.est.6b06470.

Yan, G.X., Yao, Z.S., Zheng, X.H., Liu, C.Y., 2015. Characteristics of annual nitrous and nitric oxide emissions from major cereal crops in the North China Plain under alternative fertilizer management. Agr. Ecosyst. Environ. 207, 67-78. https://doi.org/10.1016/j.agee.2015.03.030.

Yang, T., Li, F., Zhou, X., Xu, C., Feng, J., Fang, F., 2019. Impact of nitrogen fertilizer,

679 greenhouse, and crop species on yield-scaled nitrous oxide emission from vegetable
680 crops: A meta-analysis. Ecol. Indic. 105, 717-726.
681 https://doi.org/10.1016/j.ecolind.2019.02.001.

682 Yao, Z., Yan, G., Wang, R., Zheng, X., Liu, C., Butterbach-Bahl, K., 2019a. Drip irrigation or
683 reduced N-fertilizer rate can mitigate the high annual N_2O+NO fluxes from Chinese
684 intensive greenhouse vegetable systems. Atmos. Environ. 212, 183-193.
685 https://doi.org/10.1016/j.atmosenv.2019.05.056.

686 Yao, Z., Zheng, X., Wang, R., Liu, C., Lin, S., Butterbach-Bahl, K., 2019b. Benefits of integrated
687 nutrient management on N_2O and NO mitigations in water-saving ground cover rice
688 production systems. Sci. Total Environ. 646, 1155-1163.
689 https://doi.org/10.1016/j.scitotenv.2018.07.393.

690 Zhang, J., Li, H., Wang, Y., Deng, J., Wang, L., 2018. Multiple-year nitrous oxide emissions from
691 a greenhouse vegetable field in China: Effects of nitrogen management. Sci. Total
692 Environ. 616-617, 1139-1148. https://doi.org/10.1016/j.scitotenv.2017.10.206.

693 Zhang, X., Fang, Q., Zhang, T., Ma, W., Velthof, G.L., Hou, Y., Oenema, O., Zhang, F., 2020.
694 Benefits and trade-offs of replacing synthetic fertilizers by animal manures in crop
695 production in China: A meta-analysis. Glob. Change Biol. 26(2), 888-900.
696 https://doi.org/10.1111/gcb.14826.

697 Zhang, Y., Lin, F., Jin, Y., Wang, X., Liu, S., Zou, J., 2016. Response of nitric and nitrous oxide
698 fluxes to N fertilizer application in greenhouse vegetable cropping systems in southeast
699 China. Sci. Rep. 6(1): 1-11. https://doi.org/10.1038/srep20700.

Zou, J.W., Huang, Y., Jiang, J.Y., Zheng, X.H., Sass, R.L., 2005. A 3-year field measurement of methane and nitrous oxide emissions from rice paddies in China: Effects of water regime, crop residue, and fertilizer application. Glob. Biogeochem. Cycle 19(2), GB2021. https://doi.org/10.1029/2004GB002401.

Table 1 Total annual N_2O, NO, N_2O+NO emissions (kg N ha^{-1}) and their direct emission factors (EF, %), cucumber yield (t ha^{-1}), and yield-scaled N_2O, NO and N_2O+NO emissions (g N t^{-1} yield).

	N_2O				NO				N_2O+NO			
	Control	NPK	O	O+T	Control	NPK	O	O+T	Control	NPK	O	O+T
Area-scaled	4.71±0.28c	13.1±0.48a	10.1±0.78b	9.26±0.59b	1.28±0.13b	5.01±0.34a	1.64±0.12b	1.37±0.12b	5.98±0.36 c	18.2±0.13a	11.8±0.79b	10.6±0.68b
Yield	100±1.33c	117±2.85ab	111±1.71b	122±1.98a								
Yield-scaled	47.0±2.30c	113±6.99a	91.0±5.68b	75.9±3.62b	12.8±1.35b	42.9±1.93a	14.8±1.18b	11.3±0.81b	59.8±3.14d	156±5.07a	106±5.61b	87.1±4.13c
EF_{wy}		2.08±0.12a	1.34±0.19b	1.12±0.12b		0.92±0.08 a	0.09±0.03b	0.03±0.03b		3.00±0.03 a	1.43±0.19b	1.15±0.17b
EF_{gs}		1.30±0.05a	0.76±0.08b	0.65±0.08b		0.54±0.06 a	0.03±0.01b	0.01±0.01b		1.84±0.02 a	0.79±0.07b	0.66±0.09b
ΔEF		0.79	0.58	0.47		0.38	0.06	0.02		1.17	0.64	0.49

Values are means ± SEM (n=3). Different letters within the row for each variable indicate significant differences between treatments at P<0.05. Control, without fertilization; NPK, chemical fertilizer-urea; O, organic fertilizer derived from composted cattle manure with mushroom residue; O+T, bio-organic fertilizer consisting of organic fertilizer plus *Trichoderma guizhouense* NJAU 4742.

709 **Table 2** An overview of studies simultaneously investigating the impacts of synthetic and organic fertilizer application on soil N_2O and NO emissions

Site/ climate	Crop/ soil	Day (d)	N fertilizer		N emissions (kg N ha^{-1})				% of change relative to synthetic N					
			Type	N input (kg N ha^{-1})	N_2O	NO	N_2O+NO	EF_c	N_2O	NO	N_2O+NO	Yield-scaled	EF_c	
Jiangsu, China/	Vegetable/	130	CMT	180	4.12	0.61	4.73	1.15	-30	-73	-41	-44	-62	This study
Subtropical-monsoon	Clay		CM	180	4.5	0.73	5.23	1.43	-23	-67	-35	-32	-52	
			U	180	5.84	2.23	8.07	3.00						
Hubei, China/	Rice	135	50%U+50%PM	150	0.87	0.07	0.94	0.48	-77	-42	-76	-76	-80	Yao et al.,
Subtropical-monsoon	Silt loam		U	150	3.74	0.12	3.86	2.42						2019 b
			50%U+50%PM	150	1.05	0.07	1.12	0.52	-77	-59	-76	-76	-81	
			U	150	4.47	0.17	4.64	2.87						
Madrid, Spain	Maize	365	67%U+33%CPS	180	0.66	3.5	4.16	0.90	-15	13	7	61	21	Guardia et
Temperate	Sandy-clay loam		67%U+33%LFPS	180	0.53	3.9	4.43	1.05	-32	26	14	55	41	al., 2017
			U	180	0.78	3.1	3.88	0.75						
Shandong, China	Wheat	240	70%(U+DP)+30%M	180	0.74	0.3	1.04	0.38	-16	-54	-32	-10	-42	Yan et al.,
Subtropical monsoon	Silt loam		70%CRU+30%M	180	0.89	0.35	1.24	0.49	1	-46	-19	-3	-25	2015
			U+DP	180	0.88	0.65	1.53	0.65						
Madrid, Spain	Maize	142	UPS	175	8.27	0.13	8.4	1.38	-4	-43	-5	-	-14	Meijide et
Subtropical	Sandy-loam		DPS	175	7.7	0.13	7.83	1.05	-10	-43	-11	-	-34	al., 2007
			71% PS+29%U	175	9.28	0.1	9.38	1.94	8	-57	7	-	20	
			71% SW+29%U	175	7.13	0.03	7.16	0.67	17	-87	-19	-	-58	
			U	175	8.57	0.23	8.8	1.61						

Madrid, Spain	Vegetable	150	UPS	175	5.62	0.1	5.72	1.15	-23	-58	-24	-	-48	Vallejo et
Temperate	Clay loam		DPS	175	4.69	0.1	4.79	0.63	-36	-58	-37	-	-71	al., 2006
			CPS	175	6.41	0.17	6.58	1.65	-12	-29	-13	-	-25	
			71%SW+ 29%U	175	5.65	0.07	5.72	1.16	-23	-71	-24	-	-48	
			U	175	7.31	0.24	7.55	2.21						
Tsukuba, Japan	Vegetable	42	PM	150	1.71	0.12	1.83	-	494	-92	3	-	-	Akiyama
Temperate	Sandy-		SM	150	0.47	0.16	0.63	-	63	-89	-64	-	-	and Tsuruta,
marine	loam		U	150	0.29	1.49	1.78	-						2003 b
Tsukuba, Japan	Vegetable	48	OC	150	0.35	0.23	0.58	-	223	-57	-8	-	-	Akiyama
Temperate	Sandy-		CM	150	0.03	0.02	0.05	-	-77	-95	-92	-	-	and Tsuruta,
marine	loam		U	150	0.11	0.52	0.63	-						2003 a
		47	FM	150	0.35	0.21	0.55	-	64	-80	-56	-	-	
			DC	150	0.06	0.08	0.14	-	-70	-93	-89	-	-	
			U	150	0.21	1.05	1.26	-						
Japan	Vegetable	365	PM	240	0.70	0.79	1.49	-	84	-50	-23	-	-	Hayakawa et
Temperate	Sandy-		PPM	240	2.72	0.69	3.41	-	616	-56	76	-	-	al., 2009
marine	loam		AS	240	0.38	1.56	1.94	-						

710 EF_c, emission factor of N_2O+NO; CMT, Cattle manure+*Trichoderma*. *spp*; CM, Cattle manure; U, Urea; PM, Poultry manure; CPS, Composting solid

711 fraction of pig slurry; LFPS, Liquid fraction of pig slurry; DP, Diammonium phosphate; M, Manure; CRU, Controlled-release urea; UPS, Untreated pig slurry;

712 DPS, Digested pig slurry; PS, Pig slurry; SW, Solid waste; SM, Swine manure; OC, Oilcake; FM, Fishmeal; DC, Dried cattle excreta; PPM, Pelleted poultry

713 manure; AS, Ammonium sulfate.

Figure captions

Fig. 1 Soil water-filled pore space (WFPS), soil and air temperature (**a**), NH_4^+ (**b**), NO_3^- (**c**), pH (**d**), electrical conductivity (EC; **e**), dissolved organic carbon (DOC; **f**) from a greenhouse mono-cucumber field under different fertilization treatments. The gray dashed arrows indicate fertilization. Values are means ± SEM (*n*=3). Control, without fertilization; NPK, chemical fertilizer-urea; O, organic fertilizer derived from composted cattle manure with mushroom residue; O+T, bio-organic fertilizer consisting of organic fertilizer plus *Trichoderma guizhouense* NJAU 4742.

Fig. 2 Year-round fluxes of soil N_2O (**a**) and NO (**b**) from a greenhouse mono-cucumber field under different fertilization treatments. The gray dashed and solid line arrows indicate fertilization and irrigation, respectively. Grey shading marks the fallow period between the cucumber growing seasons. The field during the second fallow period was open-air because of the refurbishment of the greenhouse. Values are means ± SEM (*n*=3). Control, without fertilization; NPK, chemical fertilizer-urea; O, organic fertilizer derived from composted cattle manure with mushroom residue; O+T, bio-organic fertilizer consisting of organic fertilizer plus *Trichoderma guizhouense* NJAU 4742.


Fig. 3 Potential denitrification rates in different sampling periods under different fertilization treatments. Different letters refer to significant differences between treatments at the $P<0.05$. Control, without fertilization; NPK, chemical fertilizer-urea; O, organic fertilizer derived


735 from composted cattle manure with mushroom residue; O+T, bio-organic fertilizer consisting
736 of organic fertilizer plus *Trichoderma guizhouense* NJAU 4742.


737


738 **Fig. 4** The relative importance of environmental factors in predicting N_2O+NO emissions
739 with a linear regression model under different fertilization treatments. Control, without
740 fertilization; NPK, chemical fertilizer-urea; O, organic fertilizer derived from composted
741 cattle manure with mushroom residue; O+T, bio-organic fertilizer consisting of organic
742 fertilizer plus *Trichoderma guizhouense* NJAU 4742.

743


(a)
WFPS
Soil temperature
Air temperature
WFPS (%)
Temperature (°C)
(b)
Control
O
NPK
O+T
NH4+ (mg N kg-1)
(c)
NO3- (mg N kg-1)
(d)
Soil pH
(e)
EC (ms cm-1)
(f)
DOC (mg C kg-1)
15/08/10
15/09/24
15/11/08
15/12/23
16/02/06
16/03/22
16/05/06
16/06/20
16/08/04
16/09/18
16/11/02
16/12/17
Date (yy/mm/dd)


(a)
Control
NPK
O
O+T
3200
2400
1000
500
0
N2O fluxes (μg N m−2 h−1)
(b)
250
200
150
100
50
0
NO fluxes (μg N m−2 h−1)
15/08/10
15/09/24
15/11/08
15/12/23
16/02/06
16/03/22
16/05/06
16/06/20
16/08/04
16/09/18
16/11/02
16/12/17
Date (yy/mm/dd)


Control
NPK
O
O+T
Potential denitrification rate
(ng N2O-N g-1 h-1)
100
75
50
25
0
a
a
a
b
b
b
c
c
c
d
1st
2nd
3rd
Season


Control
NPK
O
O+T
Soil T
WFPS
NO3−
NH4+
pH
DOC
EC
R2 = 0.71
R2 = 0.80
R2 = 0.80
R2 = 0.81
0.0 0.1 0.2 0.3 0.4 0.5 0.6
Relative importance



https://dspace.org/

Soil N-oxide emissions decrease from intensive greenhouse vegetable fields by substituting synthetic N fertilizer with organic and bio-organic fertilizers

Geng, Yajun; Wang, Jinyang; Sun, Zhirong; Ji, Cheng; Huang, Mengyuan ...

2020-09-24
Elsevier
https://doi.org/10.1016/j.geoderma.2020.114730
Geng Y, Wang J, Sun Z, et al., (2021) Soil N-oxide emissions decrease from intensive greenhouse vegetable fields by substituting synthetic N fertilizer with organic and bio-organic fertilizers. Geoderma, Volume 383, February 2021, Article number 114730

Downloaded from a DSpace repository. https://dspace.org/