

5 Title:

7 Comparing the antimicrobial effectiveness of NaCl
8 and KCl with a view to salt/sodium replacement

12 Authors: Eva Bidlas¹ and Ronald J.W. Lambert^{2*}

15 ¹Quality & Safety Department, Nestlé Research Centre, Vers-chez-les-Blanc, Lausanne,
16 Switzerland 1000.

17 ²Cranfield Health, Cranfield University, Cranfield, Bedfordshire, UK, MK43 0AL

21 Running title: Salt replacement

23 *Correspondence to Ronald Lambert PhD, rjwlambert@cranfield.ac.uk*

24 **Abstract**

25 A study using a small range of pathogenic bacterial species (*Aeromonas hydrophila*, *Enterobacter*
26 *sakazakii*, *Shigella flexneri*, *Yersinia enterocolitica* and 3 strains of *Staphylococcus aureus*) has
27 shown that potassium chloride has an equivalent antimicrobial effect on these organisms when
28 calculated on a molar basis. Combined NaCl and KCl experiments were carried out and data
29 analysed using a modification to the Lambert and Lambert (2003) model for combined inhibitors
30 (J. Appl. Microbiol. 95, 734–743) and showed that in combination KCl is a direct 1:1 molar
31 replacement for the antimicrobial effect of common salt. If this is a general finding then, where salt
32 is used to help preserve a product, partial or complete replacement by KCl is possible.

33

34

35

36 Keywords: Predictive modelling, hurdles, salt, potassium chloride, NaCl, KCl

1. Introduction

Salt (NaCl) is generally added to foodstuffs to 1. improve taste and 2. as a preserving agent. Indeed, historically, salt was among the very few effective preserving methods known. With the advent of refrigeration, better processing, packaging, transport and storage, there is less need for high salt levels to maintain product integrity. Furthermore, consumers want products with reduced sodium levels (e.g. due to its relationship with hypertension), but where salt has been added as a preservation hurdle, removal or reduction of the salt will reduce shelf-life and could affect safety in more microbiologically fragile products.

The most obvious replacement for salt (NaCl) in food products is potassium chloride (KCl). [Strong, Foster and Duncan \(1970\)](#) reported that for the growth of *Clostridium perfringens*, solute identity had a bearing on the amount of growth for a given a_w , with KCl having a demonstrably greater effect than NaCl. [Beuchat \(1974\)](#), however, reported that at equivalent a_w NaCl and KCl had equivalent effects against *Vibrio parahaemolyticus*; it was reported that in fermented meat products, the replacement of NaCl with KCl did not affect the degree of inhibition and or inactivation, but did alter the taste of the foodstuffs ([Gimeno, Astiasaran, and Bello 1999](#); [Gimeno, Astiasaran, and Bello 2001](#)). More recently, [Boziaris, Skandamis, Anastasiadi, and Nychas \(2007\)](#) have reported that equal-molar concentrations of NaCl or KCl exerted similar inhibitory effects against *Listeria monocytogenes* in terms of lag phase duration, growth or death rate and that NaCl can be replaced by KCl without risking the microbiological safety, with respect to *L. monocytogenes*, of the product. They also stated that in order to confirm this observation as general, a greater number of organisms needs to be studied.

62 In the work reported herein, we simply wanted to answer the following question: can KCl
63 be a direct or partial replacement for NaCl? Since the area of investigation is potentially
64 vast, we concentrated our initial efforts on a few species of pathogenic bacteria with which
65 we already had extensive modelling expertise on and which complimented other published
66 work.

67

68 **2. Materials and methods**

69

70 *2.1. Culture Preparation*

71 *Aeromonas hydrophila* (ATCC 7092), *Enterobacter sakazakii* (1387-2NL), *Staphylococcus*
72 *aureus* (ATCC 6538, ATCC 25923 (labeled as ST121 in this report), ST55 (isolated from
73 pasta)), *Yersinia enterocolitica* (ATCC 9610) or *Shigella flexneri* (ATCC 12022) was
74 grown overnight in a flask containing 80 ml tryptone soya broth (TSB; Oxoid CM 129)
75 shaking at 30°C. The cells were harvested, centrifuged to a pellet (512g, 10 mins, 15°C),
76 washed and re-suspended in peptone water (0.1%). A standard inoculum was produced by
77 diluting the culture to an OD of 0.5 at 600nm. This standardized culture was then further
78 diluted in TSB to produce the starting inoculum (approx 1×10^5 CFU ml⁻¹ in the microtitre
79 plate).

80 *2.2. Experimental method*

81 Experiments were carried out either using half-fold dilutions using the method of [Lambert](#)
82 [& Pearson \(2000\)](#) or by using linear dilutions from stock solutions of sodium chloride or
83 potassium chloride for the effect of individual inhibitors or the method of [Lambert and](#)
84 [Lambert \(2003\)](#) for combined inhibitors.

85 *2.3. Data analyses and model fitting*

86 The data obtained from the Bioscreen are tables of optical density (OD) and time. The time
87 to detection was defined as the time to produce an OD =0.2 at 600nm. The assumption
88 being made was that at an OD =0.2, each well had identical numbers of microorganisms.
89 Furthermore, microscopic checks were performed to see if cell elongation occurred at the
90 highest salt levels used: no such elongation was observed. Data were transformed to the

reciprocal in order to stabilise data variance. Wells which showed no visible growth during the period of the experiment were removed from the analysis (censored data).

Previous publications (e.g. [Lambert and Bidlas 2007](#)) had used a general model for the fitting of time to detection data (TTD). A modified form of this model, which allows for a definitive MIC for individual inhibitors - the linear-exponential model (E-L), was used to analyse the data obtained, Eq.(1).

$$\begin{aligned} &\text{If } [x] = 0, \text{RTD} = P_0 \\ &\text{Else if } [x] < [P_1], \\ &\text{then } \text{RTD} = P_0 \exp\left(-\left(\frac{[x]}{P_1}\right)^{P_2}\right) \\ &\text{Else, } \text{RTD} = \frac{P_0}{e} (1 - P_2 (\log_e [x] - \log_e P_1)) \end{aligned} \quad (1)$$

Where $[x]$ is the concentration of the inhibitor, P_i are experimental parameters and e is the exponential, RTD is the reciprocal of the time to detection (min^{-1}).

The minimum inhibitory concentration was calculated from the parameter values obtained for each inhibitor using

$$\text{MIC} = P_1 \exp\left(\frac{1}{P_2}\right) \quad (2)$$

For combined inhibitors (combinations of NaCl and KCl) the model of [Lambert and Lambert \(2003\)](#) was modified in a similar way to Eq.(1) allowing a definitive growth-no growth boundary to be constructed for combinations, Eq.(3)

113

$$\left\{ \begin{array}{l} \text{if} \quad \sum_{i=1}^n [x_i] = 0, RTD = P_0 \\ \text{else if} \\ \quad \ln[EffC] < 0 \\ \text{then} \\ \quad RTD = P_0 \exp(-[EffC]^{P_c}) \\ \text{else} \\ \quad RTD = \frac{P_0}{e} (1 - P_c \ln[EffC]) \end{array} \right. \quad (3)$$

114 Where $[EffC] = \sum_{i=1}^n \left(\frac{[x_i]}{P_{2i-1}} \right)^{P_{2i}}$ and P_c is the multinomial exponent for the combined system

115 (Lambert and Lambert 2003).

116 Data were fitted to the equation using non-linear regression using the minimised sum of
 117 squares as the search criterion. Analyses were done using the JMP Statistical Software
 118 (SAS Institute Cary, NC).

119

120 3. Results

121 3.1. Growth inhibition by sodium chloride and potassium chloride

122 Three species – the Gram negatives *E. sakazakii* and *A. hydrophila* and the Gram positive
123 *S. aureus* are used below to highlight the results obtained. Three strains of the latter
124 organism were used, due to the importance of humectant activity to control the growth of
125 this organism. The results from these and the other two organisms examined in this study
126 are summarised in [Table 1](#).

127

128 3.2 *Enterobacter sakazakii*

129 Optical density/incubation time data were collected using half-folded dilutions of NaCl or
130 KCl. When analysed using the modified RTD model there was an excellent fit; [Figure 1](#)
131 shows the results for the effect of KCl on the RTD. At KCl concentrations less than 1×10^4
132 mg l^{-1} (1 %), there is little effect on the rate to detection, i.e. shows uninhibited growth.
133 Above 1%, inhibition increases and above $1.2 \times 10^5 \text{ mg l}^{-1}$ (12%) KCl no growth was
134 observed.

135

136 [Table 1](#) gives the experimental parameters found in percent and in mol l^{-1} . In terms of mol
137 l^{-1} the MIC of NaCl and KCl are within the 95% confidence interval of the mean
138 (calculated from the parameters P_1 and P_2) given for each humectant. [Figure 2](#) compares
139 the observed and fitted data for both the NaCl and KCl inhibition. From these results, when
140 expressed in mol l^{-1} , there is no evidence that NaCl and KCl have different inhibitory
141 effects against *E. sakazakii*. It was also found that in both cases there was evidence of a
142 threshold concentration of approx. 0.1M added salt before any observation of growth
143 inhibition.

144

145 3.3. *Staphylococcus aureus*

146 Comparison of NaCl and KCl were carried out on three strains of *S. aureus* (two standard
147 strains and a factory isolate). [Figure 3](#) shows the results of both experiments in terms of
148 mol l⁻¹ for strain ST121. The time to detection was transformed using the natural logarithm.
149 The curvature observed in Fig. 3 is typical for salt inhibition. Similar results were found
150 for the other two *S. aureus* strains examined.

151

152 It can also be observed from the figure that as conditions become more harsh, the
153 variability (or observed error) in the time to detection also increases, even although the
154 logarithm transformation has been used. There is a general observation throughout
155 predictive microbiology that the variance increases as conditions become harsher and
156 without variance stabilisation the accuracy of models is reduced the closer they approach
157 the MIC of the preservative. In this particular case the reciprocal transformation performs
158 better than the logarithmic.

159

160 3.4. *Combination NaCl and KCl experiments*

161 If the hypothesis that NaCl and KCl are mutually replaceable is correct, then the
162 antimicrobial effect of combinations of NaCl and KCl should be predictable. If the
163 calculation is performed using a combined humectant concentration in terms of mol⁻¹ then
164 the modelling is facile. If, however, the separate identities are kept, then a complication
165 arises due to the non-unity dose response. Previous work done on combined hurdles has
166 used the following equation to describe the effect on RTD (Lambert and Bidlas 2007)

167

$$RTD = P_0 \exp \left(- \left(\sum_{i=1}^n \left(\frac{[x_i]}{P_{2i-1}} \right)^{P_{2i}} \right) \right) \quad (4)$$

For combined NaCl and KCl equation (4) will only give an approximation to the observed pattern of inhibition because the parameter P_{2i} for both humectants is approximately 2 and the above equation ignores the binomial expansion of the combined response. Equation (4), which takes this into account was used to examine the observed RTD data for *A. hydrophila* from a chequerboard of NaCl/KCl mixtures (observed data [Figure 4](#), modelled data [Figure 5](#)). There is a very good fit of the model to the observed data. [Figure 6](#) shows a plot of the observed and modelled RTD against the Effective Concentration; that the *EffC* is made up from any combination of NaCl and KCl and that the observations all lie on or close to the modelled line is a simple graphical representation to show that there is no synergy between the two inhibitors, since any ‘true’ synergy would cause a mismatch. Further, the predicted values for the combined experiment (as opposed to the direct modelling of the observed data) are given in [Table 2](#) for both the *A. hydrophila* and the identical *E. sakazakii* experiments. The predicted parameters are in agreement with those modelled from the observed data.

4. Discussion

This work focuses on the ability of potassium chloride to replace salt as an antimicrobial humectant. From this study KCl is a direct 1:1 molar replacement for the antimicrobial effect of common salt. This study enhances the work done by Boziaris et al. (2007) and delivers the same conclusion for a larger spectrum of pathogens. There is still, however, clarification required for some of the other common pathogens as the paper by Strong et al. (1970) would suggest. There is also one point which needs to be made: there is no differentiation in this hypothesis of the equivalence of KCl and NaCl for the antimicrobial effect to be due to the chloride ion. The use of calcium chloride and other similar salts would immediately differentiate these two possibilities.

The model and the definition used for the effective concentration are direct applications of the Gamma hypothesis ([Zwietering](#), Wijtzes, De Wit and Van't Riet [1992](#)) - that individual inhibitors act independently against the growth of microorganisms. Recently, attempts had been made to expand the hypothesis to include the possibility of factor interaction, based on the observation of the shape of the Growth/No growth boundary for combined antimicrobials ([Augustin and Carlier 2000](#); Le Marc, Huchet, Bourgeois, Guyonnet, Mafart and Thuault 2002). The model described here adequately describes the shape of these G/NG boundaries without recourse to altering the Zwietering hypothesis.

[Figure 7](#) shows the calculated Growth/No Growth boundary for combinations of NaCl and KCl for the organisms studied. From equation (3) when the effective concentration is greater than 1, the linear function takes over the description of the level of inhibition. For a given RTD this describes a contour for a twin-mix of inhibitors, a surface for ternary-mixes and hypersurfaces for mixtures with greater numbers of components. The $RTD = 0$ contour, i.e. the absolute growth/no growth boundary is given by

$$RTD = 0 = \frac{P_0}{e} (1 - P_c \ln[EffC])$$

∴

$$EffC = \text{Exp}(1/P_c)$$

For any combination of inhibitors where the calculated effective concentration is equal to $\text{Exp}(1/P_c)$, this combination will define a point on the MIC contour (surface, etc). Any combination with $EffC > \text{Exp}(1/P_c)$ lies in a NG zone, and any with $EffC < \text{Exp}(1/P_c)$ lies in a G zone. For a given organism, conditions beyond the calculated boundary line will result in no growth. If total salt concentrations, for a given product, are within a boundary line, then other preservative factors (e.g. temperature or weak acid preservatives) have to be used to ensure no growth.

Most antimicrobial hurdles examined previously such as pH and weak acids give a linear relationship between the log of the time to detection and the concentration, hence the dose response parameter $P_2 \cong 1$ (Lambert and Bidlas 2007). Combinations of such hurdles will also give $P_c \cong 1$. The humectants used in this study, NaCl and KCl are unusual in that the dose response of each was approximately 2. Dose responses of other antimicrobials can vary substantially e.g. phenolics have dose responses (or dilution coefficients) >6 , (Lambert and Johnston 2000; Lambert and Lambert 2003).

Modelling offers a cost-effective approach to understanding the microbial growth response in foods. Indeed Zwietering (Zwietering et al.1996) has said that using a model to predict the consequences of changing a formulation on microbial growth is a factor of 1000 quicker than attempting a large scale storage trial. Of course the formulator needs access to the model in the first place and must also have an idea of its robustness. However, the use of mathematical models can help to reduce the need for storage trials, challenge tests, product reformulations and process modifications, which are labour intensive, time

232 consuming and expensive. The above example of NaCl vs. KCl is an example whereby a
233 relatively rapid method of obtaining relevant information in conjunction with a robust
234 model leads to a simple algorithm for NaCl replacement, allowing a product developer to
235 gain insight perhaps 1000 times faster.

References

- Augustin, J-C. and Carlier, V., 2000. Modelling the growth rate of *Listeria monocytogenes* with a multiplicative type model including interaction between environmental factors. International Journal of Food Microbiology 56, 53-70.
- Boziaris, I.S., Skandamis, P.N., Anastasiadi, M. and Nychas, G.-J.E., 2007. Effect of NaCl and KCl on fate and growth/no growth interfaces of *Listeria monocytogenes* Scott A at different pH and nisin concentrations. Journal of Applied Microbiology 102, 796-805
- Beuchat, L.R., 1974. Combined Effects of Water Activity, Solute, and Temperature on the Growth of *Vibrio parahaemolyticus*. Applied Microbiology 27 (6) 1075-1080.
- Gimeno, O., Astiasaran, I. and Bello, J., 1999. Influence of partial replacement of NaCl with KCl and CaCl₂ on texture and color of dry fermented sausages. Journal of Agricultural and Food Chemistry 47, 873-877.
- Gimeno, O., Astiasaran, I. and Bello, J., 2001. Influence of partial replacement of NaCl with KCl and CaCl₂ on microbiological evolution of dry fermented sausages. Food Microbiology 18, 329–334.
- Lambert, R.J.W., and Bidlas, E., 2007. An investigation of the Gamma hypothesis: A predictive modelling study of the effect of combined inhibitors (salt, pH and weak

acids) on the growth of *Aeromonas hydrophila*. International Journal of Food Microbiology 115, 12-28.

Lambert, R.J.W. and Johnston, M.D. 2000. Disinfection kinetics: a new hypothesis and model for the tailing of log-survivor/time curves. Journal of Applied Microbiology 88, 907-913.

Lambert, R.J.W., and Lambert, R., 2003. A model for the efficacy of combined inhibitors. Journal of Applied Microbiology 95, 734–743.

Lambert, R.J.W. and Pearson, J., 2000. Susceptibility testing: accurate and reproducible Minimum Inhibitory Concentration, MIC, and Non-Inhibitory Concentration, NIC, values. Journal of Applied Microbiology 88, 784-791.

Le Marc Y., Huchet V., Bourgeois, C.M., Guyonnet, J.P., Mafart, P. and Thuault, D. 2002. Modelling the growth kinetics of *Listeria* as a function of temperature, pH and organic acid concentration. International Journal of Food Microbiology 73, 219– 237.

Strong, D.H., Foster, E.F. and Duncan, C.L. 1970. Influence of water activity on the growth of *Clostridium perfringens*. Applied Microbiology 19 (6) 980-987.

Zwietering, M.H., Wiltjes, T., De Wit, J.C. and van't Riet, K., 1992. A Decision Support System for Prediction of the Microbial Spoilage in Foods. Journal of Food Protection 55, 973-979.

286 Zwietering, M. H., De Wit, J.C. and Notermans, S., 1996. Application of predictive
287 microbiology to estimate the number of *Bacillus cereus* in pasteurised milk at the point
288 of consumption. International Journal of Food Microbiology 30, 55-70.

290 Table 1. Modelled parameters values for NaCl and KCl inhibition for Pathogens used in this
 291 study

Organism	Humectant	P ₁ (mg l ⁻¹) (SErr)	P ₁ (mol l ⁻¹) (SErr)	P ₂ (SErr)	MIC (mg l ⁻¹)	MIC (mol l ⁻¹) (95% CI)
<i>E. sakazakii</i>	NaCl	48180 (160)	0.824 (0.003)	1.609 (0.010)	89810	1.537 (1.516-1.561)
	KCl	62970 (390)	0.845 (0.005)	1.587 (0.0182)	118200	1.586 (1.545-1.630)
<i>A. hydrophila</i> (ATCC 7966)	NaCl	32290 (330)	0.552 (0.006)	2.080 (0.050)	52230	0.893 (0.857-0.932)
	KCl	41870 (410)	0.562 (0.006)	1.910 (0.0395)	70670	0.948 (0.911-0.987)
<i>Y. enterocolitica</i> (ATCC 9610)	NaCl	32700 (262)	0.560 (0.0045)	2.034 (0.061)	53460	0.915 (0.873-0.961)
	KCl	37930 (172)	0.509 (0.0023)	1.859 (0.030)	64950	0.871 (0.848-0.895)
<i>Sh. flexneri</i> (ATCC 12022)	NaCl	37040 (170)	0.634 (0.003)	2.189 (0.036)	58490	1.000 (0.977-1.025)
	KCl	43410 (291)	0.582 (0.004)	1.673 (0.031)	78900	1.0589 (1.022-1.096)
<i>S. aureus 121</i>	NaCl	89600 (500)	1.533 (0.009)	2.190 (0.0476)	141500	2.421 (2.349-2.498)
	KCl	114900 (554)	1.542 (0.007)	2.114 (0.0366)	184400	2.474 (2.411-2.539)
<i>S. aureus 6538</i>	NaCl	88100 (710)	1.508 (0.012)	2.176 (0.0603)	139500	2.388 (2.292-2.488)
	KCl	115700 (590)	1.552 (0.008)	1.984 (0.0363)	191500	2.569 (2.496-2.645)
<i>S. aureus 55</i>	NaCl	87650 (470)	1.500 (0.008)	1.875 (0.0341)	149400	2.557 (2.482-2.634)
	KCl	114400 (800)	1.534 (0.010)	1.817 (0.0408)	198400	2.661 (2.558-2.770)

Table 2. Predicted and fitted parameters for the combined experiment of KCl and NaCl evaluated using equation 3.

Organism		Parameter	Predicted value	Fitted value
<i>A. hydrophila</i>	NaCl	P_1	32290	32880
		P_2	1.000	1.018
	KCl	P_3	41870	43220
		P_4	0.918	0.942
	Combined	P_c	2.080	2.010
<i>E. sakazakii</i>	NaCl	P_1	48180	47670
		P_2	1.000	1.100
	KCl	P_3	62970	58880
		P_4	0.988	1.033
	Combined	P_c	1.606	1.526

P_1 and P_3 values quoted are in mg l^{-1}

Legend to Figures

Figure 1. Effect of KCl on the rate to detection of *Enterobacter sakazakii* in TSB at 30°C.

Figure 2. Comparison of the observed (symbols) and modelled (solid line) NaCl and KCl inhibition of *Enterobacter sakazakii*; □ NaCl; ○ KCl. The models for NaCl and KCl are essentially coincident and only that for NaCl is shown.

[Figure 3](#). Comparison of the observed ln(time) to detection (mins) for NaCl (x) or KCl (○) inhibition of *Staphylococcus aureus* 121(ten replicates per humectant concentration).


Figure 4. Observed RTD for mixtures of NaCl and KCl inhibition of *Aeromonas hydrophila*

Figure 5. Modelled RTD for mixtures of NaCl and KCl inhibition of *Aeromonas hydrophila*

Figure 6. A plot of the effective concentration against the observed (symbols) and modelled RTD (solid line) for combinations of NaCl and KCl against *Aeromonas hydrophila*.


Figure 7. MIC boundaries for NaCl/KCl mixtures; ●, *Staphylococcus aureus* (Saur121); ○, *Enterobacter sakazakii*; ■ *Shigella flexneri*; □, *Aeromonas hydrophila*; ▲, *Yersinia enterocolitica*.

321 *Figure 1.*




322

323 *Figure 2.*





324

325 *Figure 3.*




326






331 *Figure 6.*



332





DSPACE

<https://dspace.org/>

Comparing the antimicrobial effectiveness of NaCl and KCl with a view to salt/sodium replacement.

Bidlas, Eva; Lambert, Ronald J. W.

2008-05-10

Elsevier

<http://dx.doi.org/10.1016/j.ijfoodmicro.2008.02.031>

Eva Bidlas, Ronald J.W. Lambert, Comparing the antimicrobial effectiveness of NaCl and KCl with a view to salt/sodium replacement, International Journal of Food Microbiology, Volume 124, Issue 1, 10 May 2008, Pages 98-102.