

# **Unveiling the effect of interacting forecasted abiotic factors on growth and Aflatoxin B1 production kinetics by *Aspergillus flavus***

**Esther GARCIA-CELA<sup>1,2†</sup> and Carol VERHEECKE-VAESSEN<sup>1,†</sup>, Maria GUTIERREZ- POZO<sup>1,3</sup>,**

**Elisavet KIAITSI<sup>1</sup>, Alessandra MARCON-GASPERINI<sup>1</sup>, Naresh MAGAN<sup>1</sup>, Angel**

**MEDINA\***

<sup>1</sup>Applied Mycology Group, Environment and AgriFood Theme, Cranfield University, Cranfield, Beds. MK43 0AL, UK

<sup>2</sup> Biological and Environmental Sciences, School of Life and Medical Sciences, University of Hertfordshire, Hatfield, AL109AB, UK

<sup>3</sup> Surface Engineering and Precision Institute, Cranfield University, Cranfield, Beds. MK43 0AL, UK

† the two authors contributed equally to the work.

\*Corresponding authors: Dr Angel Medina, Applied Mycology Group, Environment and AgriFood Theme, Cranfield University, Cranfield, Beds. MK430AL, U.K. E-mail: a.medinavaya@cranfield.ac.uk

## ABSTRACT (200 words)

The aim was to decipher the temporal impact of key interacting climate change (CC) abiotic factors of temperature (30 vs 37 °C), water activity ( $a_w$ ; 0.985 vs 0.930) and CO<sub>2</sub> exposure (400 vs 1000 ppm) on (a) growth of *Aspergillus flavus* and effects on (b) gene expression of a structural (*aflD*) and key regulatory gene (*aflR*) involved in aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) biosynthesis and (c) AFB<sub>1</sub> production on a yeast extract sucrose medium over a period of 10 days. *A. flavus* grew and produced AFB<sub>1</sub> very early with toxin detected after only 48 hours. Both growth and toxin production were significantly impacted by the interacting abiotic factors. The relative expression of the *aflD* gene was significantly influenced by temperature; *aflR* gene expression was mainly modulated by time. However, no clear relationship was observed for both genes with AFB<sub>1</sub> production over the experimental time frame. The optimum temperature for AFB<sub>1</sub> production was 30°C. Maximum AFB<sub>1</sub> production occurred between days 4-8. Exposure to higher CO<sub>2</sub> conditions simulating forecasted CC conditions, the amount of AFB<sub>1</sub> produced in elevated temperature (37 °C) was higher than with the optimum temperature (30 °C) showing a potential for increased risk for human/animal health due to higher accumulation of AFB<sub>1</sub>.

## KEYWORDS:

mycotoxins, water activity, toxin gene expression, elevated CO<sub>2</sub>, *aflD*, *aflR*.

## 1 INTRODUCTION

2 *Aspergillus flavus* is an important primary and secondary pathogen of maize, tree nuts and  
3 peanuts and can contaminate these commodities with the class 1 carcinogenic mycotoxin:  
4 Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) (IARC, 2012). It has now been shown that in some Lower Middle Income  
5 Countries (LMICs), especially in West Africa, consumption of a predominantly maize-based diet  
6 can lead to stunting in children due to exposure to AFB<sub>1</sub> (IARC, 2012).

7 Recent forecasts of climate change prediction are suggesting different scenarios of  
8 temperature and CO<sub>2</sub> changes depending on socio-economic behaviour which could lead to  
9 increases from +3.5 to 7.5°C and from 500 to 1200 ppm (Gidden et al., 2018). Moreover, there  
10 have been concerns that extreme weather events could exacerbate the contamination of staple  
11 commodities with mycotoxins including AFB<sub>1</sub>. Some studies have suggested that environmental  
12 stress may lead to increases in mycotoxin production, or indeed a switch to other toxigenic  
13 secondary metabolites (Garcia-Cela et al., 2014; Gilbert et al., 2017; Medina et al., 2014; Medina  
14 et al., 2015 a,c; Verheecke-Vaessen et al., 2019). It has been previously shown that on a conducive  
15 yeast extract sucrose (YES) medium, individual abiotic factors such as temperature, water  
16 availability, or exposure to elevated CO<sub>2</sub> alone have less modifying effects on the ecology of  
17 mycotoxigenic fungi than when combined (Medina et al., 2015a). However, interactions between  
18 these factors may have a synergistic impact on either growth and/or production of mycotoxins.  
19 Recent studies have shown variable effects of such interacting abiotic factors with little effect on  
20 growth or mycotoxin production (e.g. *A. carbonarius*, ochratoxin A) while for others such as *A.*  
21 *westerdijkiae* and *A. flavus* species colonising maize and coffee there was a stimulation of  
22 mycotoxin production, although less effects on growth (Medina et al., 2015a; Akbar et al., 2016).  
23 However, most of these studies have involved a single time point for analyses of impacts of the  
24 interacting abiotic climate-related factors. There is little knowledge of the temporal changes which

1 may occur in growth or mycotoxin production under existing or future climate-related interacting  
2 abiotic factors.

3       The objective of this study was to examine the kinetics of the impact of interacting climate-  
4 related abiotic factors of temperature (30 vs 37 °C), water activity ( $a_w$ ; 0.985 vs 0.930) and CO<sub>2</sub>  
5 exposure (400 vs 1000 ppm) on (a) lag times prior to growth and colonisation rates, (b) relative  
6 temporal expression of structural (*aflD*) and regulatory (*aflR*) biosynthetic genes involved in toxin  
7 production and (c) effects on phenotypic AFB<sub>1</sub> production by *A. flavus* on a conducive YES  
8 medium over a period of 10 days.

## MATERIALS AND METHODS

### *Fungal strains*

An aflatoxigenic type strain of *A. flavus* (NRRL 3357; Northern Regional Research Laboratories (NRRL) of the US Department of Agriculture USDA, New Orleans) was used in these experiments. The strain was maintained in glycerol:water (70:30, v/v) at -20 °C in the culture collection of the Applied Mycology Group, Cranfield University, UK.

### *Media preparation and inoculation*

The water activity ( $a_w$ ) of the media was modified by the addition of different amounts of glycerol to the water which was mixed and used as water to prepare the treatment media (0.985, 0.930). A standard curve was prepared of the amounts of glycerol/water solutions necessary to modify the YES medium to the target treatment  $a_w$  levels. This was added to the YES media (2% yeast extract, 15% sucrose, 0.05%  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ). The accuracy of the modifications was confirmed using an Aqualab 4TE instrument (Decagon, Pullman, WA, US) and found to be within  $\pm 0.01$  of the target  $a_w$  levels.

Spore suspensions of conidia were prepared from 7-days old colonies of *A. flavus* on MEA (3% malt extract, 5% mycological peptone, 1.5% agar) incubated at 25 °C. The conidial concentration was measured using a haemocytometer (Thoma Cell, Fisher Scientific) and the concentration diluted with sterile water to obtain a concentration of  $10^6$  conidia  $\text{mL}^{-1}$ .

The different  $a_w$  agar media were poured into sterile Petri dishes ( $\varnothing$  9 cm) in a sterile safety cabinet and allowed to cool. The treatments and replicates were centrally inoculated with *A. flavus* conidial spore suspensions (5  $\mu\text{l}$ ;  $10^6$  spores  $\text{mL}^{-1}$ ). Three replicates were destructively sampled

for growth measurements, mycotoxins and gene expression analysis every two days over the 10-days period.

### ***Exposure of A. flavus treatments to interacting abiotic factors***

The different treatments and replicates were placed in separate 13 L plastic environmental chambers (Verheecke-Vaessen et al., 2019). These chambers also contained an inlet and outlet with valves for flushing with CO<sub>2</sub>. The chambers contained glycerol/water solutions to maintain the ERH of the atmosphere within the individual chambers to maintain the target  $a_w$  levels. The chambers were flushed with either air or 1000 ppm CO<sub>2</sub> daily for 10 days. The gas cylinders contained either 400 ppm CO<sub>2</sub> (ambient air) or air supplemented with CO<sub>2</sub> to obtain a 1000 ppm CO<sub>2</sub> concentration from a speciality certified gas cylinder (British Oxygen Company, Guildford, Surrey, U.K.). The containers were flushed at 3 L min<sup>-1</sup> to replace 3x the volume of the incubation chamber. The chambers were incubated at either 30 or 37°C.

To calculate the fungal colonisation rates the diametric size of the colonies was measured daily in two perpendicular directions. Every two days, three replicates were destructively sampled and the fungal biomass harvested by scratching the plate surface and stored at -80 °C for gene expression analysis and 2-5 plugs (using a 5 mm surface-sterilised cork borer) across the *A. flavus* colony and media were collected. The plugs were placed in a pre-weighed 2 mL Eppendorf tube, weighed again and sealed before being stored at -20 °C for mycotoxin analysis.

### ***AFB<sub>1</sub> extraction and quantification***

Samples were thawed and extracted by mixing the agar plugs with 750 µL of methanol. The tubes were shaken for 90 min at 400 g at 25 °C in the dark. They were then centrifuged at 13,000 g for 15 min and all of supernatant removed and the plugs were re-extracted again in the same way. Extracts were dried in a miVac evaporator (Genevac, Ipswich, UK) for 6 h. The samples

were resuspended in 1000  $\mu$ L of methanol:water (50:50, v:v) and filtered through a 0.22  $\mu$ m Nylon-filter (type) into HPLC vials and stored at  $-20^{\circ}\text{C}$  until analyses.

The samples were injected into a HPLC-FLD Agilent 1200 Series HPLC system (Agilent Technologies, Palo Alto, CA, USA) equipped with post-column derivatisation with a UVE photochemical reactor with UV-Light (LCTech GmbH, Germany) and a fluorescence detector ( $\lambda_{\text{exc}}$  360 nm;  $\lambda_{\text{exc}}$  440 nm). A  $\text{C}_{18}$  column (Agilent Zorbax® Eclipse Plus, 2.1x100mm, 3.5  $\mu$ m particle size) preceded by a Phenomenex® Gemini  $\text{C}_{18}$  guard column cartridge 3 mm x 3  $\mu$ m (Phenomenex, California, USA) was used for separation. Isocratic elution with methanol:water:acetonitrile (30:60:15, v/v/v) mobile phase and the flow rate at 1.0 mL  $\text{min}^{-1}$  were used. AFB<sub>1</sub> was quantified on the base of the HPLC fluorimetric response compared with a range of mycotoxin standards supplied by Romer Labs (Romer Labs Ltd, UK) (0.5-2,010 ng mL<sup>-1</sup>,  $R^2=0.99$ ). Samples outside of the range were appropriately diluted with methanol:water (50:50, v:v). AFB<sub>1</sub> retention time was 9.5 min and the limit of quantification (LOQ) was 0.5 ng mL<sup>-1</sup> based on a signal-to-noise ratio 10:1. Signals were processed using the Agilent Chem-Station software Ver. B Rev: 03.01 [317] (Agilent Technologies, Palo Alto, CA, USA). Method validation and recovery were determined by directly spiking both media with AFB<sub>1</sub> standard solutions. Once autoclaved, the media was cooled down to 56  $^{\circ}\text{C}$ , then AFB<sub>1</sub> standard was added onto the media, mixed vigorously and poured in Petri plates ( $\varnothing$  5 mm). After media solidification three agar plugs were collected across the plate and the mycotoxins extracted as described previously. The recovery for AFB<sub>1</sub> for three replicates was 88.3% $\pm$  4.9.

### ***Relative expression of the aflD and aflR genes***

Fungal RNA was extracted from the fungal biomass using the method previously described (Verheecke-Vaessen et al., 2019). The RNA obtained was checked for integrity by electrophoresis

(a ratio of intensity 25S/18S higher than 2 was visually observed) and purity ( $\text{ratio}_{260/280}=2-2.2$ ,  $\text{ratio}_{260/230}>2$ ) by Genova Nano (Jenway, Stone, UK), and stored at  $-80\text{ }^{\circ}\text{C}$ . Reverse transcription was performed using the Omniscript<sup>®</sup> Reverse Transcription Kit (Qiagen, Hilden, Germany) using Oligo(dT)<sub>18</sub> following the manufacturers' instructions. The qPCR targeting *aflD*, *aflR* and  *$\beta$ tub* (reference gene) was performed as described previously (Medina et al., 2015c).

## ***Statistical Analysis***

JMP<sup>®</sup> 14 (SAS Institute Inc., 2016. Cary, NC, USA) was used to perform statistical analyses. Normality and homoscedasticity of each dataset (lag time, growth rate, aflatoxins and gene expression) was checked using Shapiro-Wilk test and Brown-Forsythe test. Lag time, gene expression and growth rate (square root transformed) data showed both normality and homoscedasticity; thus ANOVA was used to investigate significant differences. A Student *t*-test was done to compare each pair ( $p\text{-value} \leq 0.05$ ).

For AFB<sub>1</sub> production data and gene expression, non-normality or variance homogeneity was confirmed even after multiple transformation trials were performed. Thus, a non-parametric test, Kruskal-Wallis, was performed. When significant differences were found ( $p\text{-value} \leq 0.05$ ), each pair were compared by a *post-hoc* Wilcoxon method. For gene expression and toxin production comparison, the Spearman test was used.

## RESULTS

### *Effect of three-way interacting climate change environmental factors on A. flavus growth on YES media*

The lag phases prior to growth were shorter as  $a_w$  was increased (x 2 to x 4 fold; Table 1). The exposure to higher CO<sub>2</sub> generally reduced these lag times. This was significant at 37 °C, 0.930 with a 3.6-fold reduction in lag phase due to the higher CO<sub>2</sub> exposure. Colonisation was significantly higher at 0.985  $a_w$ . The optimal growth occurred at 30 °C and 0.985  $a_w$ . Generally, the elevated CO<sub>2</sub> significantly reduced the colonisation rate, except at 37 °C and 0.985  $a_w$ . Under drier conditions (0.93  $a_w$ ) and elevated CO<sub>2</sub> the growth rates were the lowest at both temperatures.

The statistical analysis compared the effect of interacting abiotic climate-related factors on the lag phases prior to growth and growth rates. Statistically, temperature (30 vs 37 °C),  $a_w$  (0.985 vs 0.930) and CO<sub>2</sub> exposure (400 vs 1000 ppm) effects on growth rates were compared. Both lag times prior to growth and growth rates were significantly affected ( $p$ -value = <0.0003 and  $p$ -value = <0.0001, respectively). Temperature had a significant impact on the lag phase ( $p$ -value = <0.0086) and  $a_w$  had a significant effect on growth rate ( $p$ -value = <0.0001) and CO<sub>2</sub> to a lesser extent ( $p$ -value = 0.081).

### *Influence of climate change related abiotic factors on kinetics of the expression of the AFB<sub>1</sub> biosynthetic genes aflD and aflR*

The effect of temperature (30 and 37 °C),  $a_w$  (0.930 and 0.985) and CO<sub>2</sub> exposure (400 and 1000 ppm) on the kinetics of *aflD* and *aflR* gene expression was studied and the results are presented in Figure 1 and Supplementary Figure 1. Overall, the gene expression pattern of both genes was significantly impacted by climate change interacting abiotic factors.

For the structural gene *aflD*, an increase in temperature led to a general decrease of gene expression over the experimental period, independently of the  $a_w$  studied. Comparisons between current conditions (30 °C; 0.985; 400 ppm; Figure 1b) and future scenarios (37 °C; 0.93; 1000 ppm; Figure 1c) showed a general decrease in *aflD* gene expression especially after 8 and 10 days. For the biosynthetic gene *aflR*, encoding the aflatoxin transcription factor, gene expression was generally down regulated through time except at 37 °C/0.985 in elevated CO<sub>2</sub> (Supplementary Figure 1). The down regulation was higher at 0.930  $a_w$  independently of the temperature tested. Generally, the downregulation increased over time at 30 °C, while this decreased over time at 37 °C (Figure 1). When looking into the effects of time, the gene expression of both the *aflD* and *aflR* genes was highest on day 4, regardless of the treatment (see Figure 1). By day 4, under current CO<sub>2</sub> concentration (400 ppm) and 30 °C and 0.930  $a_w$ , there was a significant increase in the expression of the *aflR* gene (Figure 1a). In contrast, the structural *aflD* gene expression was generally higher after 6 to 10 days growth when compared with day 4.

Interestingly, exposure of *A. flavus* colonies to elevated CO<sub>2</sub> resulted in a significant reduction in the *aflR* gene expression after 4 days, with no subsequent temporal changes. Conversely, *aflD* gene expression was optimum after day 4-6. At 37 °C in the 0.930  $a_w$  (see Figure 1c) treatment, the gene expression was higher after 4 days for both genes.

Further analysis was performed to study the impact of the three-way interacting climate change factors on gene expression on the YES media (Table 2). *aflD* gene expression was significantly impacted by temperature ( $p$ -value = <0.0001) and *aflR* gene expression might be mainly modulated by time ( $p$ -value = 0.0729).

***Effect of three-way interacting climate change environmental factors on the AFB<sub>1</sub> temporal production on YES***

Figure 2 shows the effect of the interacting abiotic climate-related abiotic factors on AFB<sub>1</sub> production. Overall, maximum production occurred at 30 °C. Under current CO<sub>2</sub> conditions (400 ppm), maximum production was found after 6 days at 30 °C and 0.930 a<sub>w</sub> (435.7 µg g<sup>-1</sup>). When *A. flavus* was exposed to elevated CO<sub>2</sub> conditions, the maximum production (117.5 µg g<sup>-1</sup>) was also found under the same temperature and a<sub>w</sub> levels but was lower and delayed until 8 days incubation.

Generally, it was observed that elevated CO<sub>2</sub> enhanced AFB<sub>1</sub> production at 37 °C in the dryer treatment (0.930 a<sub>w</sub>). AFB<sub>1</sub> synthesis appeared to occur very early in the development of the colonies of *A. flavus* from conidial inoculum as levels of the toxins were found under all conditions after 2 days incubation.

Over time, at 30 °C, the accumulation of AFB<sub>1</sub> with 400 ppm of atmospheric CO<sub>2</sub> was consistently higher than the accumulation of toxin under elevated CO<sub>2</sub> conditions with a few exceptions (day 8, 0.985 a<sub>w</sub>). Interestingly, this trend was completely reversed when the fungus was exposed to temperature stress conditions (37 °C), where additional atmospheric CO<sub>2</sub> led to higher (at 0.930 a<sub>w</sub>) and earlier (at day 2, 0.985 a<sub>w</sub>) accumulation of AFB<sub>1</sub> at both a<sub>w</sub> levels tested.

## DISCUSSION

This study has addressed one of the current gaps in knowledge with regards to the impact of interacting abiotic factors on the kinetics of growth and AFB<sub>1</sub> production by *A. flavus*. Most of the previous studies have only analysed effects after a single time point which may not provide the necessary insights into the mechanisms involved in growth, biosynthetic gene expression and mycotoxin production (Gilbert et al., 2017; Medina et al., 2015a,b,c; 2017; Verheecke et al., 2015).

In this work, it was found that an increase in atmospheric CO<sub>2</sub> modified the overall lag time values by 0.1 to 0.5 days and maximum growth rates by <1 mm diameter/day. It was clear that the lag times were significantly influenced, especially by drier conditions (0.930 a<sub>w</sub>) and under high temperature (37 °C) where the addition of CO<sub>2</sub> shortened the lag times significantly. However, these conditions did not influence growth significantly and the difference in growth rate between elevated and non-elevated CO<sub>2</sub> conditions was only 0.66 mm diameter day<sup>-1</sup>. This suggests that under combined water and temperature stress conditions elevated CO<sub>2</sub> induces earlier growth of *A. flavus* and hence will confer this species with a competitive advantage under such environments.

Previous studies in relation to three-way interacting climate-related abiotic changes on both YES and maize-based media and maize kernels have suggested very little effect on colonisation rates. However, in ripening maize cobs it has been shown that colonisation by *A. flavus* is similar at both 30 and 37 °C and indeed that AFB<sub>1</sub> production may be even higher at 37 °C in the milky ripe and dough stages (a<sub>w</sub> range: 0.958-0.985; Marcon-Gasperini et al., 2019). However, interaction with elevated CO<sub>2</sub> was not included in these specific studies. Previous studies with *A. flavus* on YES media under 3-way (temperature, a<sub>w</sub> and CO<sub>2</sub> concentration) climate-related abiotic conditions showed relatively minimal effect on growth rates (6.5-12 Ø mm day<sup>-1</sup>; a<sub>w</sub> 0.97-0.92 at 34 and 37 °C) (Medina et al., 2015c).

1       The relative expression of an early structural gene and a regulatory gene (*aflD*, encoding  
2   the norsolorinic acid reductase; *aflR*, encoding the main aflatoxin cluster transcription factor,  
3   respectively) were found to be very variable in the present study with little direct correlation with  
4   phenotypic AFB<sub>1</sub> production. Overall, the relative expression of the *aflD* gene was mostly  
5   modulated by temperature, with little effect of elevated CO<sub>2</sub> or drought stress. For the *aflR* gene  
6   expression, this was predominantly influenced by time of analysis over the 10 day period,  
7   regardless of the climate-related abiotic factors. Usually, biosynthesis of aflatoxins requires the  
8   switching on of the early structural genes (including *aflD*) and the biosynthetic gene regulators  
9   (*aflR*, *aflS*). The results suggest that *aflD* is switched on very early during the growth process  
10   facilitating subsequent downstream biosynthetic genes being activated. This hence suggests that  
11   the activation of *aflR*, which is the transcription factor of the cluster, and hence *aflS*, might occur  
12   even earlier than the first sampling point in this work and thus resulting in aflatoxins production  
13   by day 2. We have observed a general trend indicating that over several days, there is a reduction  
14   in expression of both *aflD* and *aflR* with a concomitant subsequent increase in toxin production  
15   when the fungus was exposed to the higher CO<sub>2</sub> levels. Thus, complementary studies including  
16   earlier time points and a larger set of biosynthetic genes would be beneficial for further elucidation  
17   of how AFB<sub>1</sub> production is modulated.

18       Previous studies have shown that early biosynthetic gene expression prior to mycotoxin  
19   production occurs in mycotoxigenic fungi generally (Geisen et al., 2004). Studies have also  
20   suggested that the impact of two-interacting abiotic factors influences the *aflR/aflS* regulatory gene  
21   ratio and this would indicate the effect on aflatoxins production (Schmidt-Heydt et al., 2010)

22       The present study has shown that AFB<sub>1</sub> production kinetics is initiated very early in the  
23   growth phase (2 days). Subsequently, there is an increase in AFB<sub>1</sub> production over the 10 day  
24   incubation period. At elevated temperature and CO<sub>2</sub> levels, under water stress conditions there was

1 a stimulation of AFB<sub>1</sub> production. This suggests that under combined stress conditions there may  
2 be a different temporal effect on mycotoxin production as suggested previously by Schmidt-Heydt  
3 et al. (2011). In addition, previously Medina et al. (2015c) showed that there was a stimulation of  
4 AFB<sub>1</sub> production under elevated CO<sub>2</sub> and 37 °C when compared to 30 °C and 400 ppm CO<sub>2</sub>. At  
5 0.930 a<sub>w</sub>, maximum production values were observed after 4-8 days, after which there was a  
6 decrease. With freely available water, results are different with little difference between climate-  
7 related treatments.

8         The results presented in this study suggest that earlier colonisation and the early production  
9 of AFB<sub>1</sub> in the life cycle could be important for their survival, especially the saprophytic  
10 component, where abiotic stresses are often extreme allowing effective competitive exclusion of  
11 other microorganisms and subsequent production of both asexual and sexual spores for infection  
12 of ripening maize crops under conducive environmental conditions.

13         In conclusion, this kinetic study has shown that genes involved in the biosynthesis of  
14 aflatoxins are switched on very early in the growth phase of *A. flavus* under both existing and  
15 future climate-related abiotic factors. Although generally there was an increase in the lag phase  
16 under 3-way interacting stress conditions growth rates were relatively unaffected. This suggests a  
17 very good resilience of *A. flavus* under climate-stress abiotic factors and will allow this species to  
18 produce afltoxins over a long period of time, even at 37 °C, elevated CO<sub>2</sub> and drought-stress  
19 conditions. There was no direct correlation between biosynthetic gene expression for a structural  
20 and biosynthetic gene and AFB<sub>1</sub> production over the 10 day experimental period. This suggests a  
21 complex interaction between interacting climate-related abiotic factors and the ecology of *A.*  
22 *flavus*. The resilience of *A. flavus* under such stress conditions suggests that there may be increased  
23 risks of mycotoxins under such conditions. Such datasets will be useful to inform predictions of

- 1 the relative risk of aflatoxins contamination of staple cereals such as maize and potential impacts
- 2 on the food safety agenda.
- 3

1   **ACKNOWLEDGEMENTS**

2   The authors of the paper would like to thank Fernando Rubio-Lopez (PhD student) for his help  
3   during the mycotoxins extractions and Rodrigo Monteirinho Leitao for his help in the RNA  
4   extractions.

5

6

## LIST OF FIGURES

**Figure 1** – Impact of three-way interacting Climate change factors on the gene expression of *aflD* and *aflR* in *Aspergillus flavus*. For each gene, the data with different letters are significantly different.

**Figure 2** – Impact of three-way interacting Climate change factors on the AFB<sub>1</sub> production on YES media for 10 days. Bars indicate standard error of 3 replicates. X indicates missing values.

## Supplementary figures

**Supplementary Figure 1:** Supplementary Figure 1 – Impact of CO<sub>2</sub> (1000 ppm) exposure on the gene expression of *aflD* and *aflR* in *Aspergillus flavus* at 37 °C , 0.985 a<sub>w</sub>. For each gene, the data with different letters are significantly different.

## LIST OF TABLES

**Table 1.** Effect of interacting abiotic factors including medium, temperature, water activity and CO<sub>2</sub> on *A. flavus* lag times prior to growth and growth rate.

**Table 2.** Comparison of the effect of environmental factors on *aflD* and *aflR* gene expression of *Aspergillus flavus*. \*denotes significant values ( $p$ -value<0.05).

## REFERENCES

- Akbar, A., Medina, A., & Magan, N. (2016). Impact of interacting climate change factors on growth and ochratoxin A production by *Aspergillus* section *Circumdati* and *Nigri* species on coffee. *World Mycotoxin Journal*, 9(5), 863-874.
- Astoreca, A., Vaamonde, G., Dalcero, A., Ramos, A.J., Marín, S., 2012. Modelling the effect of temperature and water activity of *Aspergillus flavus* isolates from corn. *Int. J. Food Microbiol.* 156, 60–67. <https://doi.org/10.1016/J.IJFOODMICRO.2012.03.002>
- Bernaldez, V., Córdoba, J.J., Magan, N., Peromingo, B., Rodríguez, A., 2017. The influence of ecophysiological factors on growth, aflR gene expression and aflatoxin B<sub>1</sub> production by a type strain of *Aspergillus flavus*. *LWT - Food Sci. Technol.* 83, 283–291. <https://doi.org/10.1016/j.lwt.2017.05.030>
- García-Cela, E., Crespo-Sempere, A., Gil-Serna, J., Porqueres, A., & Marin, S. (2015). Fungal diversity, incidence and mycotoxin contamination in grapes from two agro-climatic Spanish regions with emphasis on *Aspergillus* species. *Journal of the Science of Food and Agriculture*, 95(8), 1716-1729.
- Gasperini, A. M., Rodriguez-Sixtos, A., Verheecke-Vaessen, C., Garcia-Cela, E., Medina, A., & Magan, N. (2019). Resilience of Biocontrol for Aflatoxin Minimization Strategies: Climate Change Abiotic Factors May Affect Control in Non-GM and GM-Maize Cultivars. *Frontiers in Microbiology*, 10, 2525.
- Geisen, R., Mayer, Z., Karolewicz, A., Färber, P., 2004. Development of a Real Time PCR system for detection of *Penicillium nordicum* and for monitoring ochratoxin A production in foods by targeting the ochratoxin polyketide synthase gene. *Syst. Appl. Microbiol.* 27, 501–507. <https://doi.org/10.1078/0723202041438419>
- Gidden, M.J., Riahi, K., Smith, S.J., Fujimori, S., Luderer, G., Kriegler, E., van Vuuren, D.P., van

den Berg, M., Feng, L., Klein, D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O., Harmsen, M., Hasegawa, T., Havlik, P., Hilaire, J., Hoesly, R., Horing, J., Popp, A., Stehfest, E., Takahashi, K., 2018. Global emissions pathways under different socioeconomic scenarios for use in CMIP6: a dataset of harmonized emissions trajectories through the end of the century. *Geosci. Model Dev. Discuss.*, 1–42. <https://doi.org/10.5194/gmd-2018-266>

Gilbert, M., Medina, A., Mack, B., Lebar, M., Rodríguez, A., Bhatnagar, D., Magan, N., Obrian, G., Payne, G., 2017. Carbon Dioxide Mediates the Response to Temperature and Water Activity Levels in *Aspergillus flavus* during Infection of Maize Kernels. *Toxins (Basel)*. 10, 5. <https://doi.org/10.3390/toxins10010005>

IARC, 2012. A Review of Human Carcinogens: Chemical Agents and Related Occupations. Lyon, France.

Medina, A., Rodríguez, A., Magan, N., 2014. Effect of climate change on *Aspergillus flavus* and aflatoxin B1 production. *Front. Microbiol.* 5, 348. <https://doi.org/10.3389/fmicb.2014.00348>

Medina, Á., Rodríguez, A., Magan, N., 2015 a. Climate change and mycotoxigenic fungi: impacts on mycotoxin production. *Curr. Opin. Food Sci.* 5, 99–104. <https://doi.org/10.1016/j.cofs.2015.11.002>

Medina, A., Schmidt-Heydt, M., Rodríguez, A., Parra, R., Geisen, R., Magan, N., 2015 b. Impacts of environmental stress on growth, secondary metabolite biosynthetic gene clusters and metabolite production of xerotolerant/xerophilic fungi. *Curr. Genet.* 61, 325–334. <https://doi.org/10.1007/s00294-014-0455-9>

Medina, A., Rodríguez, A., Sultan, Y., Magan, N., 2015c. Climate change factors and *Aspergillus flavus* : effects on gene expression, growth and aflatoxin production. *World Mycotoxin J.* 8, 171–179. <https://doi.org/10.3920/WMJ2014.1726>

Medina, A., Gilbert, M.K., Mack, B.M., OBrian, G.R., Rodríguez, A., Bhatnagar, D., Payne, G.,

- 1       Magan, N., 2017. Interactions between water activity and temperature on the *Aspergillus*  
2       *flavus* transcriptome and aflatoxin B1 production. Int. J. Food Microbiol. 256, 36–44.  
3       <https://doi.org/10.1016/J.IJFOODMICRO.2017.05.020>
- 4       Schmidt-Heydt, M., Rüfer, C.E., Abdel-Hadi, A., Magan, N., Geisen, R., 2010. The production of  
5       aflatoxin B<sub>1</sub> or G<sub>1</sub> by *Aspergillus parasiticus* at various combinations of temperature and  
6       water activity is related to the ratio of aflS to aflR expression. Mycotoxin Res. 26, 241–246.  
7       <https://doi.org/10.1007/s12550-010-0062-7>
- 8       Verheecke-Vaessen, C., Diez-Gutierrez, L., Renaud, J., Sumarah, M., Medina, A., Magan, N.,  
9       2019. Interacting climate change environmental factors effects on *Fusarium langsethiae*  
10       growth, expression of Tri genes and T-2/HT-2 mycotoxin production on oat-based media and  
11       in stored oats. Fungal Biol. <https://doi.org/10.1016/j.funbio.2019.04.008>
- 12       Verheecke, C., Liboz, T., Anson, P., Diaz, R., Mathieu, F., 2015. Reduction of aflatoxin production  
13       by *Aspergillus flavus* and *Aspergillus parasiticus* in interaction with *Streptomyces*. Microbiol  
14       soc 161, 967–972. <https://doi.org/10.1099/mic.0.000070>

Table 1. Effect of interacting abiotic factors including temperature, water activity and CO<sub>2</sub> on *A. flavus* lag time and growth rate.


| Temperature | Water activity (a <sub>w</sub> ) | CO <sub>2</sub> (ppm) | Lag phase (days)             | Diametric growth rate (mm day <sup>-1</sup> ) |
|-------------|----------------------------------|-----------------------|------------------------------|-----------------------------------------------|
| 30°C        | 0.985                            | 400                   | 0.55 ± 0.04 <sup>c,d</sup>   | 12.33 ± 0.76 <sup>a</sup>                     |
|             |                                  | 1000                  | 0.44 ± 0.11 <sup>d,e</sup>   | 11.25 ± 0.21 <sup>b</sup>                     |
|             | 0.93                             | 400                   | 1.02 ± 0.03 <sup>a</sup>     | 6.76 ± 0.32 <sup>f,g</sup>                    |
|             |                                  | 1000                  | 0.84 ± 0.11 <sup>a,b</sup>   | 6.64 ± 0.46 <sup>f,g</sup>                    |
| 37°C        | 0.985                            | 400                   | <0.1 <sup>g</sup>            | 8.87 ± 0.83 <sup>d</sup>                      |
|             |                                  | 1000                  | 0.21 ± 0.06 <sup>e,f,g</sup> | 10.1 ± 0.17 <sup>c</sup>                      |
|             | 0.93                             | 400                   | 0.72 ± 0.07 <sup>b,c</sup>   | 7.18 ± 0.21 <sup>e,f</sup>                    |
|             |                                  | 1000                  | 0.20 ± 0.14 <sup>e,f,g</sup> | 6.52 ± 0.10 <sup>g</sup>                      |


<0.1 denotes that no Lag time calculated. Different letters indicate statistically significant differences between conditions.


Table 2. Results comparing the effect of factors on *aflD* and *aflR* gene expression in *Aspergillus flavus* \*denotes significant values ( $p$ -value<0.05).

| Source          | <i>aflD</i> |                | <i>aflR</i> |                |
|-----------------|-------------|----------------|-------------|----------------|
|                 | ChiSquare   | Prob>ChiSquare | ChiSquare   | Prob>ChiSquare |
| Time            | 1.4761      | 0.6878         | 6.9685      | 0.0729         |
| Temperature     | 18.1896     | <0.0001*       | 2.2509      | 0.1335         |
| Water activity  | 2.7360      | 0.0981         | 1.3864      | 0.2390         |
| CO <sub>2</sub> | 1.0561      | 0.3041         | 0.4194      | 0.5172         |

Figure 1– Impact of three-way interacting Climate change factors on the gene expression of *aflD* and *aflR* in *Aspergillus flavus*. For each gene, the data with different letters are significantly different.







Supplementary Figure 1. Impact of elevated CO<sub>2</sub> (1000 ppm) exposure on the gene expression of *aflD* and *aflR* in *Aspergillus flavus* at 37°C, 0.985. For each gene, the data with different letters are significantly different (P=0.05).



**DSPACE**

<https://dspace.org/>

## **Unveiling the effect of interacting forecasted abiotic factors on growth and Aflatoxin B1 production kinetics by *Aspergillus flavus***

**García-Cela, Esther; Verheecke-Vaessen, Carol; Gutiérrez-Pozo, Maria; Kiaitsi, Elisavet; Gasperini, Alessandra M. ...**

**2020-05-30**

Elsevier

<https://doi.org/10.1016/j.funbio.2020.05.003>

García-Cela E, Verheecke-Vaessen C, Gutiérrez-Pozo M, et al., (2021) Unveiling the effect of interacting forecasted abiotic factors on growth and Aflatoxin B1 production kinetics by *Aspergillus flavus*. *Fungal Biology*, Volume 125, Issue 2, February 2021, pp. 89-94