

Identifying uncertainty in environmental risk assessments: the development of a novel typology and its implications for risk characterisation.

Daniel JC Skinner^a, Sophie A Rocks^{b*}, Simon JT Pollard^c and Gillian H Drew^d

Cranfield University, Collaborative Centre of Excellence in Understanding and Managing Natural and Environmental Risks, Cranfield, Bedfordshire, MK43 0AL, UK.

* Corresponding author

^a tel: +44 1234 750111 email: d.j.skinner@cranfield.ac.uk

^b tel: +44 1234 750111 x2370 email: s.rocks@cranfield.ac.uk

^c tel: +44 1234 754101 email: s.pollard@cranfield.ac.uk

^d tel: +44 1234 750111 x2718 email: g.h.drew@cranfield.ac.uk

Running head: Identifying uncertainty in environmental risk assessments

ABSTRACT

Environmental risk analysts need to draw from a clear typology of uncertainties when qualifying risk estimates and/or significance statements about risk. However, categorisations of uncertainty within existing typologies are largely overlapping, contradictory, and subjective, and many typologies are not designed with environmental risk assessments (ERAs) in mind. In an attempt to rectify these issues, this research provides a new categorisation of uncertainties based, for the first time, on the appraisal of a large subset of ERAs, namely 171 peer-reviewed environmental weight-of-evidence assessments. Using this dataset, a defensible typology consisting of seven types of uncertainty (data, language, system, extrapolation, variability, model, and decision) and 20 related sub-types is developed. Relationships between uncertainties and the techniques used to manage them are also identified and statistically evaluated. A highly preferred uncertainty management option is to take no action when faced with uncertainty, although where techniques are applied they are commensurate with the uncertainty in question. Key observations are applied in the form of guidance for dealing with uncertainty, demonstrated through ERAs of genetically modified higher plants in the EU. The presented typology and accompanying guidance will have positive implications for the identification, prioritisation, and management of uncertainty during risk characterisation.

Keywords: uncertainty, typology, environmental, risk, assessment

INTRODUCTION

Uncertainties within environmental risk assessments (ERAs) need to be properly managed to enable risk estimates to be used as a sound basis for risk management actions (van der Sluijs *et al.* 2004; Refsgaard *et al.* 2007). National and international regulatory bodies stress the importance of acknowledging and dealing with uncertainty in ERAs during the risk characterisation phase (Fairman *et al.* 1998; USEPA 1998; DEFRA 2011). Implementing such guidance starts by identifying potential types of uncertainty (Morgan *et al.* 1990), at which point it is essential that environmental risk analysts are able to draw from a clear and defensible typology of uncertainties (Knol *et al.* 2009; Ramirez *et al.* 2012). Existing typologies have limitations, relating primarily to research domain transferability and content reliability (Walker *et al.* 2003; Ascough II *et al.* 2008; Knol *et al.* 2009; Troldborg 2010). In this paper, we present the development of an evidence-based typology of potential uncertainties in ERAs which, together with implementation guidance, aims to resolve the issues surrounding existing categorisations and better equip environmental risk analysts when attempting to identify and manage uncertainty.

There are a wide range of different types of evidence that can be used to formulate and evaluate risk estimates within ERAs (e.g. toxicological, biological, financial). In some situations, different lines of evidence are amalgamated and the degree to which they support or refute hypotheses about risk is evaluated (Linkov *et al.* 2009). This process, termed weight of evidence (WOE), aims to provide either a definitive course of action for decision-makers where the evidence may be contradictory, or identifies missing information needed to form a definitive conclusion (Chapman 2007). WOE can be applied to ERAs (as well as to ecological or human assessments), but is not recognised as being a specific type of ERA (Suter II and Cormier 2011) and is not consistently defined (Weed 2005). ERAs that apply WOE methods follow the same four phases (problem formulation, exposure assessment,

effects assessment, and risk characterisation; DEFRA 2011) as ERAs that do not use WOE methods, and can therefore be used to identify a useful and manageable dataset to assess how uncertainty is categorised and managed across the much larger set of available ERAs in different risk domains.

It is largely agreed that environmental uncertainty is comprised of different aspects, commonly termed dimensions (Janssen *et al.* 2003; Walker *et al.* 2003; Knol *et al.* 2009). These dimensions relate to: the inherent nature of the uncertainty, either epistemic (limitations in our knowledge) or aleatory (the randomness of natural systems and their components); the severity of the uncertainty, ranging from deterministic treatment at one end of the spectrum to indeterminacy at the other; and the location of the uncertainty, which describes where, in applied situations, the uncertainty manifests. As different uncertainties must be managed differently using different techniques (van der Sluijs *et al.* 2004; Refsgaard *et al.* 2007), identifying the different types of uncertainties that exist in applied situations is an essential part of uncertainty management (Morgan *et al.* 1990). A typology of uncertainty can aid this process by providing comprehensive, relevant, and reliable categorisations (complete with definitions) of all potential types of uncertainty that may be encountered (van Asselt and Rotmans 2002; Knol *et al.* 2009). However, existing typologies are based on small-scale literature reviews, amalgamations of existing frameworks, or researcher opinion (Table 1). As a result, the typologies often contain contradictory definitions and terms, communicate varying frequencies of uncertainty, are rarely comprehensive within their intended research domains, and do not include a clear method for the collection and collation of the evidence base. Furthermore, whilst these typologies may be applicable in a wider risk-context, they are not designed specifically for use with ERAs. Since the overall reliability of a typology relies on the legitimacy of the adopted categorisation(s), in the context of ERAs, it is crucial that this is rectified.

[TABLE 1 NEAR HERE]

This research offers a new categorisation of uncertainties based on the appraisal of a large subset of ERAs in which uncertainties have been transparently identified. As the evidence base is formed of peer-reviewed environmental ERAs that feature WOE methods, the assertions made in this article span a diverse set of interests, making the resulting typology relevant across a number of distinct risk-based research domains. The typology is combined with an analysis of the adoption of uncertainty management techniques (UMTs) used when faced with different uncertainties, and guidance for dealing with uncertainty drawn from key observations.

Uncertainty analysis should be a principal component of risk characterisation and thus ERAs. In reality, this is rarely the case. The introduction of an uncertainty typology that consults a manageable subset of the vast available ERA evidence base, coupled with prioritised guidance, will assist risk analysts in their attempts to prioritise, identify, and manage uncertainties within applied ERAs.

METHODOLOGY

Building the Evidence Base

In order to categorise uncertainty in ERAs that feature WOE methods (hereafter termed WOE-ERAs) and analyse the use of techniques in their management, an evidence base of peer-reviewed literature was established. Searches were conducted for directly labelled WOE-ERA literature, using the ISI Web of Science and Scopus academic databases, respectively, and using the terms *weight*, *evidence*, *risk*, and *uncertainty* (in the *title*, *abstract* and *keywords* field for Scopus, and in the *topic* field for ISI Web of Science). Non-labelled WOE-ERA literature was also searched for, using the terms *risk*, *assessment*, and *uncertainty* (in the *title*, *abstract* and *keywords* field for Scopus, and in the *topic* field for ISI Web of

Science). In-built filtering within the online databases was used to remove obviously non-relevant literature (non-English articles, book series articles, articles from the domains of social science, arts and humanities) before the remaining articles were assessed for inclusion based on the following criteria:

- the article must include (or be in its entirety) an ERA that applies either a qualitative, semi-quantitative, or quantitative WOE methodology (after Linkov *et al.* 2009);
- the assessment must make direct reference to the uncertainties to be recorded within this research, thereby minimising researcher-subjectivity when creating the typology;
- the assessment must be original research and not a review of previously published work, in order to avoid duplicate values; and
- an aspect of the environment must feature in at least one part of the source-pathway-receptor (S-P-R) paradigm, where the environment “... consists of all, or any, of the following media, namely the air, water, or land” (EPA 1990).

These criteria ensured that only original (i.e. non-review-based) environmentally-focused WOE-ERAs (including ecological, environmental, and human-health risk assessments) that specifically mentioned uncertainty were included within this study, whilst the general search terms used allowed representation from a wide range of research domains.

Data Collection

The articles (conforming to the selection criteria) were examined in full and relevant information was extracted and recorded in separate spreadsheet entries. A working list of definitions was kept to ensure that observations were consistent and distinctions between uncovered uncertainties were not blurred. Importantly, no upper limit was set as to the number of UMTs that could be associated with each identified uncertainty type.

Data Organisation

The uncertainty data were organised using an iterative category clustering technique (Hartigan 1975). The different objects (i.e. the uncertainties) were categorised into distinct groups, such that the degree of association between any two objects was maximal if they belonged to the same group and minimal otherwise. In this way, the articles (from the data collection stage) were organised into groups by relevance to other similar data values. To reduce the potential for subjectivity in assigning objects to groups, the process was performed iteratively, with definitions and categorisations continually refined.

Data Analysis

The frequencies with which the different locations and sub-locations of uncertainties were associated with the UMTs were recorded. These were converted to percentage values of total occurrences in order to identify the most commonly occurring relationships. A separate bivariate analysis was performed using SPSS v18 (SPSS Inc., Chicago IL) to quantify the relationships between all two-variable combinations ($P \leq 0.01$).

RESULTS

Data Frequencies and Organisation

Uncertainty typology

Analysis of the collected WOE-ERA literature (n=171 assessments), in conjunction with iterative clustering of the extracted data (Figure 1), revealed 20 separate types of uncertainty (Table 2), with a total of 385 individual occurrences. The data uncertainty (n=125 out of 385; 32.5%) and extrapolation uncertainty (n=110; 28.6%) locations were the most

frequently occurring, with the decision uncertainty (n=6; 1.6%) and language uncertainty (n=16; 4.2%) categories the least frequent.

[FIGURE 1 NEAR HERE] [TABLE 2 NEAR HERE]

Uncertainty management techniques

Data extracted from the sources highlighted the use of a variety of UMTs (n=27), with a total of 453 separate applications. Occurrence proportions of the most frequently occurring mechanisms are shown in Table 3, along with brief descriptions and associated uncertainties. Monte-Carlo simulation was adopted most frequently (n=100 out of 453; 22.1%), followed by uncertainty factors (n=75; 16.6%), sensitivity analysis (n=38; 8.4%), and 'taking no action' (n=35; 7.7%).

[TABLE 3 NEAR HERE]

Relationships Between Uncertainties and Uncertainty Management Techniques

Frequency relationships

The highest frequency relationships between the uncertainty locations and UMTs employed (Figure 2) occurred between data uncertainties and Monte-Carlo simulation (n=56 out of 453 relationships), between extrapolation uncertainties and uncertainty factors (n=40), and between extrapolation uncertainties and Monte-Carlo simulation (n=18). On a proportional basis, the highest dependencies were seen between language uncertainties and fuzzy logic (68.8%; i.e. language uncertainties were managed with fuzzy logic in 68.8% of cases), model uncertainties and sensitivity analysis (35.1%), and data uncertainties and Monte-Carlo simulation (34.4%). Overall, uncertainties were associated with at least one UMT in 92.3% of cases, and were therefore unmanaged 7.7% of the time.

[FIGURE 2 NEAR HERE]

Statistical relationships

The strongest correlations between the uncertainty types and UMTs (Figure 3) occurred between decision uncertainty and adaptive management ($\rho=0.57$), spatial (extrapolation) uncertainty and interpolation ($\rho=0.46$), and cause (system) uncertainty and causal influence ($\rho=0.40$). A similar strength correlation occurred between the portion of data uncertainties used as parameter values in computational and/or numerical models (and therefore consist of repeated values from within the data location; marked *model input* in Figure 3) and Monte-Carlo simulation ($\rho=0.32$).

Positive correlations were also observed between several uncertainty-location/UMT combinations, where all individual uncertainty types within the location shared a positive correlation with the respective UMT. The strongest of these relationships were language uncertainties with fuzzy logic ($\rho=0.45$) and fuzzy-stochastic systems ($\rho=0.24$), and model uncertainty with sensitivity analysis ($\rho=0.29$).

[FIGURE 3 NEAR HERE]

DISCUSSION

An Improved (Evidence-Based) Uncertainty Typology

The existing uncertainty typologies (Table 1) are predominantly based within specific research areas, using categorisations that are primarily relevant to those fields. They communicate varying frequencies of uncertainties, often in a contradictory fashion, and use a number of different approaches in their construction, including small-scale literature reviews (e.g. Regan *et al.* 2002) and amalgamations of existing frameworks (e.g. Ascough II *et al.* 2008). This has led to overlapping and contradictory sets of categorisations. The uncertainty typology presented in Table 2 addresses the following issues:

- The set of articles analysed included ecological, environmental, and human-health risk assessments. Although the specific requirements of these assessments differ, they do contain the same four phases and many of the same processes (US EPA 1998; Zhang et al. 2010; DEFRA 2011). Therefore, the developed typology does not restrict observations to narrowly-defined research domains (e.g. conservation biology) but instead extends the focus to all concerns of an environmental nature, enabling the typology to be more transferrable and relevant to a larger number of risk analysts.
- Using WOE-ERAs, which contain a variety of ERA techniques as well as distinct forms of evidence, increases the potential for a larger spectrum of uncertainties to exist. This is reflected in the typology which, containing 20 distinct forms of location-based uncertainties arranged according to their natures, is the most extensive to date.
- By constructing and interrogating a large supporting evidence base of peer-reviewed articles (n=171) all uncertainty categorisations within the typology are supported and defensible.

It is also pertinent to address the potential limitations associated with the method used to construct the typology and its resulting categorisations:

- *Dependence on existing assessments to contain reliable information.* This limitation may have been realised where incorrect information was presented within the sourced materials, though the peer-review process was expected to resolve these errors. Perhaps of more concern was the potential omission (rather than incorrect inclusion) of important uncertainties; key uncertainties that went unidentified in the source materials could not feature in the typology. However, the evidence base of 171 assessments was believed to be extensive enough to account for all potential uncertainties.

- *Subjectivity in the information clustering process.* The clustering process used to form categorisations within and between the different types of uncertainty, whilst efficient and effective, did require an element of subjectivity on the part of the researcher. This type of qualitative clustering has the potential to blur definitions, thereby reducing the clarity of the clustered output. This potential limitation was managed as far as possible by making the clustering process transparent (see Figure 1).
- *Representativeness of the typology for application to ERAs.* Limiting the included studies to WOE-ERAs may have led to biases within the evidence base, which would have been transferred into the typology. One potential bias was a focus on risk domains in which WOE-ERAs are commonly used. This potential limitation may result in a lack of representativeness when applying the typology to non-WOE-ERA scenarios. However, when weighed against other viable alternatives, such as building an evidence base of ERAs based in specific risk domains, the WOE-ERA approach was deemed to be the most representative for future application of the typology.

The outlined advantages together with the management of potential limitations ensures that the presented typology addresses the issues associated with existing categorisations.

Defining Uncertainty

The nature of uncertainty

Interrogation and analysis of the WOE-ERAs (n=171) identified a total of three types of nature-based uncertainty, which are discussed in detail here.

Aleatory uncertainty

Aleatory uncertainty represents the inherent randomness displayed in human and natural systems (Bedford and Cook 2001; Ascough II *et al.* 2008). Aleatory uncertainty

cannot be reduced, although additional research may help to better understand the complexities of the system(s) of interest. Whilst such systems may in actuality be chaotic rather than random (and are therefore in principle understandable; Regan *et al.* 2002), risk analysts find it useful to treat the associated uncertainties from the latter position. For example, stochastic numerical techniques (such as Monte-Carlo simulation and Latin Hypercube sampling) act as realistic representations of real-world processes, which are either viewed as being too complex for deterministic interpretation (e.g. seismic activity) or as inherently random (e.g. weather systems). However, in mimicking nature, stochastic models can produce results that are consistently more representative than their deterministic counterparts (Hromkovic 2005).

Epistemic uncertainty

Epistemic uncertainty (Bedford and Cooke 2001; Walker *et al.* 2003; Petersen 2006; Ascough II *et al.* 2008; Knol *et al.* 2009) represents the imperfection of knowledge concerning a system of interest. Epistemic uncertainty can be quantified, reduced, and possibly eliminated, depending on the specific situation. However, whilst epistemic uncertainty is in principle reducible by increasing relevant knowledge, this new information can reveal the true depths of our ignorance, only serving to increase the associated uncertainty (Janssen *et al.* 2003; van der Keur 2008).

Combined uncertainty

This research introduces a combined epistemic and aleatory category, reflecting the potential for the location-based uncertainties contained within it to incorporate both epistemic and aleatory aspects, and forcing a separation from those sets. For example, model uncertainty may incorporate system uncertainty, which can reduce confidence in the structure

of a model, as well as variability uncertainty, which may cast doubt over the validity of the model's output. For this combined category, reducing secondary uncertainties associated with incorporated groups is just as important as managing the primary failings.

The location of uncertainty

Interrogation and analysis of the WOE-ERAs (n=171) identified a total of 7 main types of location-based uncertainty and 20 related sub-types, which are discussed in detail here.

Data uncertainty

Data are used extensively in risk assessments, not least environmental WOE-ERAs. For example, data may be used to draw attention to a source of environmental danger, to assess the degree of harm imposed upon a valued asset, or to support or refute damaging claims made against an individual, organisation, or even nation. Whether empirical or experimental, all data carries a level of inherent confidence associated with its truth and correctness. Identifying potential sources of uncertainty within data can help to distinguish between the reliable and the unreliable.

Data uncertainties can be further arranged into three groups: *availability*, referring to the incompleteness, scarcity, or absence of data (i.e. data is not available); *precision*, concerning the lack of accuracy in obtained data (i.e. data is not precise); and *reliability*, reflecting its trustworthiness (i.e. data is unreliable, possibly due to errors associated with its processing, statistical analysis, or presentation). The data reliability sub-location, which accounts for 20.8% of all uncertainties within the WOE-ERA evidence base, primarily reflects the measurement and systematic sub-categories seen within existing typologies (Table 1).

Language uncertainty

Language is used both in conjunction with and separately to data. The uncertainties associated with language arise for a number of reasons, but stem primarily from a lack of clarity. Language can be used to express ideas and commands or to communicate the final results of assessments; its use is unavoidable and necessary.

Linguistic uncertainties are comprised of three types: *ambiguity*, where multiple meanings are possible; *underspecificity*, where meanings are not exact; and *vagueness*, where meanings are not clear and understandable. The use of a single field-specific term can carry all three linguistic uncertainties: it may not be clearly defined and therefore have many meanings throughout the community (ambiguous); its use may be superseded by a more relevant and accurate term (underspecific), and certain members, especially those from outside the field, may have heard of the term, but have a limited understanding of its true meaning (vague; Acosta *et al.* 2010).

In previous typologies, language uncertainties (if included at all) were typically separated into their own category (e.g. Morgan *et al.* 1990; Regan *et al.* 2002; Ascough II *et al.* 2008), but are here deemed to be epistemic. The uncertainties associated with language arise for a number of reasons, but stem primarily from a lack of clarity (Morgan *et al.* 1990). However, the definitions, contexts, and applications associated with language can be controlled (Regan *et al.* 2002). Theoretically, language uncertainties can be quantified, reduced or even removed – techniques such as fuzzy logic are testament to this – equating them with the other uncertainties (data and system) within the epistemic set. Despite their relatively low levels of occurrence within the WOE evidence base (of just 4.7%; Figure 4.2), communicating the epistemic quality of language uncertainties allows analysts to approach them with reduction and elimination in mind, which may previously not have been the case.

System uncertainty

System uncertainty tallies closely to scientific understanding; if the understanding is low the uncertainty will be high, and vice-versa. However, a field which develops rapidly, such as biotechnology, will contain high levels of knowledge as well as some system uncertainty, due largely to the unknowns that progress brings.

System uncertainties can be more clearly defined according to the source-pathway-receptor relationship, which constitutes the three main phases of system understanding: *cause*, which concerns a lack of clarity regarding the source(s) of harm; *effect*, relating to the influence a particular stressor (source) has upon the receptor(s); and *process*, which concerns either not understanding the risks or not identifying something vital to a successful assessment.

Process uncertainty correlates with the pathway stage of the relationship, which can be anything between the source(s) of harm and asset(s) of value. It can contain a variety of uncertainties, such as not identifying the critical dose needed for an adverse effect to result (Meek and Hughes 1995). The risks associated with certain nanotechnologies, a rapidly developing field, are unclear because of a lack of process understanding, which in some cases may be coupled with high effect uncertainty. For example, the contribution of physical structure to a nanoparticle's toxicity may not be fully understood (Gottschalk *et al.* 2010), whilst its effects upon different receptors may simply be unknown (Zalk *et al.* 2009).

Variability uncertainty

Also described as random and stochastic, variability uncertainty is the inherent unpredictability of any human or natural system. *Human* variability in ERAs results primarily from intentionally biased and subjective actions (Khan *et al.* 2002), but extends to all

qualities of humans which are, either literally or from the viewpoint of the risk analyst, stochastic in nature. Irrespective of their position or seniority, humans involved in the assessment process may display bias when they have something to gain, or subjectivity when they believe their own views to be more correct than those of others (Chen *et al.* 2007). Human variability can be exhibited by those with close links to a project, such as decision-makers, stakeholders, and scientists, as well as those with no vested interest, such as hired laboratory technicians or computer modellers (Croke *et al.* 2007).

The *natural* element may be considered unexpected and free from intentional bias (Jørgensen *et al.* 2009). It pertains to the chaotic traits of natural systems. Natural variability is also the primary cause of uncertainties associated with extrapolation; a process that becomes necessary when faced with limited knowledge (e.g. limited data or limited process understanding).

Extrapolation uncertainty

Extrapolation can occur across a variety of means, and is usually present wherever there is missing information or knowledge (Luttik *et al.* 2005), but is not necessarily associated with numeric data. In the developed typology, extrapolation uncertainty is a sub-category of the aleatory category, where previously it has either been grouped with model uncertainties (Walker *et al.* 2003; Regan *et al.* 2002; Finkel 1990), treated as a branch of variability (Huijbregts 2001), or more commonly ignored altogether. Extrapolation can be considered an attempt at rectifying availability issues: if information were readily available, extrapolation would not be necessary. However, when it is required, the process is deemed uncertain due to the natural variability involved (e.g. spatially and temporally extrapolating meteorological data beyond the physical limits of an existing network of measuring stations to a study site). Extrapolation can therefore be considered the result of epistemic failings,

with the connected uncertainties driven through aleatory means. Whilst an increase in relevant epistemic knowledge may prevent the need for extrapolation (thereby providing a distinction from variability uncertainty, which can be neither eliminated nor reduced), when it is required it is the aleatory-based failings that must be addressed. These observations confirm extrapolation uncertainties to be aleatory in nature, and indicate that they should be considered separately from the variability location.

Extrapolation is identified in six forms: *intraspecies*, where information specific to members of a species is used to represent other members of the same species; *interspecies*, where information specific to members of a species is used to represent members of a different species; *laboratory*, where information specific to laboratory conditions is used to represent real-world scenarios; *quantity*, where information specific to one quantity is used to represent another; *spatial*, where information specific to one spatial scale is used to represent another; and *temporal*, where information specific to one timescale is used to represent another.

Model uncertainty

With regard to a system of interest, modelling is an attempt to understand processes, predict responses, evaluate management alternatives, and support the policy and decision-making process (Arhonditsis *et al.* 2007). Modelling procedures vary according to the system of study and desired outcomes, though they invariably involve an initial conceptualisation stage, which is then developed into a numerical and/or computational representation (Stephens *et al.* 1993). Simplifications and assumptions are usually necessary features of the structural process, since natural features and dependencies are complex and numerous. The initial conceptualisation stage is arguably the most important. Any uncertainties that exist here will likely be propagated throughout the rest of the modelling procedure. The conceptual

representation also needs to be fit for purpose: an oversimplification may result in a failure to capture essential features, leading in turn to inadequate numerical or computational simulations. Conversely, an undersimplification may yield a model that is too complex, and therefore time-intensive, or even prohibitive, to build and execute (El-Ghonemy *et al.* 2005).

Model uncertainties relate to the different stages of the process: *structure*, which concerns the representation of real-world processes in model form; and *output*, which reflects the level of confidence in the results. The model structure sub-location, which accounts for 3.9% of all uncertainties within the WOE-ERA evidence base, primarily reflects the structural and technical sub-categories seen within existing typologies (Table 3.1).

Decision uncertainty

Decision uncertainty exists when doubt surrounds an optimal course of action, often in the face of differing objectives. There may be multiple options which satisfy at least a part of the criteria for the decision, but also possible is the existence of no such alternatives. For example, management of ecological and environmental resources requires decision-makers to evaluate multiple and often conflicting strategies, whilst balancing objectives of productivity and sustainability (Ducey and Larson 1999). Decision uncertainty is potentially comprised of all uncertainties identified up to and including this stage of the WOE-ERA process.

The level of uncertainty

Every identified uncertainty with a defined nature- and location-type must also be considered in terms of its level (i.e. severity; Janssen *et al.* 2003; Walker *et al.* 2003; Refsgaard *et al.* 2007). The level of an identified uncertainty is highly context-dependant and cannot, at present, be ascribed *a priori* along with its nature and location. Due to this, there is a reduced need (compared with the nature and location) for an uncertainty typology to make

specific reference to potential levels within its main structure. It may simply be more appropriate to do it in an accompanying narrative, as is the case here.

Humans exhibit a variety of distinct levels of knowledge, ranging from determinism (perfect knowledge) to indeterminacy (lack of knowledge; Wynne 1992). The further we move from a deterministic understanding of a system, the more severe the uncertainty becomes (Walker *et al.* 2003). The level of uncertainty is described according to two factors, namely the degree of confidence attached to the likelihood of an event occurring, and the degree of confidence attached to the severity of outcomes should that event occur (Wynne 1992; Stirling 1999). These metrics are used to convey the level of understanding, and therefore the level of the associated uncertainty. Recognised levels of uncertainty include: *deterministic uncertainty*, in which we are confident about the likelihoods and outcomes; *statistical uncertainty*, where we can confidently assign probabilities to events but have little understanding of the ramifications of the events; *scenario uncertainty*, where there is confidence about the outcomes but not likelihoods of an event (i.e. the reverse of statistical uncertainty); *recognised ignorance*, where it is not possible to define probabilities or a complete set of outcomes; and *total ignorance*, which is the uncertainty of which we know nothing and to which we are ignorant (i.e. the inverse of deterministic uncertainty).

When the focus shifts from uncertainty identification (i.e. the purpose of the typology presented here) to uncertainty management, an effective typology should also aim to communicate methods for quantification and/or reduction. In that instance, communicating the uncertainty levels is essential as a change in level can cause a change in the optimal UMT. In terms of data uncertainties, for example, when there is a level of statistical uncertainty the associated data uncertainty can be tackled through sensitivity analysis. However, if we were in the range of scenario uncertainty, scenario analysis, for example,

would be more appropriate (Refsgaard *et al.* 2007). Ultimately, selection of a suitable UMT is dependent on the mix of all three uncertainty dimensions: location, nature and level.

Dealing with Uncertainty

The appropriateness of UMTs employed

UMTs should be used in concert with specific types of uncertainty (Refsgaard *et al.* 2007). The correct adoption of any one UMT is therefore dependent upon the uncertainties present (Stirling 2012). The occurrence frequency analysis and statistical analysis conducted between the uncertainty types and UMTs highlighted several relationships, the vast majority of which show the UMTs being used to tackle appropriate uncertainties. This observation extends to frequently occurring uncertainty and UMT combinations (e.g. Monte-Carlo simulation being used to tackle data reliability uncertainty; Figure 3) as well as those combinations which occur less frequently, but are no less appropriate (e.g. MCDA being used to tackle decision uncertainty; Figure 3; Linkov and Moberg 2011). This is a positive finding, since the incorrect utilisation of a UMT may be considered just as important as choosing not to use one at all, which was the fourth most-adopted option in the studied data set. We have defined taking ‘no action’ as the publication author(s) recognising uncertainties but not taking action, with or without offering justification (e.g. Wright-Walters *et al.* 2011). As well as indicating the inappropriate use of this technique with reference to specific uncertainties (primarily model and variability), the occurrence frequency analysis and resulting dependency model (Figure 2) convey a more important point: dealing with uncertainties should be a major priority within these assessments. The fact that the ‘no action’ mechanism appears so often suggests that this is not currently the case.

Separating uncertainty and variability

The categorisation of uncertainties as being either epistemic, aleatory, or a combination of the two, might imply that each of the identified UMTs can equally be assigned to one of these groups. This is not the case, nor is there a single mechanism that offers comprehensive solutions to all of the identified uncertainties.

Whilst uncertainties appear to fall easily into the aforementioned groupings, the boundary can be less well defined in applied situations (Merz and Thieken 2009). The most pertinent example of this is the use of Monte-Carlo Simulation in an attempt to cope with both forms of uncertainty. Since epistemic and aleatory uncertainties can both be described by probability distributions, many assessments involving a first-order Monte-Carlo procedure claim to successfully handle both (Wu and Tsang 2004). However, the ensuing single distribution (which may combine data reliability uncertainty with inherent natural variability) incorrectly implies that uncertainty and variability are the same, and that they can be dealt with as one (Wu and Tsang 2004). Problems may still exist even when a distinction is made: incorrectly treating variability as if it were uncertainty may yield a meaningless distribution when a single figure is required (Vose 2000). Effectively, the techniques that are employed to manage uncertainty can, if executed incorrectly, introduce further errors.

It is increasingly recognised that uncertainty and variability need to be treated separately (Kelly and Campbell 2000; Li *et al.* 2008; Kumar *et al.* 2009; Qin and Huang 2009; Helton *et al.* 2011). Once separated, both aleatory variability and epistemic uncertainty can be quantified, and steps can be taken to reduce and potentially remove epistemic uncertainty. Techniques such as second-order Monte-Carlo (Griffin *et al.* 1999; Wu and Tsang 2004) and integrated fuzzy-stochastic systems (Li *et al.* 2007; Kumar *et al.* 2009; Qin and Huang 2009) have emerged that can manage both aleatory and epistemic uncertainties. Moreover, through correct uncertainty management, they attempt to eliminate the inferred,

and potentially unjustifiable, level of confidence that can incorrectly be assigned to risk estimates.

Guidance for practitioners

In order to help practitioners better prioritise, identify, and manage uncertainties in assessments, we propose combining the uncertainty typology (Table 2) with the uncertainty-based frequency and dependency (Figure 2) data. The resulting list of potential uncertainties (Table 4), which is organised by uncertainty location and sub-location, is ranked according to the frequency with which the uncertainties appear in the evidence base (of 171 WOE-ERAs). These rankings correspond to the order in which practitioners may wish to consider uncertainties in their assessments. The individual uncertainties are further categorised according to their nature. In addition, several options for managing each uncertainty are presented, ordered according to the strength of the dependencies between an uncertainty sub-location and its respective UMTs within the evidence base (where one is the optimal UMT and three is the least optimal).

[TABLE 4 NEAR HERE]

Applying the guidance: the case of genetically modified higher plants

In the European Union (EU) the introduction of Genetically Modified Organisms (GMOs) for experimental purposes and for placing on the market for cultivation, importation or processing is regulated by European Commission Directive 2001/18/EC (EC 2001). In order to obtain consent for purposes of deliberate release into the environment, applicants must submit a comprehensive dossier containing relevant information about the GMO, including an ERA. However, submitted ERAs rarely consider uncertainties, and where uncertainty is acknowledged it is primarily handled by adopting (favourable) worst-case estimates (Hart *et al.* 2007).

A well-researched example of GMOs in the environment is the potential risk of *Bacillus thuringiensis* (Bt) modified maize to non-target Monarch butterflies, with research in USA investigating levels of risk under differing exposure scenarios. For the purposes of this research, this specific case can reasonably be expanded to a more generalised relationship of potential Genetically Modified Higher Plant (GMHP) risk to Lepidoptera. Whilst 81 examples of this scenario exist within the publically available dossiers submitted by applicants under Directive 2001/18/EC, the dossiers do not include evidence to support attempts to identify or manage uncertainties within their respective ERAs, which seems to contradict the instruction in the enforcing regulation to do so.

Directive 2001/18/EC promotes a six-step ERA procedure for applicants to follow, where the first four steps correspond to the ERA and the final two to risk management options beyond the assessment. The first four steps are commonly known as problem formulation, effects assessment, exposure assessment, and risk characterisation (DEFRA 2011). An ERA carried out by an applicant can be expected to consist of these four phases, which, on the basis of information contained within relevant governmental guidance documents (Fairman *et al.* 1998; USEPA 1998; DEFRA 2011), and in the context of potential GMHP risk to Lepidoptera, could contain most or all of the major elements listed in Table 5. The presented uncertainty typology (Table 2) and guidance (Table 4) can be applied to this standard ERA structure to determine potential locations of uncertainty and relevant options for their management (Table 6). For example, problems may exist when attempting to determine aspects of the dose of the GMHP stressor (e.g. modified protein) received by the Lepidoptera receptor during the effects assessment phase of the ERA. Such issues could feasibly correspond to: uncertainty in applying relevant data about the duration, frequency, or intensity of the dose (leading to data reliability, availability, and/or precision uncertainty); variability about the situation (natural variability); forced extrapolations from the available

data to other points of interest (interspecies, spatial, intraspecies, temporal, laboratory, and/or quantity extrapolation uncertainty), or; defining parameters in models that reflect the data utilised (model structure uncertainty) and using those models to quantify the dose received (model output uncertainty).

[TABLE 5 NEAR HERE] [TABLE 6 NEAR HERE]

By applying the uncertainty typology (Table 2) and guidance (Table 4) to the rest of the ERA structure in Table 5, we formulated a list of 43 potential uncertainties. These uncertainties are categorised according to the four phases of the ERA and the main locations in which uncertainty can exist (e.g. data, variability), which are in turn organised in order of their highest ranked uncertainty sub-location (e.g. data reliability, natural variability). System uncertainties are likely to dominate the problem formulation phase, with data, variability, extrapolation, and model uncertainties the focus in the middle analysis phase (effects and exposure assessments), and language and decision uncertainties playing more of a role at the final risk characterisation step. The responsibility for determining whether the potential uncertainties exist, and at what level of severity, will rest with the relevant applicant(s). Prioritised techniques for the management of each uncertainty sub-location (brought forward from Table 4) are also included. When implementing these UMTs applicants should ensure that epistemic and aleatory uncertainties are approached in the correct way.

This simple example demonstrates how potential uncertainties can be identified using the presented uncertainty typology and guidance. This may allow for more considered uncertainty analyses in both established risk domains and highly regulated emerging fields, such as GMHPs, leading to more robust ERAs. Environmental decision-making at some of the highest strategic levels (e.g. the European Union) may ultimately benefit. However, the researchers recognise that application of the presented typology will inevitably require some end-user subjectivity, and that consistent reproduction of results may be hard to achieve. To

that end, the researchers are currently investigating, applying and validating methods to improve the uncertainty identification process within ERAs, which build on the presented typology and reduce the reliance on the skill, experience and ability of the end-user.

CONCLUSION

Uncertainty typologies aim to foster understanding, further acting as tools to aid uncertainty identification during risk characterisation. The categorisations and definitions presented within uncertainty typologies must be comprehensive and reliable, but existing typologies have been found to be lacking in a number of ways, especially in an ERA context.

This research presented a typology of uncertainties based, for the first time, on the analysis of a large evidence base, namely 171 peer-reviewed environmental WOE-ERAs. In creating the typology, which consists of 7 main types of location-based uncertainty (data, language, system, extrapolation, variability, model, and decision) and 20 related sub-types, several key issues surrounding existing typologies, including research domain transferability and content reliability issues, have been resolved. In addition, whilst the techniques used by analysts to manage these uncertainties were implemented appropriately, we have shown that in some cases the validity of a risk estimate is negatively impacted as uncertainty management is excluded. The practical guidance that we have introduced here will help resolve this issue by providing a robust method for dealing with uncertainty, as demonstrated through an applied case study focussing on ERAs of genetically modified higher plants in the EU. This case study also highlights the relationships between different uncertainties and the various phases and tasks within ERAs. Moving forward, we are currently exploring these relationships in more detail, with the aim of adding value to the uncertainty identification process.

The typology presented here and accompanying guidance, which should be utilised by risk analysts during the formative stages of uncertainty analyses, will have positive implications for the identification, prioritisation, and management of uncertainty during risk characterisation.

ACKNOWLEDGEMENTS

The Collaborative Centre of Excellence in Understanding and Managing Natural and Environmental Risks is funded jointly by EPSRC/ESRC/NERC/Defra under grant EP/G022682/1. DJCS is funded by Cranfield University through an EPSRC Doctoral Training Account.

REFERENCES

- Acosta H, Wu D, and Forrest BM. 2010. Fuzzy experts on recreational vessels, a risk modelling approach for marine invasions. *Ecol Model* 2215:850-863
- Agüero A, Pinedo P., Simón I, *et al.* 2008. Application of the Spanish methodological approach for biosphere assessment to a generic high-level waste disposal site. *Sci Total Environ* 403(1-3): 34-58
- Ahlers J, Stock F, and Werschkun B. 2008. Integrated testing and intelligent assessment - New challenges under REACH. *Environ Sci Pollut R* 15(7): 565-572
- Alcamo J and Bartnicki J. 1987. A framework for error analysis of a long-range transport model with emphasis on parameter uncertainty. *Atmos Environ*, 21:2121-2131
- Alden III RW, Hall Jr. LW, Dauer DM, *et al.* 2005. An integrated case study for evaluating the impacts of an oil refinery effluent on aquatic biota in the Delaware River: Integration and analysis of study components. *Hum Ecol Risk Assess* 11(4): 879-936
- Alvarez-Guerra M, Viguri J, and Voulvoulis N. 2009. A multicriteria-based methodology for site prioritisation in sediment management. *Environ Int* 35(6): 920-930
- An YJ, Yoon CG, Jung KW, *et al.* 2007. Estimating the microbial risk of E. coli in reclaimed wastewater irrigation on paddy field. *Environ Monit Assess* 129(1-3): 53-60
- Apitz SE, Barnati A, Bernstein AG, *et al.* 2007. The assessment of sediment screening risk in Venice Lagoon and other coastal areas using international sediment quality guidelines. *J Soil Sediment* 7(5): 326-341
- ApSimon HM, Warren RF, and Kayin S. 2002. Addressing uncertainty in environmental modelling: A case study of integrated assessment of strategies to combat long-range transboundary air pollution. *Atmos Environ* 36(35): 5417-5426

668 Arhonditsis G, Qian S, Stow C, *et al.* 2007. Eutrophication risk assessment using Bayesian
669 calibration of process-based models: Application to a mesotrophic lake. *Ecol model*
670 208(2-4):215-229

671 Ascough II JC, Maier HR, Ravalico JK, *et al.* 2008. Future research challenges for
672 incorporation of uncertainty in environmental and ecological decision-making. *Ecol*
673 *Model* 2193-4:383-399

674 Aspinall W, Woo G, Voight B, *et al.* 2003. Evidence-based volcanology: application to
675 eruption crises. *J Volcanol Geoth Res* 128:273-285

676 van Asselt M and Rotmans J. 2002. Uncertainty in integrated assessment modelling. *Climatic*
677 *Change* 54:75-105

678 Avagliano S and Parrella L. 2009. Managing uncertainty in risk-based corrective action
679 design: Global sensitivity analysis of contaminant fate and exposure models used in the
680 dose assessment. *Environ Model Assess* 141:47-57

681 Avagliano S, Vecchio A, and Belgiorno V. 2005. Sensitive parameters in predicting exposure
682 contaminants concentration in a risk assessment process. *Environ Monit Assess* 111(1-
683 3): 133-148

684 Babendreier JE and Castleton KJ. 2005. Investigating uncertainty and sensitivity in
685 integrated, multimedia environmental models: Tools for FRAMES-3MRA. *Environ*
686 *Modell Softw* 20(8): 1043-1055

687 Baccou J, Chojnacki E, Mercat-Rommens C, *et al.* 2008. Extending Monte Carlo simulations
688 to represent and propagate uncertainties in presence of incomplete knowledge:
689 Application to the transfer of a radionuclide in the environment. *J Environ Eng* 134(5):
690 362-368

691 Barron MG, Duvall SE, and Barron KJ. 2004. Retrospective and current risks of mercury to
692 panthers in the Florida everglades. *Ecotoxicology* 13(3): 223-229

693 Batley GE, Burton GA, Chapman PM, *et al.* 2002. Uncertainties in sediment quality weight-
694 of-evidence (WOE) assessments. *Hum Ecol Risk Assess* 8(7): 1517-1547

695 Batzias F and Siontorou CG. 2007. A novel system for environmental monitoring through a
696 cooperative/synergistic scheme between bioindicators and biosensors. *Environ Manage*
697 82(2): 221-239

698 Baudrit C, Guyonnet D, and Dubois D. 2007. Joint propagation of variability and imprecision
699 in assessing the risk of groundwater contamination. *J Contam Hydrol* 93(1-4): 72-84

700 Beck M. 1987. Water quality modeling: a review of the analysis of uncertainty. *Water Resour*
701 *Res* 23:1393-1442

702 Bedford T and Cooke RM. 2001. *Probabilistic Risk Analysis: Foundations and Methods*.
703 Cambridge University Press, Cambridge, UK

704 Beer T. 2006. Ecological risk assessment and quantitative consequence analysis. *Hum Ecol*
705 *Risk Assess* 12:51-65

706 Benekos ID, Shoemaker CA, and Stedinger JR. 2007. Probabilistic risk and uncertainty
707 analysis for bioremediation of four chlorinated ethenes in groundwater. *Stoch Env Res*
708 *Risk A* 21(4): 375-390

709 Benke KK and Hamilton AJ. 2008. Quantitative microbial risk assessment: Uncertainty and
710 measures of central tendency for skewed distributions. *Stoch Env Res Risk A* 22(4):
711 533-539

712 Bennett JR, Kaufman CA, Koch I, *et al.* 2007. Ecological risk assessment of lead
713 contamination at rifle and pistol ranges using techniques to account for site
714 characteristics. *Sci Total Environ* 374(1): 91-101

715 Bevington P and Robinson D. 2002. *Data reduction and error analysis for the physical*
716 *sciences, Vol. 3*. McGraw-Hill, New York

717 Beyer C, Altfelder S, Duijnisveld WHM, *et al.* 2009. Modelling spatial variability and
 718 uncertainty of cadmium leaching to groundwater in an urban region. *J Hydrol* 369(3-4):
 719 274-283

720 Bittueva MM, Abilev SK, and Tarasov VA. 2007. Efficiency of the prediction of
 721 carcinogenic activities of chemical substances based on scoring somatic mutations in
 722 the soybean Glycine max (L.) Merrill. *Russ J Genet* 43(1): 64-72

723 Blazkova S and Beven K. 2004. Flood frequency estimation by continuous simulation of
 724 subcatchment rainfalls and discharges with the aim of improving dam safety assessment
 725 in a large basin in the Czech Republic. *J Hydrol* 292(1-4): 153-172

726 Borsuk ME, Reichert P, Peter A, *et al.* 2006. Assessing the decline of brown trout (*Salmo*
 727 *trutta*) in Swiss rivers using a Bayesian probability network. *Ecol Modell* 192(1-2):
 728 224-244

729 Bosgra S, van der Voet H, Boon PE, *et al.* 2009. An integrated probabilistic framework for
 730 cumulative risk assessment of common mechanism chemicals in food: An example
 731 with organophosphorus pesticides. *Regul Toxicol Pharm* 54(2): 124-133

732 Bosgra S, Bos PMJ, Vermeire TG, *et al.* 2005. Probabilistic risk characterization: An
 733 example with di(2-ethylhexyl) phthalate. *Regul Toxicol Pharm* 43(1): 104-113

734 Brèchignac F and Doi M. 2009. Challenging the current strategy of radiological protection of
 735 the environment: arguments for an ecosystem approach. *J Environ Radioact* 100(12):
 736 1125-1134

737 van den Brink C, Zaadnoordijk WJ, Burgers S, *et al.* 2008. Stochastic uncertainties and
 738 sensitivities of a regional-scale transport model of nitrate in groundwater. *J Hydrol*
 739 361(3-4): 309-318

740 Brouwer R and De Blois C. 2008. Integrated modelling of risk and uncertainty underlying the
 741 cost and effectiveness of water quality measures. *Environ Modell Softw* 23:922-937

742 Brown J. 2004. Knowledge, uncertainty and physical geography: towards the development of
743 methodologies for questioning belief. *T I Brit Geogr* 29:367-381

744 Buekers J, Steen Redeker E, and Smolders E. 2009. Lead toxicity to wildlife: Derivation of a
745 critical blood concentration for wildlife monitoring based on literature data. *Sci Total*
746 *Environ* 407(11): 3431-3438

747 Burgman MA, Keith DA, and Walshe TV. 1999. Uncertainty in comparative Risk Anal for
748 threatened Australian plant species. *Risk Anal* 19(4): 585-598

749 Burton Jr. GA, Greenberg MS, Rowland CD, *et al.* 2005. In situ exposures using caged
750 organisms: A multi-compartment approach to detect aquatic toxicity and
751 bioaccumulation. *Environ Pollut* 134(1): 133-144

752 Calabrese EJ, Baldwin LA, Kostecki PT, *et al.* 1997. A toxicologically based weight-of-
753 evidence methodology for the relative ranking of chemicals of endocrine disruption
754 potential. *Regul Toxicol Pharmacol* 261 I:36-40

755 Caley P, Lonsdale WM, and Pheloung PC. 2006. Quantifying uncertainty in predictions of
756 invasiveness. *Biol Invasions* 8(2): 277-286

757 Campbell JE and Longsine DE. 1990. Application of generic risk assessment software to
758 radioactive waste disposal. *Reliab Eng Sys Safe* 30(1-3): 183-193

759 Cañellas-Boltà S, Strand R, and Killie B. 2005. Management of environmental uncertainty in
760 maintenance dredging of polluted harbours in Norway. *Water Sci Technol* 52(6): 93-98

761 Carlon C, Pizzol L, Critto A, *et al.* 2008. A spatial risk assessment methodology to support
762 the remediation of contaminated land. *Environ Int* 34(3): 397-411

763 Carrington CD, Cramer GM, and Bolger PM. 1997. A risk assessment for methylmercury in
764 tuna. *Water, Air, Soil Pollut* 97(3-4): 273-283

765 Cesar A, Marin A, Marin-Guirao L, *et al.* 2009. *Ecotox Environ Safe* 72:1832-1841

766 Chapman PM. 2007. Determining when contamination is pollution - Weight of evidence
767 determinations for sediments and effluents. *Environ Int* 334:492-501

768 Chen YC and Ma HW. 2007. Combining the cost of reducing uncertainty with the selection
769 of risk assessment models for remediation decision of site contamination. *Journal of J*
770 *Hazard Mater* 141(1): 17-26

771 Chen C, Ma H, and Reckhow K. 2007. Assessment of water quality management with a
772 systematic qualitative uncertainty analysis. *Sci total environ* 374(1):13-25

773 Chowdhury S, Champagne P, and McLellan PJ. 2009. Uncertainty characterization
774 approaches for risk assessment of DBPs in drinking water: A review. *J Environ*
775 *Manage* 90(5): 1680-1691

776 Chowdhury R and Flentje P. 2003. Role of slope reliability analysis in landslide risk
777 management. *Bul Eng Geol Environ* 62(1): 41-46

778 Collins JF, Alexeeff GV, Lewis DC, *et al.* 2004. Development of acute inhalation reference
779 exposure levels (RELs) to protect the public from predictable excursions of airborne
780 toxicants. *J Appl Toxicol* 24(2): 155-166

781 Collins G, Kremer JN, and Valiela I. 2000. Assessing uncertainty in estimates of nitrogen
782 loading to estuaries for research, planning, and risk assessment. *Environ Manage* 25(6):
783 635-645

784 Cothorn CR, Coniglio WA, and Marcus WL. 1986. Development of quantitative estimates of
785 uncertainty in environmental risk assessments when the scientific data base is
786 inadequate. *Environ Int* 12(6): 643-647

787 Crane JL and MacDonald DD. 2003. Applications of Numerical Sediment Quality Targets for
788 Assessing Sediment Quality Conditions in a US Great Lakes Area of Concern. *Environ*
789 *Manage* 32(1): 128-140

- Critto A, Torresan S, Semenzin E, *et al.* 2007. Development of a site-specific ecological risk assessment for contaminated sites: Part I. A multi-criteria based system for the selection of ecotoxicological tests and ecological observations. *ScTEen* 3791:16-33
- Croke B, Ticehurst T, Letcher R, *et al.* 2007. Integrated assessment of water resources: Australian experiences. *Water resour manag* 21(1):351-373
- Culp JM, Podemski CL, Cash KJ, *et al.* 2000. A research strategy for using stream microcosms in ecotoxicology: Integrating experiments at different levels of biological organization with field data. *J Aquat Ecosyst Stress Recovery* 7(2): 167-176
- Cupit M, Larsson O, De Meeûs C, G *et al* n. 2002. Assessment and management of risks arising from exposure to cadmium in fertilisers - II. *Sci Total Environ* 291(1-3): 189-206
- Daniels JI, Bogen KT, and Hall LC. 2000. Analysis of uncertainty and variability in exposure to characterize risk: Case study involving trichloroethylene groundwater contamination at Beale Air Force Base in California. *Water, Air, Soil Pollut* 123(1-4): 273-298
- Department for Environment, Food and Rural Affairs. 2011. *Guidelines for environmental risk assessment and management: green leaves III*. HMSO, London
- DelValls TA, and Riba I. 2007. A weight of evidence approach to assess sediment quality in the Guadalquivir estuary. *Aquat Ecosyst Health* 10(1): 101-106
- Dey WP, Jinks SM, and Lauer GJ. 2000. The 316(b) assessment process: Evolution towards a risk-based approach. *Environ Sci Policy* 3SUPPL.
- Dewulf A, Craps M, Bouwen R, *et al.* 2005. Integrated management of natural resources: dealing with ambiguous issues, multiple actors and diverging frames. *Water sci technol* 52:115-124

- Diodato N and Ceccarelli M. 2005. Environinformatics in ecological risk assessment of agroecosystems pollutant leaching. *Stoch Env Res Risk A* 19(4): 292-300
- Ducey M and Larson B. 1999. A fuzzy set approach to the problem of sustainability. *Forest Ecol Manag* 115(1):29-40
- Dunham JB, Young MK, Gresswell RE, *et al.* 2003. Effects of fire on fish populations: Landscape perspectives on persistence of native fishes and nonnative fish invasions. *Forest Ecol Manag* 178(1-2): 183-196
- Dussault ÉB, Balakrishnan VK, Sverko E, *et al.* 2008. Toxicity of human pharmaceuticals and personal care products to benthic invertebrates. *Environ Toxicol Chem* 27(2): 425-432
- Echevarria G, Sheppard MI, and Morel J. 2001. Effect of pH on the sorption of uranium in soils. *J Environ Radioact* 53(2): 257-264
- Efroymson RA, Jones DS, and Gold AJ. 2007. An ecological risk assessment framework for effects of onsite wastewater treatment systems and other localized sources of nutrients on aquatic ecosystems. *Hum Ecol Risk Assess* 13(3): 574-614
- El-Ghonemy H, Watts L, and Fowler L. 2005. Treatment of uncertainty and developing conceptual models for environmental risk assessments and radioactive waste disposal safety cases. *Environ int* 31(1):89-97
- Enick OV and Moore MM. 2007. Assessing the assessments: Pharmaceuticals in the environment. *Environ Impact Assess Rev* 27(8): 707-729
- Environmental Protection Act. 1990. *Environmental Protection Act: chapter 43, section 1.* HMSO, London
- European Commission Directive 2001/18/EC of the European Parliament and of the Council of 12 March 2001 on the deliberate release into the environment of genetically

modified organisms and repealing Council Directive 90/220/EEC. *Official Journal of the European Communities*, L106:1-39.

Fairman R, Mead C, Williams W, *et al.* 1998. *Environmental risk assessment: approaches, experiences and information sources*. Office for Official Publications of the European Communities

Faucheux S and Froger G. 1995. Decision-making under environmental uncertainty. *Ecol Econ* 15:29-42

Fewtrell L, Macgill SM, Kay D, *et al.* 2001. Uncertainties in risk assessment for the determination of drinking water pollutant concentrations: Cryptosporidium case study. *Water Res* 35(2): 441-447

Fiksel J. 1985. Quantitative Risk Anal for toxic chemicals in the environment. *J Hazard Mater* 10(2-3): 227-240

Filipsson M, Lindström M, Peltola P, *et al.* 2009. Exposure to contaminated sediments during recreational activities at a public bathing place. *J Hazard Mater* 171(1-3): 200-207

Finkel A. 1990. *Confronting uncertainty in risk management: a guide for decision-makers*. Center for Risk Management, Resources for the Future, Washington, DC.

Fischer DL 2005. Accounting for differing exposure patterns between laboratory tests and the field in the assessment of long-term risks of pesticides to terrestrial vertebrates. *Ecotoxicology* 14(8): 853-862

Fish R, Winter M, Oliver DM, *et al.* 2009. Unruly pathogens: eliciting values for environmental risk in the context of heterogeneous expert knowledge. *Environ Sci Policy* 12(3): 281-296

Forbes VE and Calow P. 2002. Applying weight-of-evidence in retrospective ecological risk assessment when quantitative data are limited. *Hum Ecol Risk Assess* 8(7): 1625-1639

862 Fuhrer J. 2009. Ozone risk for crops and pastures in present and future climates. *Naturwiss*
863 96(2): 173-194

864 Funtowicz SO and Ravetz JR. 1990. *Uncertainty and Quality in Science for Policy*. Kluwer,
865 Dordrecht

866 Godduhn A and Duffy LK. 2003. Multi-generation health risks of persistent organic pollution
867 in the far north: Use of the precautionary approach in the Stockholm Convention.
868 *Environ Sci Policy* 64:341-353

869 Golden RJ, Holm SE, Robinson DE, *et al.* 1997. Chloroform mode of action: Implications for
870 cancer risk assessment. *Regul Toxicol Pharm* 26(2): 142-155

871 Goodman J, McConnell E, Sipes I, *et al.* 2006. An updated weight of the evidence evaluation
872 of reproductive and developmental effects of low doses of bisphenol A. *Crit Rev*
873 *Toxicol* 36(5): 387-457

874 Gottschalk F, Sonderer T, Scholz RW, *et al.* 2010. Possibilities and limitations of modeling
875 environmental exposure to engineered nanomaterials by probabilistic material flow
876 analysis. *Environ Toxicol Chem* 29:1036-48

877 Greenberg MM. 1997. The central nervous system and exposure to toluene: A risk
878 characterization. *Environ Res* 72(1): 1-7

879 Griffin S, Goodrum PE, Diamond GL, *et al.* 1999. Application of a probabilistic risk
880 assessment methodology to a lead smelter site. *Hum Ecol Risk Assess* 5-4:845-868

881 Grist EPM, Crane M, Jones C, *et al.* 2003. Estimation of demographic toxicity through the
882 double bootstrap. *Water Res* 37(3): 618-626

883 Gurjar BR and Mohan M. 2003. Potential health risks due to toxic contamination in the
884 ambient environment of certain Indian states. *Environ Monit Assess* 82(2): 203-223

- Hacon S, Rochedo ERR, Campos RRR, *et al.* 1997. Mercury exposure through fish consumption in the urban area of Alta Floresta in the Amazon Basin. *J Geochem Explor* 58(2-3): 209-216
- Hamilton AJ, Stagnitti F, Premier R, *et al.* 2006. Quantitative microbial risk assessment models for consumption of raw vegetables irrigated with reclaimed water. *Appl Environ Microbiol* 72(5): 3284-3290
- Hart A, Roelofs W, Crocker J, *et al.* 2007. *Defra Research Contract CPEC38: Quantitative approaches to the Risk Assessment of GM crops*. Central Science Laboratory: York.
- Hartigan JA. 1975. *Clustering algorithms*. John Wiley & Sons Inc., New York
- Hayes K, Regan H, Burgman M, *et al.* 2006. Methods to address uncertainty in ecological risk assessment of Genetically Modified Organisms, *in* Ninth International Symposium on the Biosafety of Genetically Modified Organisms, Jeju Island, South Korea, Sept. 24 29 2006
- Hayes EH and Landis WG. 2004. Regional ecological risk assessment of a near shore marine environment: Cherry Point, WA. *Hum Ecol Risk Assess* 10(2): 299-325
- Hays SM, Aylward LL, Gagné M, *et al.* 2009. Derivation of Biomonitoring Equivalents for cyfluthrin. *Regul Toxicol Pharm* 55(3): 268-275
- Hays SM, Aylward LL, Gagné M, *et al.* 2009. Derivation of Biomonitoring Equivalents for cyfluthrin. *Regul Toxicol Pharm* 55(3): 268-275
- Helton JC, Johnson JD, and Sallaberry CJ. 2011. Quantification of margins and uncertainties: Example analyses from reactor safety and radioactive waste disposal involving the separation of aleatory and epistemic uncertainty. *Reliab Eng Syst Safe* 96(9):1014-1033
- Helton JC. 1994. Treatment of uncertainty in performance assessments for complex systems. *Risk Anal* 14:483-511

- Henning-De Jong I, van Zelm R, Huijbregts MAJ, *et al.* 2008. Ranking of agricultural pesticides in the Rhine-Meuse-Scheldt basin based on toxic pressure in marine ecosystems. *Environ Toxicol Chem* 27(3): 737-745
- Henrion M and Fischhoff B. 1986. Assessing uncertainty in physical constants. *Am J Phys* 54:791
- Hoffman F and Hammonds J. 1994. Propagation of uncertainty in risk assessments: the need to distinguish between uncertainty due to lack of knowledge and uncertainty due to variability. *Risk Anal* 14:707-712
- Hromkovic J. 2005. *Design and analysis of randomized algorithms: introduction to design paradigms*. Springer-Verlag Inc., New York
- Hughes K, Meek ME, Walker M, *et al.* 2003. 1,3-Butadiene: Exposure estimation, hazard characterization, and exposure-response analysis. *J Toxicol Environ Health Part B* 6(1): 55-83
- Huijbregts MAJ, Norris G, Bretz R, *et al.* 2001. Framework for modelling data uncertainty in life cycle inventories. *Int J Life Cycle Ass* 63:127-132
- Hung ML, Wu SY, Chen YC, *et al.* 2009. The Health Risk Assessment of Pb and Cr leached from fly ash monolith landfill. *Journal of Hazard Mater* 172(1): 316-323
- Huysmans M, Madarász T, and Dassargues A. 2006. Risk assessment of groundwater pollution using sensitivity analysis and a worst-case scenario analysis. *Environ Geol* 502:180-193
- Jackson LE, Bird SL, Matheny RW, *et al.* 2004. A regional approach to projecting land-use change and resulting ecological vulnerability. *Environ Monit Assess* 94(1-3): 231-248
- Janssen P, Petersen A, van der Sluijs J, *et al.* 2003. *RIVM/MNP guidance for uncertainty assessment and communication: Quickscan hints & actions list*. RIVM/MNP Guidance for Uncertainty Assessment and Communication Series 2

- Jones DR, Peters JL, Rushton L, *et al.* 2009. Interspecies extrapolation in environmental exposure standard setting: A Bayesian synthesis approach. *Regul Toxicol Pharm* 53(3): 217-225
- Jorgensen R, Hauser T, D'Hertefeldt T, *et al.* 2009. The variability of processes involved in transgene dispersal - case studies from Brassica and related genera. *Environ Sci Pollut R* 16(4):389-395
- Kaloudis S, Tocatlidou A, Lorentzos NA, *et al.* 2005. Assessing wildfire destruction danger: a decision support system incorporating uncertainty. *Ecol Modell* 181(1): 25-38
- Kandlikar M, Ramachandran G, Maynard A, *et al.* 2007. Health risk assessment for nanoparticles: A case for using expert judgment. *JNR* 91:137-156
- Kapo KE and Burton Jr. GA. 2006. A geographic information systems-based, weights-of-evidence approach for diagnosing aquatic ecosystem impairment. *Environ Toxicol Chem* 25(8): 2237-2249
- Keiter S, Braunbeck T, Heise S, *et al.* 2009. A fuzzy logic-classification of sediments based on data from in vitro biotests. *J Soil Sediment* 9(3): 168-179
- Kelly M, Bennion H, Burgess A, *et al.* 2009. Uncertainty in ecological status assessments of lakes and rivers using diatoms. *Hydrobiologia* 633(1): 5-15
- Kelly E and Campbell K. 2000. Separating variability and uncertainty in environmental risk assessment - making choices. *Hum Ecol Risk Assess* 6(1):1-13
- Kentel E and Aral MM. 2007. Risk tolerance measure for decision-making in fuzzy analysis: A health risk assessment perspective. *Stoch Env Res Risk A* 21(4): 405-417
- van der Keur P, Henriksen H, Refsgaard J, *et al.* 2008. Identification of major sources of uncertainty in current IWRM practice. Illustrated for the Rhine basin. *Water resour manag* 22(11):1677-1708

958 Khan F, Sadiq R, and Husain T. 2002. GreenPro-I: a risk-based life cycle assessment and
 959 decision-making methodology for process plant design. *Environ Modell Softw* 17(8):
 960 669-692

961 King RS and Richardson CJ. 2003. Integrating bioassessment and ecological risk assessment:
 962 an approach to developing numerical water-quality criteria. *Environ Manage* 31: 795-
 963 809

964 Klier C, Grundmann S, Gayler S, *et al.* 2008. Modelling the environmental fate of the
 965 herbicide glyphosate in soil lysimeters. *Water, Air, and Soil Pollution: Focus* 82:187-
 966 207

967 Knol AB, Petersen AC, van der Sluijs JP, *et al.* 2009. Dealing with uncertainties in
 968 environmental burden of disease assessment. *Environ Health* 8:21

969 Kooistra L, Huijbregts MAJ, Ragas AMJ, *et al.* 2005. Spatial variability and uncertainty
 970 ecological risk assessment: A case study on the potential risk of cadmium for the little
 971 owl in a Dutch river flood plain. *Environ Sci Technol* 39(7): 2177-2187

972 Kraymer von Krauss M, Casman EA, and Small MJ. 2004. Elicitation of expert judgments of
 973 uncertainty in the risk assessment of herbicide-tolerant oilseed crops. *Risk Anal*
 974 24:1515-1527

975 Kumar V, Mari M, Schuhmacher M, *et al.* 2009. Partitioning total variance in risk
 976 assessment: Application to a municipal solid waste incinerator. *Environ Modell Softw*
 977 242:247-261

978 Landis WG, Duncan PB, Hayes EH, *et al.* 2004. A regional retrospective assessment of the
 979 potential stressors causing the decline of the cherry point pacific herring run. *Hum Ecol*
 980 *Risk Assess* 10(2): 271-297

981 Lee JJ, Jang CS, Liang CP, *et al.* 2008. Assessing carcinogenic risks associated with
 982 ingesting arsenic in farmed smeltfish (*Ayu, Plecoglossus altirelis*) in aseniasis-endemic
 983 area of Taiwan. *Sci Total Environ* 403 1-3): 68-79
 984 Lemke LD and Bahrou AS. 2009. Partitioned multiobjective risk modeling of carcinogenic
 985 compounds in groundwater. *Stoch Env Res Risk A* 23(1): 27-39
 986 Li HL, Huang GH, and Zou Y. 2008. An integrated fuzzy-stochastic modeling approach for
 987 assessing health-impact risk from air pollution. *Stoch Environ Res Risk Assess* 226:789-
 988 803
 989 Li J, Huang GH, Zeng G, *et al.* 2007. An integrated fuzzy-stochastic modeling approach for
 990 risk assessment of groundwater contamination. *J Environ Manage* 822:173-188
 991 Li J, Liu L, Huang G, *et al.* 2006. A fuzzy-set approach for addressing uncertainties in risk
 992 assessment of hydrocarbon-contaminated site. *Water, Air, Soil Pollut* 171(1-4): 5-18
 993 Liao CM and Chou BYH. 2005. Predictive risk thresholds for survival protection of farmed
 994 abalone, *Haliotis diversicolor supertexta*, exposed to waterborne zinc. *Environmental*
 995 *Toxicology* 20(2): 202-211
 996 Lindenschmidt KE, Huang S, and Baborowski M. 2008. A quasi-2D flood modeling
 997 approach to simulate substance transport in polder systems for environment flood risk
 998 assessment. *Sci Total Environ* 397(1-3): 86-102
 999 Linkov I and Moberg E. 2011. Multi-Criteria Decision Analysis: Environmental Applications
 1000 and Case Studies. CRC Press, Boca Raton, Florida
 1001 Linkov I, Loney D, Cormier S, *et al.* 2009. Weight-of-evidence evaluation in environmental
 1002 assessment: Review of qualitative and quantitative approaches. *ScTEn* 40719:5199-
 1003 5205
 1004 Linkov I, Satterstrom FK, Steevens J, *et al.* 2007. Multi-criteria decision analysis and
 1005 environmental risk assessment for nanomaterials. *JNR* 94:543-554

1006 Linkov I and Burmistrov D. 2005. Sources of uncertainty in model predictions: Lessons
 1007 learned from the IAEA Forest and Fruit Working Group model intercomparisons. *J*
 1008 *Environ Radioact* 84(2): 297-314

1009 Linkov I, Von Stackelberg KE, Burmistrov D, *et al.* 2001. Uncertainty and variability in risk
 1010 from trophic transfer of contaminants in dredged sediments. *ScTEn* 2741-3:255-269

1011 Liu ZQ, Zhang YH, Li GH, *et al.* 2007. Sensitivity of key factors and uncertainties in health
 1012 risk assessment of benzene pollutant. *Journal of Environmental Sciences* 19(10): 1272-
 1013 1280

1014 Loos M, Ragas AMJ, Plasmeijer R, *et al.* 2010. Eco-SpaCE: An object-oriented, spatially
 1015 explicit model to assess the risk of multiple environmental stressors on terrestrial
 1016 vertebrate populations. *Sci Total Environ* 408(18): 3908-3917

1017 Lu H, Axe L, and Tyson TA. 2003. Development and application of computer simulation
 1018 tools for ecological risk assessment. *Environmental Modeling and Assessment* 8(4):
 1019 311-322

1020 Luttik R, Mineau P, and Roelofs W. 2005. A review of interspecies toxicity extrapolation in
 1021 birds and mammals and a proposal for long-term toxicity data. *Ecotoxicology* 14(8):
 1022 817-832

1023 Ma HW. 2002. Stochastic multimedia risk assessment for a site with contaminated
 1024 groundwater. *J Environ Manage* 166:464-478

1025 Ma WC and van der Voet H. 1993. A risk-assessment model for toxic exposure of small
 1026 mammalian carnivores to cadmium in contaminated natural environments. *Sci Total*
 1027 *Environ* (Suppl. 2): 1701-1714

1028 Maier HR, Ascough II JC, Wattenbach M, *et al.* 2008. Chapter Five - Uncertainty in
 1029 Environmental Decision Making: Issues, Challenges and Future Directions,
 1030 Developments in Integrated Environmental Assessment 3:69-85

1031 Matson CW, Gillespie AM, CMcCarthy C, *et al.* 2009. Wildlife toxicology: Biomarkers of
 1032 genotoxic exposures at a hazardous waste site. *Ecotoxicology* 18(7): 886-898

1033 Maxim LD and McConnell EE. 2001. Interspecies comparisons of the toxicity of asbestos
 1034 and synthetic vitreous fibers: A weight-of-the-evidence approach. *Regul Toxicol Pharm*
 1035 33(3): 319-342

1036 Maxwell RM and Kastenbergs WE. 1999. Stochastic environmental risk analysis: An
 1037 integrated methodology for predicting cancer risk from contaminated groundwater.
 1038 *Stoch Env Res Risk A* 13(1-2): 27-47

1039 Maycock BJ and Benford DJ. 2007. Risk assessment of dietary exposure to methylmercury in
 1040 fish in the UK. *Human and Experimental Toxicology* 26(3): 185-190

1041 McDonald BG and Wilcockson JB. 2003. Improving the use of toxicity reference values in
 1042 wildlife food chain modeling and ecological risk assessment. *Hum Ecol Risk Assess*
 1043 9(7): 1585-1594

1044 Meek ME, Beauchamp R, Long G, *et al.* 2002. Chloroform: Exposure estimation, hazard
 1045 characterization, and exposure-response analysis. *J Toxicol Environ Health Part B* 5(3):
 1046 283-334

1047 Meek ME and Hughes K. 1995. Approach to health risk determination for metals and their
 1048 compounds under the Canadian Environmental Protection Act. *Regul Toxicol Pharm*
 1049 22(3):206-212

1050 Merz B and Thielen AH. 2009. Flood risk curves and uncertainty bounds. *Nat Hazards* 513:
 1051 437-458

1052 Meyer V, Scheuer S, and Haase D. 2009. A multicriteria approach for flood risk mapping
 1053 exemplified at the Mulde river, Germany. *Nat Hazards* 48(1): 17-39

1054 Morgan M, Henrion M and Small M. 1990. *Uncertainty: a guide to dealing with uncertainty*
 1055 *in quantitative risk and policy analysis*. Cambridge University Press, Cambridge, UK

1056 Mugglestone MA, Stutt ED, and Rushton L. 2001. Setting microbiological water quality
1057 standards for sea bathing - A critical evaluation. *Water Sci Technol* 43(12): 9-18

1058 Mukhtasor HT, Veitch B, and Bose N. 2004. An ecological risk assessment methodology for
1059 screening discharge alternatives of produced water. *Hum Ecol Risk Assess* 10(3): 505-
1060 524

1061 Naito W, Gamo Y, and Yoshida K. 2006. Screening-level risk assessment of Di(2-
1062 ethylhexyl)phthalate for aquatic organisms using monitoring data in Japan. *Environ*
1063 *Monit Assess* 115(1-3): 451-471

1064 Nasiri F, Huang G, and Fuller N. 2007. Prioritizing groundwater remediation policies: A
1065 fuzzy compatibility analysis decision aid. *J Environ Manage* 82(1): 13-23

1066 de Nazelle A and Rodríguez DA. 2009. Tradeoffs in incremental changes towards pedestrian-
1067 friendly environments: Physical activity and pollution exposure. *Transpor Res D-TR E*
1068 14(4): 255-263

1069 Nazir M, Khan F, Amyotte P, *et al.* 2008. Subsea release of oil from a riser: an ecological
1070 risk assessment. *Risk Anal* 28(5): 1173-119

1071 Neuhäuser B and Terhorst B. 2007. Landslide susceptibility assessment using "weights-of-
1072 evidence" applied to a study area at the Jurassic escarpment (SW-Germany).
1073 *Geomorphology* 86(1-2): 12-24

1074 de Nijs TCM, Toet C, Vermeire TG, *et al.* 1993. Dutch risk assessment system for new
1075 chemicals: DRANC. *Sci Total Environ* (Suppl. 2): 1729-1748

1076 Oughton DH, Agüero A, Avila R, *et al.* 2008. Addressing uncertainties in the ERICA
1077 Integrated Approach. *J. Environ. Radioactivity* 99:1384-1392

1078 Park RA, Clough JS, and Wellman MC. 2008. AQUATOX: Modeling environmental fate and
1079 ecological effects in aquatic ecosystems. *Ecol Modell* 213(1): 1-15

1080 Pascoe GA, Blanchet RJ, and Linder G. 1993. Ecological risk assessment of a metals-
1081 contaminated wetland: Reducing uncertainty. *Sci Total Environ* (Suppl. 2): 1715-1728

1082 Persson K and Destouni G. 2009. Propagation of water pollution uncertainty and risk from
1083 the subsurface to the surface water system of a catchment. *J Hydrol* 377(3-4): 434-444

1084 Petersen AC. 2006. *Simulating nature: a philosophical study of computer-simulation*
1085 *uncertainties and their role in climate science and policy advice*. Aksant Academic
1086 Pub, Amsterdam

1087 Phillips KP, Foster WG, Leiss W, *et al.* 2008. Assessing and managing risks arising from
1088 exposure to endocrine-active chemicals. *J Toxicol Env Heal B: Critical Reviews* 113-
1089 4:351-372

1090 Pollino CA, Woodberry O, Nicholson A, *et al.* 2007. Parameterisation and evaluation of a
1091 Bayesian network for use in an ecological risk assessment. *Environ Modell Softw* 22(8):
1092 1140-1152

1093 Proctor DM, Shay EC, Fehling KA, *et al.* 2002. Assessment of human health and ecological
1094 risks posed by the uses of steel-industry slags in the environment. *Hum Ecol Risk*
1095 *Assess* 8(4): 681-711

1096 Prudhomme C, Jakob D, and Svensson C. 2003. Uncertainty and climate change impact on
1097 the flood regime of small UK catchments. *J Hydrol* 277(1-2): 1-23

1098 Qin XS and Huang GH. 2009. Characterizing uncertainties associated with contaminant
1099 transport modeling through a coupled fuzzy-stochastic approach. *Water Air Soil Poll*
1100 1971-4:331-348

1101 Ramirez AJ, Jensen AC, and Cheng BHC. 2012. A taxonomy of uncertainty for dynamically
1102 adaptive systems. *ICSE Workshop on Software Engineering for Adaptive and Self-*
1103 *Managing Systems*:99-108

1104 Ranke J. 2002. Persistence of antifouling agents in the marine biosphere. *Environ Sci Technol*
1105 36(7): 1539-1545

1106 Refsgaard J, van Der Sluijs J, Hojberg A, *et al.* 2007. Uncertainty in the environmental
1107 modelling process-A framework and guidance. *Environ Modell Softw* 22:1543-1556

1108 Regan HM, Colyvan M, and Burgman MA. 2002. A taxonomy and treatment of uncertainty
1109 for ecology and conservation biology. *Ecol Appl* 12-2:618-628

1110 Rowe W. 1994. Understanding uncertainty. *Risk Anal* 14:743-750

1111 Rutherford L, Garron C, Ernst W, *et al.* 2003. The aquatic environment and textile mill
1112 effluents - An ecological risk assessment. *Hum Ecol Risk Assess* 9(2): 589-606

1113 Sander P and Öberg T. 2006. Comparing deterministic and probabilistic risk assessments: A
1114 case study at a closed steel mill in southern Sweden. *J Soil Sediment* 6(1): 55-61

1115 Sanderson H, Laird B, Pope L, *et al.* 2007. Assessment of the environmental fate and effects
1116 of ivermectin in aquatic mesocosms. *Aquatic Toxicology* 85(4): 229-240

1117 Sanderson H, Dyer SD, Price BB, *et al.* 2006. Occurrence and weight-of-evidence risk
1118 assessment of alkyl sulfates, alkyl ethoxysulfates, and linear alkylbenzene sulfonates
1119 (LAS) in river water and sediments. *Sci Total Environ* 368(2-3): 695-712

1120 Scherm H. 2000. Simulating uncertainty in climate-pest models with fuzzy numbers. *Environ*
1121 *Pollut* 108(3): 373-379

1122 Schoeny R, Haber L, and Dourson M. 2006. Data considerations for regulation of water
1123 contaminants. *Toxicology* 221(2-3): 217-224

1124 Schwartz S, Berding V, and Matthies M. 2000. Aquatic fate assessment of the polycyclic
1125 musk fragrance HHCB scenario and variability analysis in accordance with the EU risk
1126 assessment guidelines. *Chemosphere* 41(5): 671-679

1127 Scott MJ, Brandt CA, Bunn AL, *et al.* 2005. Modeling long-term risk to environmental and
 1128 human systems at the Hanford nuclear reservation: Scope and findings from the initial
 1129 model. *Environ Manage* 35(1): 84-98

1130 Shakhawat C, Tahir H, and Neil B. 2006. Fuzzy rule-based modelling for human health risk
 1131 from naturally occurring radioactive materials in produced water. *J Environ Radioact*
 1132 89(1): 1-17

1133 van der Sluijs J. 1997. *Anchoring amid uncertainty: on the management of uncertainties in*
 1134 *risk assessment of anthropogenic climate change*. PhD thesis, Utrecht University

1135 Smith JT, Walker LA, Shore RF, *et al.* 2009. Do estuaries pose a toxic contamination risk for
 1136 wading birds? *Ecotoxicology* 18(7): 906-917

1137 Smith PN, Cobb GP, Godard-Codding C, *et al.* 2007. Contaminant exposure in terrestrial
 1138 vertebrates. *Environ Pollut* 150(1): 41-64

1139 Son HV, Ishihara S, and Watanabe H. 2006. Exposure risk assessment and evaluation of the
 1140 best management practice for controlling pesticide runoff from paddy fields. Part 1:
 1141 Paddy watershed monitoring. *Pest Manag Sci* 62(12): 1193-1206

1142 van Sprang PA, Verdonck FAM, van Assche F, *et al.* 2009. Environmental risk assessment of
 1143 zinc in European freshwaters: A critical appraisal. *Sci Total Environ* 407(20): 5373-
 1144 5391

1145 Staples CA, Woodburn K, Caspers N, *et al.* 2002. A weight of evidence approach to the
 1146 aquatic hazard assessment of bisphenol A. *Hum Ecol Risk Assess* 8(5): 1083-1105

1147 Stephens M, Goodwin B, and Andres T. 1993. Deriving parameter probability density
 1148 functions. *Reliab Eng Sys Safe* 42(2-3):271-291

1149 Stevens G, de Foy B, West JJ, *et al.* 2007. Developing intake fraction estimates with limited
 1150 data: Comparison of methods in Mexico City. *Atmos Environ* 41(17): 3672-3683

1151 Stirling A. 2012. Opening up the politics of knowledge and power in bioscience. *PLoS Biol*
1152 10(1): e1001233

1153 Stirling A. 1999. Risk at a turning point? *J Risk Res* 1:97-109

1154 Suter II GW and Cormier SM. 2011. Why and how to combine evidence in environmental
1155 assessments: weighing evidence and building cases. *Sci Total Environ* 409:1406-
1156 1417

1157 Teunis PFM, Medema GJ, Kruidenier L, *et al.* 1997. Assessment of the risk of infection by
1158 *Cryptosporidium* or *Giardia* in drinking water from a surface water source. *Water Res*
1159 31(6): 1333-1346

1160 Therriault TW and Herborg LM. 2008. A qualitative biological risk assessment for vase
1161 tunicate *Ciona intestinalis* in Canadian waters: Using expert knowledge. *ICES J Mar*
1162 *Sci* 65(5): 781-787

1163 Thorsen M, Refsgaard JC, Hansen S, *et al.* 2001. Assessment of uncertainty in simulation of
1164 nitrate leaching to aquifers at catchment scale. *J Hydrol* 242(3-4): 210-227

1165 Tillman Jr. FD and Weaver JM. 2006. Uncertainty from synergistic effects of multiple
1166 parameters in the Johnson and Ettinger (1991) vapor intrusion model. *Atmos Environ*
1167 40(22): 4098-4112

1168 Troldborg, M. 2010. *Risk assessment models and uncertainty estimation of groundwater*
1169 *contamination from point sources*. PhD Thesis, Technical University of Denmark

1170 Tsuji JS, Benson R, Schoof RA, *et al.* 2004. Health effect levels for risk assessment of
1171 childhood exposure to arsenic. *Regul Toxicol Pharm* 39(2): 99-110

1172 Twining JR and Cameron RF. 1997. Decision-making processes in ecological risk assessment
1173 using copper pollution of macquarie harbour from Mt. Lyell, Tasmania, as a case study.
1174 *Hydrobiologia* 352(1-3): 207-218

1175 United States Environmental Protection Agency. 1998. *Guidelines for ecological risk*
1176 *assessment*. Washington: United States Environmental Protection Agency, Risk
1177 Assessment Forum.

1178 Vallack HW, Bakker DJ, Brandt I, *et al.* 1998. Controlling persistent organic pollutants-what
1179 next? *Environ Toxicol Pharmacol* 6(3): 143-175

1180 Verdonck FAM, van Sprang PA, and Vanrolleghem PA. 2008. An intelligent data collection
1181 tool for chemical safety/risk assessment. *Chemosphere* 70(10): 1818-1826

1182 Vesely W and Rasmuson D. 1984. Uncertainties in nuclear probabilistic risk analyses. *Risk*
1183 *Anal* 4:313-322

1184 Vose D. 2000. *Risk Analysis*. John Wiley & Sons, Chichester

1185 Walker WE, Harremoës P, Rotmans J, *et al.* 2003. Defining Uncertainty: A Conceptual Basis
1186 for Uncertainty Management in Model-Based Decision Support. *Integrated Assessment*
1187 41:5-17

1188 Walker R, Landis W, and Brown P. 2001. Developing a regional ecological risk assessment:
1189 A case study of a Tasmanian agricultural catchment. *Hum Ecol Risk Assess* 7(2): 417-
1190 439

1191 Wang B, Yu G, Huang J, *et al.* 2009. Tiered aquatic ecological risk assessment of
1192 organochlorine pesticides and their mixture in Jiangsu reach of Huaihe River, China.
1193 *Environ Monit Assess* 157(1-4): 29-42

1194 Weyers A, Sokull-Klüttgen B, Knacker T, *et al.* 2004. Use of Terrestrial Model Ecosystem
1195 Data in Environmental Risk Assessment for Industrial Chemicals, Biocides and Plant
1196 Protection Products in the EU. *Ecotoxicology* 13(1-2): 163-176

1197 Wiegers JK, Feder HM, Mortensen LS, *et al.* 1998. A regional multiple-stressor rank-based
1198 ecological risk assessment for the fjord of Port Valdez, Alaska. *Hum Ecol Risk Assess*
1199 4(5): 1125-1173

1200 Williams BK, Szaro RC, and Shapiro CD. 2009. *Adaptive Management: The U.S.*
1201 *Department of the Interior Technical Guide*. Adaptive Management Working Group,
1202 Washington, DC

1203 Wright-Walters M, Volz C, Talbott E, *et al.* 2011. An updated weight of evidence approach
1204 to the aquatic hazard assessment of Bisphenol A and the derivation a new predicted no
1205 effect concentration (Pnec) using a non-parametric methodology. *Sci Total Environ*
1206 409(4):676-685

1207 Wu FC and Tsang YP. 2004. Second-order Monte Carlo uncertainty/variability analysis using
1208 correlated model parameters: Application to salmonid embryo survival risk assessment.
1209 *Ecol Model* 1773-4:393-414

1210 Wynne B. 1992. Uncertainty and environmental learning: Reconceiving science and policy in
1211 the preventive paradigm. *Global Environ Chang* 2:111-127

1212 Xiao X, Zhang Y, and Li S. 2008. Ecological risk assessment of DDT accumulation in
1213 aquatic organisms of Taihu Lake, China. *Hum Ecol Risk Assess* 14(4): 819-834

1214 Zadeh LA. 1965. Fuzzy sets. *Inform Control* 83:338-353

1215 Zalk D, Paik S, and Swuste P. 2009. Evaluating the Control Banding Nanotool: a qualitative
1216 risk assessment method for controlling nanoparticle exposures. *J Nanopart Res*
1217 11(7):1685-1704

1218 Zhang H, Huang GH, and Zeng GM. 2009. Health risks from arsenic-contaminated soil in
1219 Flin Flon-Creighton, Canada: Integrating geostatistical simulation and dose-response
1220 model. *Environ Pollut* 157(8-9): 2413-2420

1221 Zimmermann HJ. 2001. *Fuzzy Set Theory and Its Applications*. Kluwer Academic Publishers,
1222 Norwell, Massachusetts

Table 1 Uncertainty categorisations offered by existing typologies within the published environmental risk-based literature.

Source reference	Uncertainties included within source
Vesely and Rasmuson 1984	Data; Model (understanding, approximation); Completeness; Physical variability
Henrion and Fischhoff 1986	Random; Systematic
Alcamo and Bartnicki 1987	Model (structure, parameters, forcing, initial state, operation)
Beck 1987	Model (aggregation, structure, numerical, parameter); Variability; Errors;
Morgan and Henrion 1990	Statistical variation; Systematic error; Linguistic; Variability; Inherent randomness; Disagreement; Model (approximation, form)
Finkel 1990	Model; Parameter; Decision; Natural variability
Funtowicz and Ravetz 1990	Inexactness; Unreliability; Border with ignorance
Wynne 1992	Risk; Uncertainty; Ignorance; Indeterminacy;
Helton 1994	Stochastic; Subjective
Hoffman and Hammonds 1994	Lack of knowledge; Variability
Rowe 1994	Temporal; Structural; Metrical; Translational
Faucheux and Froger 1995	Ignorance; Strong uncertainty; Uncertainty; Certainty
van der Sluijs 1997	Inexactness; Unreliability; Ignorance; Model (input data, conceptual model structure, technical model structure, bugs, model completeness)
Stirling 1999	Risk; Uncertainty; Ambiguity; Ignorance
Bedford and Cooke 2001	Aleatory; Epistemic; Parameter; Data; Model; Ambiguity; Volitional
Huijbregts <i>et al.</i> 2001	Parameter; Model; Choices; Variability (spatial, temporal, between source and object)
Bevington and Robinson 2002	Systematic errors; Random errors

Regan <i>et al.</i> 2002	Epistemic (measurement error, systematic error, natural variation, inherent randomness, model, subjective judgement); Linguistic (vagueness, context dependence, ambiguity, underspecificity, indeterminacy of theoretical terms)
van Asselt and Rotmans 2002	Variability (nature, cognitive, behavioural, societal, technological); Knowledge (inexactness, lack of measurements, practically immeasurable, conflicting evidence, ignorance, indeterminacy)
Janssen <i>et al.</i> 2003	Statistical; Scenario; Recognised ignorance; Knowledge-based; Variability-based; Context; Expert judgement; Model (structure, technical, parameters, input); Data; Outputs
Walker <i>et al.</i> 2003	Statistical; Scenario; Recognised ignorance; Total ignorance; Epistemic; Variability; Context; Model (structure, technical, parameters, input, outputs)
Brown 2004	Bounded uncertainty; Unbounded uncertainty; Indeterminacy; Ignorance
Dewulf <i>et al.</i> 2005	Inherent nature of phenomena; Lack of knowledge; Ambiguity in system understanding
Beer 2006	Probabilistic; Ambiguity; Incertitude; Ignorance; Indeterminacy
Petersen 2006	Location; Nature; Range; Recognised ignorance; Methodological unreliability; Value diversity
Hayes <i>et al.</i> 2006	Linguistic; Variability; Incertitude
Maier <i>et al.</i> 2008	Data (measurement error, type of data, length of record, analysis); Model (method, record quality, calibration, validation, experience); Human (stakeholder, politics)
Ascough II <i>et al.</i> 2008	Knowledge; Variability; Linguistic; Process; Model; Variability; Linguistic; Decision
Brouwer and Blois 2008	Statistical; Scenario; Qualitative; Recognised ignorance
Knol <i>et al.</i> 2009	Statistical; Scenario; Recognised ignorance; Epistemic; Ontic (process, normative); Model (structure, parameters, input data); Methodological; Analyst uncertainty

Table 2 Novel typology of uncertainties (including definitions) resulting from the analysis and iterative clustering of data obtained from 171 ERAs that applied WOE methods.

Nature	Location	Sub-location	Definition
Epistemic	Data	Availability	referring to the incompleteness, scarcity, or absence of data
		Precision	concerning the lack of accuracy or precision in obtained data
		Reliability	reflecting its trustworthiness i.e. data is erroneous for some specified reason
	Language	Ambiguity	where multiple meanings are possible
		Underspecificity	where meanings are not exact
		Vagueness	where meanings are not clear and understandable
	System	Cause	concerning a lack of clarity regarding the source(s) of harm
		Effect	relating to the influence a particular stressor (source) has upon the receptor(s)
		Process	where the risks are not understood or a process vital to a successful assessment is not identified
Aleatory	Variability	Human	results primarily from intentionally biased and subjective actions, but extends to all qualities of humans which are, either literally or from the viewpoint of the risk analyst, stochastic in nature
		Natural	pertains to the stochastic traits of natural systems
	Extrapolation	Intraspecies	where information specific to members of a species is used to represent other members of the same species
		Interspecies	where information specific to members of a species is used to represent members of a different species
		Laboratory	where information specific to laboratory conditions is used to represent real-world scenarios

		Quantity	where information specific to one quantity is used to represent another
		Spatial	where information specific to one spatial scale is used to represent another
		Temporal	where information specific to one timescale is used to represent another
Combined	Model	Structure	concerning the representation of real-world processes in model form
		Output	reflecting the level of confidence in the produced results
	Decision	Decision	where doubt surrounds an optimal course of action, often in the face of differing objectives.

Table 3 Descriptions of the most frequently occurring uncertainty management techniques, organised according to the percentage rates with which they occur in the evidence base of 171 ERAs that applied WOE methods, along with their associated uncertainties.

Uncertainty management technique	Description	Associated uncertainty locations	Referenced in:
Monte-Carlo simulation (22.1%)	Utilises repeated executions of numerical models to simulate stochastic processes.	Data, Extrapolation, Variability, Model, System	Ma 2002 Qin and Huang 2009
Uncertainty factor (16.6%)	Attaches a factor-based correction to the data being used which reflects the level of uncertainty within it.	Extrapolation, System, Data, Variability	Calabrese 1994 Phillips <i>et al.</i> 2008
Sensitivity analysis (8.4%)	Tests the sensitivity of a chosen output variable to variations in quantities relating to input variables.	Data, Model, Extrapolation, System	Huysmans <i>et al.</i> 2006 Oughton <i>et al.</i> 2008
No action (7.7%)	Not attempting to quantify, reduce, or manage uncertainties, whether recognised by the publication author(s) or identified through this research.	Data, Extrapolation, System, Variability, Model	Cesar <i>et al.</i> 2009
Further data collection (7.3%)	The collection of increased quantities of data.	Extrapolation, Data, Variability	Avagliano and Parella 2009
Fuzzy logic (6.8%)	A form of multi-valued logic that allows its components to be approximate rather than precise.	Data, Language, Model, Variability	Zadeh 1965 Acosta <i>et al.</i> 2010
Expert elicitation (4.6%)	Seeks to capture the knowledge of one or more experts in a field with regard to a specific matter.	Data, System , Variability	Kandlikar <i>et al.</i> 2007
Probability density function ¹	Describes the frequency of occurrence for different	Data, Variability	Oughton <i>et al.</i> 2008

(4.0%)	parameter values over a given range.		
Latin hypercube sampling (3.5%)	Splits a distribution into distinct intervals for sampling and use as inputs to a numerical model.	Data, Variability	Klier <i>et al.</i> 2008 Kumar <i>et al.</i> 2009
Bayesian belief network (3.1%)	A graphical representation of a system, in which relationships between uncertain characteristics are expressed through probability values.	Variability, Data, System	Aspinall <i>et al.</i> 2003
Fuzzy-stochastic system (3.1%)	A hybrid approach for incorporating epistemic and stochastic uncertainties separately.	Data, Extrapolation, Language	Li <i>et al.</i> 2007 Kumar <i>et al.</i> 2009
Precautionary management (1.8%)	Management based upon the application of the Precautionary Principle.	Extrapolation, System	Godduhn and Duffy 2003
Multi-criteria decision analysis (1.1%)	Brings together criteria and performance scores to provide a basis for integrating risk and uncertainty levels.	Decision	Linkov <i>et al.</i> 2007 Critto <i>et al.</i> 2007
Adaptive management (0.4%)	Incorporate the needs of many into an iterative system where differing alternatives and objectives are present.	Decision	Dey <i>et al.</i> 2000 Williams <i>et al.</i> 2009
¹ Refers to probability density functions that are applied independently of the Monte-Carlo simulation and Latin-hypercube sampling techniques.			

Table 4 Ranked potential uncertainties (according to the percentage proportion with which they occur in the evidence base of 171 ERAs that applied WOE methods) for risk analysts to consider, detailing uncertainty locations, sub-locations, and natures, along with related uncertainty management techniques in order of decreasing appropriateness. Row shadings correspond to the uncertainties that can be quantified, reduced and potentially removed (epistemic ) , quantified at best (aleatory ) , and those that must be considered on a case-by-case basis (combined ) .

Rank	Location of uncertainty	Sub-location of uncertainty	Nature of uncertainty	Uncertainty management technique #1	Uncertainty management technique #2	Uncertainty management technique #3
1 (20.8%)	Data	Reliability	Epistemic	Monte-Carlo simulation	Sensitivity analysis	Uncertainty factors
2 (10.1%)	Data	Availability	Epistemic	Monte-Carlo simulation	Sensitivity analysis	Uncertainty factors
3 (9.9%)	Variability	Natural	Aleatory	Monte-Carlo simulation	Further data collection	Uncertainty factors
4 (7.8%)	Extrapolation	Interspecies	Aleatory	Uncertainty factors	Monte-Carlo simulation	Further data collection
=5 (6.0%)	Extrapolation	Spatial	Aleatory	Interpolation	Monte-Carlo simulation	Uncertainty factors
=5 (6.0%)	System	Process	Epistemic	Uncertainty factors	Monte-Carlo simulation	Expert elicitation
7 (5.2%)	Extrapolation	Intraspecies	Aleatory	Uncertainty factors	Monte-Carlo simulation	Further data collection
=8 (4.2%)	Extrapolation	Temporal	Aleatory	Further data collection	Uncertainty factors	Monte-Carlo simulation
=8 (4.2%)	Model	Output	Combined	Sensitivity analysis	Monte-Carlo simulation	Fuzzy logic
=10 (3.9%)	Model	Structure	Combined	Sensitivity analysis	Monte-Carlo simulation	Fuzzy logic
=10 (3.9%)	System	Effect	Epistemic	Uncertainty factors	Expert elicitation	Monte-Carlo simulation
=12 (3.1%)	Extrapolation	Laboratory	Aleatory	Uncertainty factors	Further data collection	Monte-Carlo simulation
=12 (3.1%)	System	Cause	Epistemic	Uncertainty factors	Further data collection	Monte-Carlo simulation
=14 (2.3%)	Extrapolation	Quantity	Aleatory	Uncertainty factors	Further data collection	Monte-Carlo simulation
=14 (2.3%)	Variability	Human	Aleatory	Bayesian belief networks	Expert elicitation	Sensitivity analysis

=16 (1.6%)	Data	Precision	Epistemic	Fuzzy logic	Expert elicitation	Monte-Carlo simulation
=16 (1.6%)	Decision	Decision	Combined	Adaptive management	MCDA	Bayesian belief networks
=16 (1.6%)	Language	Ambiguity	Epistemic	Fuzzy logic	Fuzzy-stochastic	N/A
=16 (1.6%)	Language	Vagueness	Epistemic	Fuzzy logic	Fuzzy-stochastic	N/A
20 (1.0%)	Language	Underspecificity	Epistemic	Fuzzy logic	Fuzzy-stochastic	N/A

Table 5 Major elements of an environmental risk assessment, derived from government guidance documents (Fairman *et al.* 1998; USEPA 1998; DEFRA 2011).

Assessment phase	Assessment task	Assessment sub-task
Problem formulation	Build a conceptual model	Define risk relationships e.g. source-pathway-receptor paradigm
		Select assessment endpoints e.g. development; behaviour; survival; fecundity; abundance
		Consider appropriateness of assessment endpoints e.g. to other endpoints; to receptor
	Form work/analysis plan	Factors affecting fate and transport of stressor e.g. physical; chemical; atmospheric; biotic Data considerations/requirements e.g. gaps; collection; synthesis; analysis
Effects assessment	Analyse the stressor-response relationship	Determine the dose received e.g. duration, intensity Examine assessment endpoints e.g. development; behaviour; survival; fecundity; abundance
	Create stressor-response (effects) profile(s) e.g. single-point; distribution	
Exposure assessment	Collect data/information relating to:	The stressor e.g. composition; distribution; release
		The fate and transport of the stressor (i.e. pathways) e.g. biological; chemical; physical; receiving media
		The receptor e.g. composition; distribution
	Evaluate stressor-receptor contact	Co-occurrence e.g. frequency; duration; intensity

		Nature of contact e.g. ingestion; inhalation; dermal
	Create exposure profile(s) e.g. worst-case; conservative; probabilistic	
Risk characterisation	Select relevant effects/exposure profiles	
	Estimate risk e.g. single-point comparison; cumulative distribution	
	Aggregate risk e.g. combine stressor-based risk estimates; combine endpoint-based risk estimates	
	Evaluate risk	Confidence in risk estimate(s; i.e. uncertainty analysis) e.g. qualitative; semi-quantitative; quantitative
		Significance of risk estimate(s) using e.g. regulation; stakeholders; receptor recovery potential
	Communicate risk e.g. to risk professionals; to laypersons; to stakeholders; to regulators	

Table 6 Potential uncertainties for the ERA scenario of GMHP risk to Lepidoptera, organised according to the four phases of ERAs in which they will occur: problem formulation, effects assessment, exposure assessment, and risk characterisation. Analyst(s) should consider each listed potential uncertainty against all corresponding sub-locations (which are ranked according to the frequency with which they occur within the evidence base of 171 ERAs that applied WOE methods). Prioritised uncertainty management techniques are also displayed for each uncertainty sub-location, should a related uncertainty be deemed to exist. The potential level of uncertainty must be assessed by the analyst on a case-by-case basis.

Uncertainty location/ sub-location	Problem formulation	Effects assessment	Exposure assessment	Risk characterisation
Data (epistemic)	<ul style="list-style-type: none"> Factors affecting fate and transport of stressor e.g. can we get the required data? Data considerations and requirements e.g. identifying data collection, synthesis, and analysis techniques; 	<ul style="list-style-type: none"> Determine the dose received by receptor e.g. data about the duration, frequency, or intensity of dose; Examine assessment endpoints e.g. data about receptor development, behaviour, survival, fecundity, abundance; Create exposure profile(s) e.g. distributions (of stressor intensity Vs. response magnitude) using analysed data 	<ul style="list-style-type: none"> Stressor info e.g. data about its composition, distribution, or release; Fate/transport info e.g. data about the dispersion or deposition of the receptor; about atmospheric, terrestrial, or biotic conditions; Receptor info e.g. data about dietary, breeding, migratory, or predatory patterns; Create exposure profile(s) e.g. using direct monitoring data; 	<ul style="list-style-type: none"> Assessing the significance of the risk e.g. using data regarding regulatory-enforced or stakeholder-derived acceptability levels;

Reliability (1) Monte Carlo Simulation; Sensitivity analysis; Uncertainty factors; Availability (2) Monte Carlo Simulation; Sensitivity analysis; Uncertainty factors; Precision (=16) Fuzzy logic; Expert elicitation; Monte Carlo Simulation;				
Variability (aleatory)	<ul style="list-style-type: none"> Factors affecting fate and transport of stressor e.g. variability in/between identified processes; 	<ul style="list-style-type: none"> Determine the dose received by receptor e.g. variability in the duration, frequency, or intensity of dose; Examine assessment endpoints e.g. variability in receptor development, behaviour, survival, fecundity, abundance; Create exposure profile(s) e.g. variability in single point (e.g. LC₅₀, EC₅₀) estimates; 	<ul style="list-style-type: none"> Stressor info e.g. variability in spatial/temporal distribution; variability in intensity or quantity of release; Fate/transport info e.g. variability in dispersion or deposition of the receptor; variability in atmospheric, terrestrial, or biotic conditions; Receptor info e.g. variability in dietary, breeding, migratory, or predatory patterns; Stressor-receptor contact e.g. variability in spatial, temporal or intensity of overlap; 	<ul style="list-style-type: none"> Risk estimation e.g. variability in single-point comparisons of PEC Vs. LC₅₀/EC₅₀; variability in cumulative distributions of stressor intensity Vs. response magnitude; Assessing the significance of the risk e.g. variability in regulatory-enforced or stakeholder-derived acceptability levels; variability in receptor recovery potential;
Natural (3) Monte Carlo Simulation; Further data collection; Uncertainty factors; Human (=14) Bayesian belief networks; Expert elicitation; Sensitivity analysis;				
Extrapolation (aleatory)	<ul style="list-style-type: none"> Consider appropriateness of assessment endpoints e.g. extrapolating generic endpoints for use with this receptor; 	<ul style="list-style-type: none"> Determine the dose received by receptor e.g. extrapolating knowledge for the duration, frequency, or intensity of dose; 	<ul style="list-style-type: none"> Stressor info e.g. forced extrapolation of release intensity or quantity information; Create exposure profile(s) e.g. 	<ul style="list-style-type: none"> Risk estimation e.g. extrapolating from single-point comparisons of PEC Vs. LC₅₀/EC₅₀; extrapolating from cumulative

			extrapolating to create conservative estimates; extrapolating to create worst-case estimates;	distributions of stressor intensity Vs. response magnitude; <ul style="list-style-type: none"> Assessing the significance of the risk e.g. extrapolating from regulatory-enforced or stakeholder-derived acceptability levels;
Interspecies (4)	Uncertainty factors; Monte Carlo Simulation; Further data collection;			
Spatial (=5)	Interpolation; Monte Carlo Simulation; Uncertainty factors;			
Intraspecies (7)	Uncertainty factors; Monte Carlo Simulation; Further data collection;			
Temporal (=8)	Further data collection; Uncertainty factors; Monte Carlo Simulation;			
Lab. (=12)	Uncertainty factors; Further data collection; Monte Carlo Simulation;			
Quantity (=14)	Uncertainty factors; Further data collection; Monte Carlo Simulation;			
System (epistemic)	<ul style="list-style-type: none"> Define risk relationships e.g. missing a stressor, pathway, or receptor; Select assessment endpoints e.g. missing an endpoint; Consider appropriateness of assessment endpoints e.g. relevance to other endpoints; relevance to receptor; Identifying fate/transport factors e.g. are there any 			

	missing?			
Process (=5)	Uncertainty factors; Monte Carlo Simulation; Expert elicitation;			
Effect (=10)	Uncertainty factors; Expert elicitation; Monte Carlo Simulation;			
Cause (=12)	Uncertainty factors; Further data collection; Monte Carlo Simulation;			
Model (combined)		<ul style="list-style-type: none"> • Determine the dose received by receptor e.g. model parameters for the duration, frequency, or intensity of dose; • Examine assessment endpoints e.g. model parameters for receptor development, behaviour, survival, fecundity, abundance; • Create effects profiles e.g. using model output; 	<ul style="list-style-type: none"> • Stressor info e.g. model parameters for the composition, distribution, or release; • Fate/transport info e.g. model parameters for the dispersion or deposition of the receptor; • Receptor info e.g. model parameters for the dietary, breeding, migratory, or predatory patterns; • Stressor-receptor contact e.g. model parameters for the spatial, temporal or intensity of overlap; • Create exposure profile(s) e.g. using dispersion models; using probabilistic models; 	
Output (=8)	Sensitivity analysis; Monte Carlo Simulation; Fuzzy logic;			
Structure (=10)	Sensitivity analysis; Monte Carlo Simulation; Fuzzy logic;			
Decision				<ul style="list-style-type: none"> • Selecting relevant

(combined)				effects/exposure profiles; <ul style="list-style-type: none"> • Risk aggregation e.g. combining selected estimates to form one overall risk estimate;
Decision (=16)	Adaptive management; Multi-criteria decision analysis; Bayesian belief networks;			
Language (epistemic)	<ul style="list-style-type: none"> • Defining the scope of the ERA e.g. communicating with stakeholders 			<ul style="list-style-type: none"> • Assessing the significance of the risk e.g. with regulators or stakeholders; • Communicating the risk e.g. to risk professionals; to laypersons; to stakeholders; to regulators;
Ambiguity (=16)	Fuzzy logic; Fuzzy-stochastic system;			
Vagueness (=16)	Fuzzy logic; Fuzzy-stochastic system;			
Underspec. (20)	Fuzzy logic; Fuzzy-stochastic system;			

Figure Captions

Figure 1 Overview of the clustering process applied to the uncertainty data extracted from the collected evidence base (n=171 ERAs that applied WOE methods), showing: a) all 36 recorded location-based uncertainty types; b) all 36 recorded location-based uncertainty types organised according to their nature; and c) final 20 location-based uncertainty types organised according to their nature. The superscript Greek letters in b) are matched to the superscript Greek letters in c), representing clustering into like groups. For example, model structure, model parameters, computer software/hardware, model calibration, and model simplification uncertainties, denoted by the Greek letter Kappa (κ), in b) are clustered into model structure uncertainty, also denoted by κ , in c).

Figure 2 Model showing the occurrence frequencies of the individual location-based uncertainty types identified (light grey squares; n=20), management techniques utilised (dark grey circles; n=10), and the relationships between them (black lines) within the collected evidence base (n=171 ERAs that applied WOE methods). The areas of the squares and circles (which depict the respective occurrence frequencies) are relative to each other, as are the widths of the dependency lines, where an increasing (square or circle) area and (line) width indicates an increasing frequency.

Figure 3 Matrix plot showing the correlation values (ρ) between the uncertainties and their respective management techniques within the collected evidence base (n=171 ERAs that applied WOE methods), where a higher value indicates a stronger correlation.



DSPACE

<https://dspace.org/>

Identifying uncertainty in environmental risk assessments: the development of a novel typology and its implications for risk characterisation

Skinner, Daniel J. C.; Rocks, Sophie A.; Pollard, Simon J. T.; Drew, Gillian H.

2014-12-31T00:00:00Z

Taylor & Francis

<http://dx.doi.org/10.1080/10807039.2013.779899>

Daniel J. C. Skinner, Sophie A. Rocks, Simon J. T. Pollard & Gillian H. Drew, Identifying uncertainty in environmental risk assessments: the development of a novel typology and its implications for risk characterisation, Human and Ecological Risk Assessment: An International Journal, Volume 20, Issue 3, 2014, Pages 607-640.