

Supplementary materials

The title: Freshwater organic micropollutants alter submerged macrophyte litter decomposition dynamics and microbial community in the surface sediments

Author's/Co-authors: Fan Wu^{1, 2}, Ziyi Yang¹, Tao Lyu^{2*}, Xiaohong Zhou^{1*}, Qingjie Xie¹

Affiliation (s)

1 School of the Environment and Safety Engineering, Jiangsu University, Zhenjiang 212013, China.

2 Faculty of Engineering and Applied Sciences, Cranfield University, Bedfordshire MK43 0AL, UK.

* Corresponding author: Xiaohong Zhou, Tao Lyu

E-mail: xhzhou0214@ujs.edu.cn, T.Lyu@cranfield.ac.uk

Contents of this file

Number of pages: 15

Number of tables: 3 (Tables S1 to S3)

Number of Figures: 12 (Figures S1 to S12)

Table S1 Summarized results of physicochemical parameters in freshwater sediment

	NH₄⁺-N	NO₃⁻-N	O.M.	TP	TC	TN	pH
	mg·kg ⁻¹	mg·kg ⁻¹	%	g·kg ⁻¹	g·kg ⁻¹	g·kg ⁻¹	
Initial sediment	0.08±0.00	1.58±0.11	0.13±0.03	0.25±0.00	7.95±0.11	1.07±0.00	7.22±0.27

Table S2 P-value of analysis of similarities (ANOSIM) and permutational multivariate analysis of variance (PERMANOVA) test results at the OTU level

Group	ANOSIM	PERMANOVA
CK and LD group	0.001**	0.001**
CK and ALD group	0.001**	0.001**
LD and ALD group	0.861	0.462
N, D and ND-ALD group	0.978	0.938
CK Time group	0.267	0.155
LD Time group	0.001**	0.004**
N-ALD time group	0.001**	0.007**
D-ALD time group	0.001**	0.006**
ND-ALD time group	0.001**	0.004**

(Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD). * and ** indicate significant differences ($P<0.05$ and $P<0.01$))

Table S3 Mantel test results in both the microbial community matrix (at the OTU level) and influence factor matrix based on Bray-curtis distance

Matrix1 (Species)	Matrix2	Mantel <i>r</i> statistic	<i>P</i> value
CK group	Physicochemical parameters of the sediment	-0.20618	0.339
LD group	Litter-related factors	0.74982	0.001**
	Physicochemical parameters of the sediment	0.73302	0.001**
	Litter-related factors	0.77288	0.001**
ALD group	Physicochemical parameters of the sediment	0.58788	0.001**
	Antibiotic concentration	0.61103	0.001**
	Litter-related factors	0.77367	0.001**
N_ALD group	Physicochemical parameters of the sediment	0.55843	0.002**
	Antibiotic concentration	0.76642	0.001**
	Litter-related factors	0.91055	0.001**
D_ALD group	Physicochemical parameters of the sediment	0.70549	0.001**
	Antibiotic concentration	0.88108	0.001**
	Litter-related factors	0.66212	0.001**
ND_ALD group	Physicochemical parameters of the sediment	0.6567	0.001**
	Antibiotic concentration	0.64786	0.001**

(Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD). * and ** indicate significant differences ($P<0.05$ and $P<0.01$))




Fig. S1 Shannon curves of bacterial microorganisms in freshwater sediment. (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD). * and ** indicate significant differences ($P<0.05$ and $P<0.01$))




Fig. S2 Physicochemical parameters (TC (a), TN (b), TP (c), O.M. (d), $\text{NH}_4^+\text{-N}$ (e), $\text{NO}_3^-\text{-N}$ (f) and pH (g)) of sediment during the litter decomposition processing. (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD). Means with a common letter are significantly different ($P < 0.05$))




Fig. S3 Kruskal-Wallis H test bar plot of alpha diversity indices of N-ALD, D-ALD and ND-ALD group in sediment. (Norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD)). * and ** indicate significant differences ($P < 0.05$ and $P < 0.01$)




Fig. S4 Kruskal-Wallis H test bar plot of alpha diversity indices of CK, LD, N-ALD, D-ALD and ND-ALD group 1, 35 and 110 of litter decomposition. (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD)).

* and ** indicate significant differences ($P < 0.05$ and $P < 0.01$)




Fig. S5 Principal co-ordinate analysis (PCoA) of the bacterial community structure of CK (a), LD (b), ALD (c), N-ALD (d), D-ALD (e) and ND-ALD (f) group on day 1, 35 and 110 of litter decomposition. (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD))




Fig. S6 Venn diagram of CK (a), LD (b), ALD (c), N-ALD (d), D-ALD (e) and ND-ALD (f) and group on day 1, 35 and 110 of litter decomposition. (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD).)




Fig.S7 The relative abundance of bacterial communities at Class (a-b) and genus (c-d) level. (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD))




Fig.S8 Kruskal-wallis H test bar plot of community composition between CK, LD and ALD group (a), and N-ALD, D-ALD and ND-ALD group (b) and day 1, 35 and 110 (c) at phylum level (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD)). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$)




Fig.S9 Bacterial functional pathways showing significant differences at Level 2 (a) by PICRUST2 analysis based on the KEGG database in the N-ALD, D-ALD and ND-ALD group and ecologically relevant functional groups of bacteria showing significant differences via FAPROTAX analysis (b-d) (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD)). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$)




Fig S10 Spearman correlation heatmap of Phylum species with litters nutrients and the physicochemical parameters of sediment in the CK (a), LD (b) and ALD (c) groups. (Control group (CK), litter decomposition group (LD), and antibiotic-litter decomposition group (ALD)). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$)




Fig S11 Correlation between sediment physicochemical properties (a-c), litter nutrient (d-f) and antibiotics concentration (g-i) on sediment community structure, microbial alpha diversity indices and phylum species. ((Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD)). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$)




Fig S12 Physicochemical indicators DO of overlying water. (Control group (CK), litter decomposition group (LD), antibiotic-litter decomposition group (ALD), norfloxacin-litter decomposition group (N-ALD), doxycycline-litter decomposition group (D-ALD), a combination of both antibiotics-litter decomposition group (ND-ALD))



DSPACE

<https://dspace.org/>

Freshwater organic micropollutants alter submerged macrophyte litter decomposition dynamics and microbial community in the surface sediments

Wu, Fan; Yang, Ziyi; Lyu, Tao; Zhou, Xiaohong; Xie, Qingjie

2026-06-06

Springer

<https://doi.org/10.1007/s11368-026-04421-7>

Wu F, Yang Z, Lyu T, et al., (2026) Freshwater organic micropollutants alter submerged macrophyte litter decomposition dynamics and microbial community in the surface sediments. *Journal of Soils and Sediments*, Volume 26, Issue 6, June 2026, Article number 217