

Understanding the biochemical characteristics of struvite bio-mineralising microorganisms and their future in nutrient recovery

Yirong Leng, Robert Colston, Ana Soares*

Cranfield Water Science Institute, Cranfield University, Bedfordshire, MK43 0AL, UK

Abstract

The biochemical properties of selected microorganisms (*Bacillus pumilus*, *Brevibacterium antiquum*, *Myxococcus xanthus*, *Halobacterium salinarum* and *Idiomarina loihiensis*), known for their ability to produce struvite through biomineralisation, were investigated. All five microorganisms grew at mesophilic temperature ranges (22–34°C), produced urease (except *I. loihiensis*) and used bovine serum albumin as a carbon source. *I. loihiensis* was characterised as a facultative anaerobe able to use O₂ and NO₃ as an electron acceptor. A growth rate of 0.15 1/h was estimated for *I. loihiensis* at pH 8.0 and NaCl 3.5% w/v. The growth rates for the other microorganisms tested were 0.14–0.43 1/h at pH 7–7.3 and NaCl ≤1% w/v. All the microorganisms produced struvite, as identified by morphological and X-ray Powder Diffraction (XRD) analysis, under aerobic conditions. The biological struvite yield was between 1.5–1.7 g/L of media, the ortho-phosphate removal and recovery were 55–76% and 46–54%, respectively, the Mg²⁺ removal and recovery was 92–98% and 83–95%,

* Corresponding author at Cranfield Water Science Institute, Cranfield University, Vincent Building, Cranfield, Bedfordshire, MK43 0AL, UK. Tel.: +44 (0) 1234 758121. E-mail address: a.soares@cranfield.ac.uk.

respectively. Large crystals (>300 µm) were observed, with coffin-lid and long-bar shapes being the dominant morphology of biological struvite crystals. The characterisation of the biochemical properties of the studied microorganisms is critical for reactor and process design, as well as operational conditions, to promote phosphorus recovery from waste streams.

Keywords: biomineral formation; struvite; biochemical properties; phosphorus recovery; statistical design

1 Introduction

Biological struvite (bio-struvite) has been identified as a route to recover phosphorus (P) from municipal wastewater streams (Soares et al., 2014). Microorganisms play an important role in struvite bio-mineralisation through different metabolic activities (Sinha et al., 2014) and by precipitation of specific structures or substances for microbial processes (Arias et al., 2017). Five microbial strains, *Halobacterium salinarum*, *Bacillus pumilus*, *Brevibacterium antiquum*, *Myxococcus xanthus*, and *Idiomarina loihiensis*, have been reported to be involved in biologically driven struvite formation in liquid streams (Table 1, González-Muñoz et al., 2008; Soares et al., 2014). *M. xanthus*, *I. loihiensis* and *H. salinarum* were reported to produce extracellular polymeric substances (EPS), which may fix cations and contribute to mineral heterogeneous nucleation and precipitation (González-Muñoz et al., 2010, 2008; Merroun et al., 2003). Most of the selected microorganisms can use O₂ as an electron acceptor (Table 1). *H. salinarum* has been reported to be able to use dimethyl sulfoxide

(DMSO) as an electron acceptor under anaerobic conditions, and use photophosphorylation in the presence of light (Table 1).

B. pumilus and *M. xanthus* can use carbohydrates as a carbon source but this does not apply to *B. antiquum* and *H. salinarum*. According to the literature, all the selected microorganisms can use protein/amino acids as a carbon source (Robinson, 2014; Trujillo and Goodfellow, 2015). The utilisation of organic carbon sources depended on enzyme production, and the rates of enzyme-catalysed reactions optimally performed under appropriate temperature, pH and salinity ranges (Silva et al., 2016). The selected microbial strains have been reported to grow in pHs from 5.5 to 9, and temperatures ranging from 20–45 °C (Table 1). The halotolerant microorganisms *B. antiquum*, *H. salinarum* and *I. loihiensis* can live in environments containing high NaCl (Gavrish et al., 2004; González-Muñoz et al., 2008; Mesbah and Wiegel, 2005), particularly *H. salinarum*, which can survive at extremely high NaCl concentrations (17.4~30.16 %, Table 1).

Although some of the biochemical properties and growth conditions of selected microorganism have been reported in the literature, some of the values are controversial and further verification and characterisation is required. Statistical experimental design is recognised as an approach widely used for parameter screening in optimisation studies (Massey et al., 2009). By using such design, Simoes et al. (2017) investigated the significant factors required for *B. antiquum* growth, and maximised the growth rates in wastewater streams by screening and optimising a number of factors.

This study aims to investigate the biochemical properties of the selected microorganisms owing to their capability to produce struvite through biomineralisation. For industrial exploitation of microorganisms, the investigation of biochemical characterisation is critical for appropriate processes design and meeting microbial requirements by optimising reactor operational conditions. The temperature, pH, electron acceptor, and organic carbon source are among the most important environmental parameters affecting microbial growth and organic substance synthesis (Silva et al., 2016). Knowledge of such parameters will allow the design of reactors/processes and operational conditions to ensure proliferation of the selected microorganisms, and even out-compete other microorganisms in mixed cultures, for eventual enhanced P recovery by struvite from waste streams.

73 **Table 1 Biochemical properties of the five tested microorganisms**

		<i>B. pumilus</i>	<i>M. xanthus</i>	<i>B. antiquum</i>	<i>I. loihiensis</i>	<i>H. salinarum</i>
Strain		MTCC 1640	CECT 422	DSM 21545	MAH1 /CECT 5996	DSM 671
Type		Bacteria	Bacteria	Bacteria	Bacteria	Archaea
Gram reaction		+	-	+	-	-
Cell shape		Rod	Rod	Short rod/ coccoid	Rod	Rod
Size		0.6~0.7 x 2.0~3.0 µm	0.5 x 6 µm	0.6~1 µm	0.3~0.5 x 0.6~2 µm	0.5-1 x 1~6 µm
Motility		+	+	-	+	+
Endospore forming		+	+	-	-	-
O ₂ requirement/tolerance		Aerobic	Aerobic	Aerobic	Aerobic	Facultative anaerobic, photophosphorylation at low O ₂ concentration with light
Electron acceptor		O ₂	O ₂	O ₂	O ₂	O ₂ , dimethyl sulfoxide
Extracellular polymeric substances synthesis		Not documented	+	Not documented	+	+
Preferred organic carbon source	carbohydrate	Arabinose, mannitol, xylose, glucose, lactose, acetone	Glucose	Not able to directly use	Not documented	Not able to directly use
	protein/amino acid	Casein, lysine,	Amino acids	Casein, amino acid	Amino acid	Lysine, ornithine, arginine
	Other	Citrate, sucrose, D-trehalose, starch, D-glucose, D-arabinose, D-xylose, gelatin	Not documented	Gluconate, urea, gelatin, salicin, sorbitol	L-alaninamide	Gelatin
Growth temperature		20~40 °C, optimum 30~35 °C	14~40°C, optimum 34~36°C	7°C, <37°C; optimum 24~26°C	2 ~ 43°C; optimum 28 ~37 °C	20~55°C, optimum 35~50°C
Growth pH		6~8, optimum at 7	5.5~9.0, optimum at 7	5.5~10, optimum at 7	not documented	5.5~8, optimum at 7
Growth in NaCl		0~2 %	not documented	0~18 %, optimum 3%	0.7~20 %, optimum 2~6 %	17.4~30.16 %, optimum 20.3 %
References		(Robinson, 2014; Shivaji et al., 2006)	(González-Muñoz et al., 2010; Janssen et al., 1977; Merroun et al., 2003; Poza et al., 2004; Robinson, 2014)	(Gavrish et al., 2004; Robinson, 2014; Simoes et al., 2017; Trujillo and Goodfellow, 2015)	(González-Muñoz et al., 2008)	(Losensky et al., 2017; Mesbah and Wiegel, 2005; Mormile et al., 2003; Zinder and Dworkin, 2013)

2 Material and methods

2.1 Microbial strains and culture solution

Five microbial strains were used in this study: *H. salinarum* & *B. antiquum* (DSM 671 & DSM 21545, German Resource Centre for Biological Material, Germany), *B. pumilus* (GB43, LGC Standards, Middlesex, UK), *M. xanthus* & *I. loihiensis* (CECT 422 & MAH1 /CECT 5996, Spanish Type Culture Collection, University of Valencia, Paterna, Spain). The microorganism were grown in synthetic B41 solution comprising 4 g/L yeast extract, 2 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and 2 g/L K_2HPO_4 (Da Silva et al., 2000). The solution was autoclaved at 121°C for 20 minutes and cooled to room temperature (20–22°C). For inoculation of each microbial strain; 100 ml synthetic B41 solution in a 250 ml E-flask was inoculated with 0.9% w/v NaCl pre-washed pure cultures of the selected microorganisms that were grown for 96 hours. The E-flasks were sealed with foam stoppers and incubated on an orbital shaker (Stuart model SSL1, Fisher Scientific, UK) at 150 rpm at room temperature. The halophile *I. loihiensis* was grown in B41 solution with 1% w/v NaCl (González-Muñoz et al., 2008).

2.2 Gram staining and enzyme production

Microorganisms, in their early exponential phase of growth (0-8 hours), were Gram-stained using standard methods (Claus, 1992). A KB002™ HiAssorted Biochemical Test Kit (HiMedia Laboratories Pvt. Ltd, India) was used to characterise the pure cultures, according to the manufacturer's instructions. All tests were completed in triplicate and a non-inoculated control was maintained under identical conditions.

95 2.3 Statistical design of experiments

96 To investigate the impact of growth conditions on microorganisms, a full factorial
97 experiment (FFD) was designed with five factors: temperature, initial pH, NaCl, Ca²⁺,
98 (by CaCl₂) and bovine serum albumin (BSA) as an additional carbon source (Table S1).
99 As factors that are key in optimising the industrial processes involving microbes (Silva
100 et al., 2016), additional variations in NaCl and Ca²⁺ were carried out as they have been
101 inferred to be detrimental to bacterial function and abiotic struvite growth (Le Corre et
102 al., 2005; Rivadeneyra et al., 2006). The tests were based upon low, medium and high
103 levels in relation to characterisation of municipal wastewater and sludge dewatering
104 liquors (Table S1). Temperatures varying from 6–34°C and pH 5.5–8.5 to cover the
105 range of temperatures and pH of municipal wastewater (Tchobanoglous et al., 2003). Ca
106 concentration was adjusted to 28 mg/L (Gassama et al., 2015). The NaCl content varied
107 between 0.5–3.5% w/v, based on characterisation of municipal wastewater and sludge
108 dewatering liquors in different full scale sites in the UK (Simoes et al 2017). (Table
109 S1). Three factors (temperature, NaCl and initial pH) at the 3-level and two factors (Ca²⁺
110 and BSA) at the 2-level corresponded to $3^3 \times 2^2$ combinations of recipes, which were
111 studied in duplicate and thus generated $3^3 \times 2^2 \times 2 = 216$ tests for each microorganism.
112 The initial and final intact cell counts were examined to generate the overall cell
113 increase that was used as a response to the factors investigated. The experimental data
114 were fitted to a first-order linear regression model or second-order polynomial
115 regression model considering linear and quadratic forms of the independent factors. The
116 response surface methodology (RSM, (Bezerra et al., 2008)) was applied to examine the
117 significant relationship ($p < 0.01$) between cell increase and the five growth factors, as

well as the significant two-factor interactions ($p < 0.01$). The RSM was also used to determine the optimal conditions that jointly maximise the cell increase by applying a multiple response optimisation. All statistical design and analysis was performed using Minitab 17 (Minitab, 2010).

2.4 Microbial cultivation under investigated growth conditions

Microorganisms were grown in 96-well sterile microplates with working volume about 250 μ l per well (Corning™, Fisher scientific, UK). Each well contained 234 μ l solution and 26 μ l inoculum. To prepare the solutions corresponding to the FFD recipes (Table S1), synthetic B41 solution with different NaCl concentrations was autoclaved and mixed with 0.22 μ m sterile filtered (Sartorius Stedim Biotech, Germany) BSA and CaCl₂ concentrated solutions. The initial pH was adjusted by 0.1 M NaOH and 0.1 M HCl sterile solutions. To minimise liquid evaporation from each well, only the central wells (10 x 6) of the microplate were used for microbial inoculation, and the edge and corner wells of the microplate were used for the non-inoculated controls (Syberg, 2016). Breathable rayon film (VWR Collection, VWR, UK) was used to seal the microplates to stop cross-contamination and to achieve uniform air and gas exchange, while also reducing liquid evaporation for each well. The sealed microplates were then placed inside a cube humidity chamber with four ventilation holes at each bottom corner and with a water reservoir inside. The humidity chamber was kept at constant temperatures of 6, 20 and 34 °C, incubated for 106, 66 and 48 hours, respectively. The application of a humidity chamber was found to reduce liquid evaporation from 150 to 20–25 μ l/well by the end of the incubation period.

2.5 Microbial cultivation at different dissolved oxygen levels

After investigating microbial growth with the different FFD recipes, the conditions that resulted in the highest increase of intact cell count were repeated but this time, incubation took at two dissolved oxygen (DO) levels using sacrificial glass vials. For microbial cultivation under aerobic conditions, 30 ml sacrificial glass vials containing 9 ml synthetic B41 solution pre-adjusted in terms of pH, NaCl, Ca^{2+} and BSA were inoculated with 1 ml inoculum and sealed with breathable film. The DO in the 30 mL vials varied between 6–8 mg/L.

For microbial cultivation under anoxic/anaerobic conditions, synthetic B41 solution was pre-adjusted in terms of pH, NaCl and bubbled with $\text{N}_{2(\text{g})}$ after passing through a 0.22 μm sterile filter (Sartorius Stedim Biotech, Germany) at a rate of 30 L/min. The 10 ml sacrificial glass vials were sealed with a nontoxic butyl rubber stopper and autoclaved (121°C for 20 min). The DO in the 10 ml vials was close to 0 mg/L. Concentrated solution of CaCl_2 and BSA, and 1 ml inoculum were then added using a sterile disposable syringe and needle (VWR, UK). The capability of *I. loihiensis* to grow under anoxic conditions was examined in synthetic solutions with absence of O_2 but with 0.5 g/L NaNO_3 .

The glass vials were placed inside humidity chambers and incubated on an orbital shaker-incubator (MAXQ5000 M6, Thermo Scientific, UK) at 150 rpm for 120 hours. Samples were taken for examination at regular intervals (4–24 hours) through sacrificial vials. All tests were completed in triplicate, and non-inoculated controls were maintained under identical conditions.

2.6 Abiotic struvite formation

Abiotic struvite was prepared by mixing 200 mL 0.05 M $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ with 100 ml 0.2 M $\text{NH}_4\text{H}_2\text{PO}_4$, both pre-adjusted to pH 9 with 1 M NaOH (Le Corre et al., 2005). Concentrated BSA solution was added to the mixture at a rate of 4 g/L and incubated on an orbital shaker at 150 rpm at room temperature for 24 hours.

2.7 Crystal isolation, purification and determination

The microorganisms were inoculated in 500 mL B41 media in sterile, 1 L Duran bottles, sealed by a breathable film, and incubated under the investigated optimal growth conditions at agitation rate 150 rpm for 120 hours. At the end of the incubation period, the samples were filtered through a 10 μm nylon-mesh filter (Plastok, UK) and the crystals were washed with deionised water twice. The isolated crystals were air-dried at 37°C for 2 hours and weighed to determine crystal yields. The pure crystals were then identified by X-ray powder diffractometer (XRD, D5000, Siemens / Bruker, Germany).

2.8 Analytical methods

The intact cell count was examined by the SYBR Green I - propidium iodide co-staining method using a flow cytometer (BD accuri C6, BD Biosciences, US, (Nocker et al., 2017)). Solution DO and pH values were determined with a portable DO- meter (HQ40D, HACH, UK) and digital pH-meter (Jenway 3540, Bibby Scientific, UK). The concentrations of soluble chemical oxygen demand (SCOD), $\text{PO}_4\text{-P}$, $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}_2$ were monitored with Merck Spectroquant® test kits. Mg^{2+} was measured by atomic absorption spectroscopy (AAS, Analyst 800, PerkinElmer, UK) equipped with flaming

and electrothermal spectrometers. A high-resolution microscope (L-series upright compound microscope, Division of GT vision Ltd, UK) was applied for observation of Gram-stained cultures and crystal morphology in microbial cultures.

3 Results and discussion

3.1 Microbial properties and enzyme production

B. pumilus and *B. antiquum* were identified as Gram-positive and *M. xanthus*, *H. salinarum* and *I. loihiensis* as Gram-negative, which agrees with previously published information (Table 1). In particular, *B. pumilus* formed crusted two-cell clusters or tetrads in B41 media, which were not observed in the other four microbial cultures. Such cell structures did not grow in size but had the potential to aggregate together or onto the crystal surface. Similar cell structures were observed as mineralised *Thiomargarita* embryo-infesting cells (Bailey et al., 2007), and as silica spheroids onto the cell sheath in microbial silicification (Yee et al., 2003). Thus, the crusted cell structures observed during *B. pumilus* growth in this study is proposed associated with the bio-mineralisation.

It is suggested that *B. pumilus* can form mineral particles along the cell surface during exponential phase in solutions rich in $\text{PO}_4\text{-P}$ and Mg^{2+} , and the mineral particles firmly attach to the cell surface to form completely encrusted cell minerals. Based on previous work on microbial mineralization occurring at the peptidoglycan wall due to negative charges associated in gram-negative and gram-positive, concentrating cations such as Mg^{2+} (Orange et al., 2009) and the consequential interaction with secreted phosphate

204 group and carboxyl groups which bind into the peptidoglycan framework of gram-
205 positive bacteria (Schultze-Lam et al., 1996).

206 The biochemical characterisation tests demonstrated varied enzyme production amongst
207 the microorganisms investigated. Nevertheless, it was quite remarkable to observe that
208 *H. salinarum*, *B. antiquum*, *B. pumilus* and *M. xanthus* were capable of using ornithine
209 as a carbon source and produced urease. Urease activity, as well as the degradation of
210 proteins, can generate energy for microbial growth and produce NH_3 as a by-product,
211 which raises the pH and release $\text{NH}_4\text{-N}$ to combine with $\text{PO}_4\text{-P}$ and Mg^{2+} for struvite
212 precipitation (Sadowski et al., 2014). Bio-mineralisation of struvite by urease-producing
213 microorganisms in the urinary tract has been reported, leading to the formation of
214 kidney stones, that typically contain 15–20% struvite (Arias et al., 2017; Coe et al.,
215 2005; Prywer and Torzewska, 2010).

216 *I. loihiensis* was the only microorganism investigated in this study that did not produce
217 urease and also the only one that showed a positive reaction of NO_3^- reduction to NO_2^- .
218 The latter is a common phenomenon in anoxic respiration, where NO_3^- was used as an
219 electron acceptor.

220 All five microorganisms showed neither positive nor negative results in terms of lysine
221 utilisation. They were also found to be negative for their ability to use citrate and
222 carbohydrates (including glucose, arabinose, lactose, adonitol and sorbitol) as a carbon
223 source, and for phenylalanine deamination and hydrogen sulfide production. The only
224 exception was that *B. pumilus* showed a 33% positive result for glucose utilisation. The

results obtained in this study partially agree with the organic carbon source utilisation presented in Table 1.

3.2 Identification of significant factors to microbial growth

By applying the multi-response surface methodology, each microorganism was grown in optimum conditions (Table 2) within the range of chemical conditions of wastewaters and sludge dewatering liquors (Table S1). BSA was identified to have a significant positive linear correlation ($p < 0.01$) with microbial growth. All selected microorganisms were able to use BSA as a carbon source (Table 2). Temperature, pH and NaCl were also identified as being significant for microbial growth for all selected microorganisms. A Ca^{2+} of 28 mg/L was identified to be required for growth of *I. loihiensis* but not for *M. xanthus* growth, and was a non-significant factor for the other three microorganisms (Table 2). In addition, temperature correlated with other factors (carbon source and p , $p < 0.01$) within the investigated range of 6–34°C. The growth of *B. pumilus*, *M. xanthus*, and *I. loihiensis* had a positive linear correlation with the temperature and reached a peak value at 34°C, while the relationship between temperature and cell count for *H. salinarum* and *B. antiquum* fitted a quadratic trend and the growth peak occurred between 22–24°C. Thus, the optimal growth temperature and enzyme activity for the investigated microorganisms was within the mesophilic range of temperatures (Table 2). Quadratic relationships between pH and microbial growth were also observed. *B. pumilus*, *H. salinarum*, *B. antiquum* and *M. xanthus* preferred neutral pH (7.1–7.3), while *I. loihiensis* was observed to adapt to a mild alkaline pH of 8.0 (Table 2). Furthermore, *I. loihiensis*, as a halophile, distinguished itself from the other four microorganisms by its ability to adapt to grow at high NaCl concentration (3.5% w/v),

highlighting its ability to control the increased osmotic pressure due to higher salt concentrations (Robinson, 2014). Whereas the other four microorganisms preferred a reduced NaCl concentration (0.5–1% w/v, Table 2). A coefficient of determination (r^2) was introduced to display the degree that the regression model approximates the real data points, with an $r^2 > 0.7$ typically being considered good (Grace-Martin, 2012). In this study, the coefficient of determination was within the range of 0.71–0.94 (Table 2), and thus the regression model could well explain the divergence of data points from a trend.

Table 2 Significant growth factors (main effect, $p < 0.01$) and preferred growing conditions defined by multi-response surface methodology

	Temperature (°C)	NaCl (% w/v)	pH	BSA (g/L)	Ca ²⁺ (mg/L)	r^2
<i>B. pumilus</i>	34	0.5	7.3	4	Ns	0.94
<i>H. salinarum</i>	24	0.5	7.1	4	Ns	0.80
<i>B. antiquum</i>	22	0.5	7.3	4	Ns	0.85
<i>M. xanthus</i>	34	1	7.2	4	0	0.73
<i>I. loihiensis</i>	34	3.5	8.0	4	28	0.71

Ns - Non-significant correlation to microbial growth

$a - r^2$, ranging from 0 to 1, indicated the proportion of variation that can be explained by the regression model. $r^2 = 1$ indicates that the regression line perfectly fits the data.

3.3 Microbial growth at different dissolved oxygen levels

No lag phase of microbial growth was observed under aerobic conditions (DO = 6–8 mg/L) and the exponential phase occurred within 24/48 hours of incubation starting. The growth rates (μ) for the different microorganisms varied between 0.14 and 0.43

1/hour (Figure 1a). The relatively high growth rate of *B. pumilus* (0.35 1/hour) and *M. xanthus* (0.24 1/hour) under anaerobic conditions distinguished themselves from the other three microbial strains ($\mu \leq 0.04$ 1/hour, Figure 1a). The growth rate of *I. loihiensis* under anoxic condition was 0.12 1/hour (Figure 1a), and >99.5% of NO₃-N was reduced by the end of the incubation time. The final microbial intact cell counts for *B. pumilus*, *B. antiquum*, *M. xanthus*, *H. salinarum*, *I. loihiensis* were 80–94% lower under anaerobic conditions and 66% lower under anoxic conditions, than those under aerobic conditions (Figure 1b). The SCOD removal was 20–27% under aerobic conditions, 0–2.4% under anaerobic conditions. SCOD removal by *I. loihiensis* under anoxic conditions was only 6% (Figure 1c). Aerobic respiration, using O₂ as an electron acceptor, is known to enable microorganisms to convert energy from carbon sources to adenosine triphosphate production more efficiently than using other electron acceptors (Kader & Saltveit, 2003). Hence, it was unsurprising that higher cell counts and SCOD removal were observed under aerobic conditions (Figure 1b-c). None of the intact cell microbial growth or SCOD removal was observed in non-inoculated controls.

I. loihiensis has been reported to be an aerobic organism (González-Muñoz et al., 2008). However, in this study it was identified as a facultative anaerobe, able to use both O₂ and NO₃ as an electron acceptor. Although *B. pumilus* and *M. xanthus* have been recognised as obligate aerobes (Robinson, 2014), in this study they were found to be facultative anaerobes. There was no report related to *M. xanthus* being a facultative anaerobe, although genome sequencing demonstrated that its common ancestor was a facultative anaerobe (Thomas et al., 2008). Several *B. pumilus* strains have been reported as facultative anaerobes, yet the electron acceptor has not been identified

289 (Alcaraz, 2015). *B. antiquum* was observed to be a strict aerobe in this study with a
290 specific growth rate of 0.14 1/hour, agreeing with previously reported growth rates in
291 wastewater with NaCl (3% w/v) and using acetate as the major carbon source
292 (equivalent to 1124 mg chemical oxygen demand/L, (Simoes et al., 2017)). Besides
293 carbon source and electron acceptor, exhaustion of macro/micro-nutrients (Maathuis,
294 2009) or formation of toxic metabolism by-products (Trinh and Srienc, 2009) cannot be
295 excluded as factors affecting the microbial growth.




Figure 1 Microbial growth rate (1/h) during exponential growth (0 - 24 /48 h) (a), and intact cell counts (b) and SCOD removal (c). After 120 hour incubation period under aerobic (□), anoxic (Δ) and anaerobic conditions (○). Error bars represent standard deviation obtained from duplicates.


300 3.4 Identification of struvite crystals

301 All the selected microorganisms produced crystals under aerobic conditions. The XRD
302 diffractions results showed that the curves of isolated purified crystal products met the
303 peak profile of the standard struvite crystals curve (pattern: COD 9007674). The
304 crystals produced by all the microorganisms tested were hence identified as struvite
305 (here called bio-struvite as it was formed through bio-mineralisation mechanism). The
306 bio-struvite and the abiotic struvite crystals presented with the same dominant faces,
307 with miller indices of [011], [111] and $[00\bar{1}]$ (Table S2). Besides these three faces,
308 [010] was also found predominant for the bio-struvite produced by *M. xanthus*, *H.*
309 *salinarum*, *B. antiquum*, and *I. loihiensis* (Table S2).

310 The dominant morphology of the bio-struvite crystals was coffin-lid shape (Figure 2a-b,
311 d-e) and long bar shape (Figure 2c), which have been reported to be among the most
312 typical struvite forms (Tansel et al., 2018). The shape and size of these bio-struvite
313 crystals were different from the relatively small dendritic abiotic struvite (Figure 2h).
314 Abiotic struvite of such dendritic X-shape is typically formed at high pH ≥ 9 (Ronteltap
315 et al., 2010; Ye et al., 2014). In most microbial cultures grown under aerobic conditions,
316 crystals were observed as early as after 4 hours of incubation (Figure 2g) and these grew
317 larger to more than 300 μm during stationary phase (Figure 2a-b, d-f). In *B. pumilus*
318 culture, a considerable number of crusted tetrad clusters were observed aggregated on
319 the specific bio-struvite crystal surface, particularly the [011] faces (Figure 2g). Bacteria
320 (e.g. *Proteus mirabilis*) was reported to exert control on the bio-struvite crystal
321 morphology (Torzewska et al., 2003). Prywer and Torzewska (2009) proposed a

potential of specific molecular interactions, which related the *P. mirabilis* capability of binding to positively charged molecules (e.g. $\text{Mg}[\text{H}_2\text{O}]_6^{2+}$ octahedra) in the crystal surface structure. Such molecular interactions varied with the composition of the microbial secreted biomolecules (e.g. polysaccharide) and its affinity for cations (Prywer and Torzewska, 2009), as well as the charged molecules' type and density on the crystal surface (Sadowski et al., 2014). In this study, the microbial growth may have potential to enhance specific faces of the bio-struvite crystals (e.g. [011], [111], $[00\bar{1}]$ miller indices) and therefore lead to the different crystal morphology (e.g. coffin-lid shape).

The self-assembly of crystals such as contact twinning (Figure 2i-j) and penetration twinning (Figure 2k) were observed, along with the parallel grouping of coffin-lid shaped crystals (Figure 2l) and long-bar shaped crystals (Figure 2m). Some bio-struvite crystals were observed with truncated apices, which was related to enhanced [111] end caps (Figure 2n). Similar struvite crystals were observed at low or moderate pH (8–8.5, (Sadowski et al., 2014)).



337 Figure 2 Coffin-lid and long-bar shaped bio-struvite produced in stationary phase by (a) -
 338 *B. pumilus*; (b, c) - *M. xanthus*; (d) - *H. salinarum*; (e) - *B. antiquum* and (f) - *I. loihiensis*.
 339 (g) Crusted cell cluster aggregated on *B. pumilus* bio-struvite crystal surface (4 h
 340 incubation); (h) - dendritic abiotic struvite crystals, bio-struvite crystals contact twinning
 341 (i-k), parallel grouping (l-m), bio-struvite crystals with truncated apices (n), Black bar
 342 scale – 88.32 μm , white bar scale – 35.93 μm , grey bar scale – 10.19 μm .

343 3.5 Removal and recovery of ortho-phosphate and magnesium

344 The bio-struvite crystal yields under aerobic conditions varied between 1521 and 1746
345 mg crystals per litre synthetic solution (Table 3). No crystal was collected under
346 oxygen-limiting conditions (Table 3). The removal of $\text{PO}_4\text{-P}$ and Mg^{2+} by the end of the
347 120 hour incubation time varied with DO levels. Under aerated conditions, the removal
348 of $\text{PO}_4\text{-P}$ and Mg^{2+} was between 55–76% and 92–98%, respectively. Under anaerobic
349 conditions, the removal of $\text{PO}_4\text{-P}$ and Mg^{2+} varied between 1–2% and 2–8% of Mg^{2+} ,
350 respectively. Under anoxic conditions, *I. loihiensis* was able to remove 2% of $\text{PO}_4\text{-P}$
351 and 32% of Mg^{2+} , from the synthetic media (Table 3).

352 A mass balance to the nutrients in solution (liquid and crystals $<10\text{ }\mu\text{m}$) demonstrated
353 that considerable amounts of $\text{PO}_4\text{-P}$ and Mg^{2+} recovered were as bio-struvite (46–54%
354 and 83–95%, respectively (Table 3). Although *B. antiquum* removed a relatively high
355 content of $\text{PO}_4\text{-P}$ (314 mg/L, 76%) from the synthetic solution, the $\text{PO}_4\text{-P}$ recovery
356 (48%) by bio-struvite crystals was lower than those for *B. pumilus*, *M. xanthus* and *H.*
357 *salinarum* (52–54%). Moreover, the Mg^{2+} recovery by *B. antiquum* (84%) and *I.*
358 *loihiensis* (83%) was observed to be lower than for the other three microorganisms (92–
359 95%).

360 **Table 3 Removal and recovery of PO₄-P and Mg²⁺ at two DO levels by the end of 120 hour**
361 **incubation period.**

	DO (mg/L)	Bio-struvite ^a production (mg bio- struvite/L synthetic solution)	PO ₄ -P removal	Mg ²⁺ removal	PO ₄ -P recovered by bio-struvite ^a	Mg ²⁺ recovered by bio-struvite ^a
<i>B. pumilus</i>	7.2	1700	265 ± 3 mg/L 64%	176 ± 1 mg/L 98%	215 mg/L 52%	167 mg/L 93%
	0	0	8 ± 3 mg/L 2%	5 ± 1 mg/L 2%	-	-
<i>M. xanthus</i>	7.2	1746	272 ± 1 mg/L 66%	177 ± 0 mg/L 98%	221 mg/L 54%	171mg/L 95%
	0	0	5 ± 1 mg/L 1%	9 ± 1 mg/L 5%	-	-
<i>H. salinarum</i>	7.8	1692	276 ± 1 mg/L 67%	170 ± 0 mg/L 94%	215 mg/L 52%	166 mg/L 92%
	0	0	7 ± 0 mg/L 2%	7 ± 1 mg/L 4%	-	-
<i>B. antiquum</i>	8.0	1550	314 ± 1 mg/L 76%	173 ± 0 mg/L 96%	196 mg/L 48%	152 mg/L 84%
	0	0	7 ± 3 mg/L 1%	10 ± 1 mg/L 6%	-	-
<i>I. loihiensis</i>	6.2	1521	229 ± 1 mg/L 55%	166 ± 0 mg/L 92%	192 mg/L 46%	149 mg/L 83%
	0	0	4 ± 2 mg/L 1%	14 ± 5 mg/L 8%	-	-
	0 ^b	0	9 ± 3 mg/L 2%	58 ± 1 mg/L 32%	-	-
control	-	0	0	0	-	-

362 a - Bio-struvite crystals >10 µm

363 b - Anoxic condition with 0.5 g/L NaNO₃

364 The synthesis of bio-struvite and removal of $\text{PO}_4\text{-P}$ and of Mg^{2+} have been reported to
365 depend on microbial growth and metabolism pathways (Sinha et al., 2014). The significant
366 difference of $\text{PO}_4\text{-P}$ removal and bio-struvite crystal yields between aerobic and anaerobic
367 conditions in this study indicates the importance of DO for P removal and bio-struvite
368 production. Furthermore, the capability of the selected microorganisms, particularly *I.*
369 *loihiensis*, to produce bio-struvite and remove $\text{PO}_4\text{-P}$ in this study might be underestimated
370 due to the NaCl concentration of 3.5% w/v. It was reported that the increased NaCl could
371 increase the solubility of the struvite phase and therefore lead to inhibition of the bio-struvite
372 crystal size (Rivadeneyra et al., 2006). Significant prevention of bio-struvite production was
373 also observed on sludge dewatering liquors with 3% w/v NaCl (Simoes et al., 2017). The
374 molar ratio of the removed $\text{PO}_4\text{-P}$ to Mg^{2+} by *B. antiquum* under aerobic conditions ($[\text{PO}_4\text{-P}]/[\text{Mg}^{2+}] = 1.4$)
375 was relatively higher than the standard stoichiometric ratio $[\text{PO}_4\text{-P}]/[\text{Mg}^{2+}]$
376 of struvite, indicating that *B. antiquum* may absorb considerable amounts of $\text{PO}_4\text{-P}$ into cells.
377 Such $\text{PO}_4\text{-P}$ accumulation within *B. antiquum* cells was reported to be relative to the
378 formation of intracellular bio-struvite (Smirnov et al., 2005).

379 *M. xanthus* displayed a higher recovery rate of bio-struvite, in comparison with the other
380 microbial strains investigated. Although it removes less P than other strains (66% compared
381 to 76% by *B. antiquum*) less of the resource was lost inside biomass or as small crystals.
382 Furthermore, *M. xanthus* presented high growth rates (0.43 1/h) and competitive SCOD
383 removal among the others tested (Figure 1). On the other side, its final intact cell count was
384 amongst the lowest, indicating it may be more susceptible to changing conditions
385 experienced in a batch reactor.

386 3.6 **Implication to the wastewater industry**

387 Similar to most biological processes in conventional wastewater treatment, bio-struvite
388 production will be ideally applied in open, mixed-culture conditions. The microorganisms
389 enrolled in bio-struvite production are required to out-compete others and become the
390 dominant species in a mixed-microbial culture. The investigation of microbial capabilities
391 and growth of the selected microorganisms in this study can help identify the suitable types
392 of streams (e.g. municipal wastewater, urine, addition of seawater to wastewater) for optimal
393 resource recovery. Streamline reactor and process design, with the most appropriate
394 operational conditions regarding temperature, pH, availability of certain nutrients, and
395 concentrations of NaCl, Ca²⁺ and DO (Table 2). By tailoring processes based on these results
396 the chance for the selected bio-struvite-producing bacteria out-competing other microbial
397 communities in wastewaters increases, whilst efficiency controlling the system can reduce
398 energy and additive costs.

399 The findings of biochemical characterisation in this study (

Table 4) can be compared with existing information (Table 1). This study's findings indicate that *B. pumilus*, *M. xanthus*, *B. antiquum* and *H. salinarum* have the potential to grow in urine due to their ability to produce urease and adapt to lower pH (urine pH 5-7), suggesting that these bacterium would be viable options for urine-separated stream treatment and resource recovery. This also has the benefit of reducing the uncertainty of these bacteria being out competed by mixed-cultures in wastewaters, as urine is typically sterile, improving decentralised system's efficiency and reliability. *I. loihiensis* has the ability to grow under anoxic conditions, alkaline pH, high concentrations of NaCl and Ca^{2+} (e.g. seawater) and can possibly be used in selective chemical pressures for competitive growth. *B. pumilus*, *M. xanthus* and *I. loihiensis* have the potential to grow in effluents from mesophilic digesters of temperature around 35 °C. Furthermore, specific wastewater streams characterised by high load of protein/amino acids (e.g. dairy processing wastewater) are proposed as preferred wastewater sources to grow the microorganisms. In all scenarios a well-aerated environment was identified as being essential for bio-struvite production, which can be achieved by pre-existing infrastructure in wastewater treatment plants as secondary treatment process are aerobic, with forced or passive aeration (Tchobanoglous et al., 2003).

With the increased knowledge in struvite recovery, researchers have also been investigating its suitability as a fertiliser. This study has shown that bio-struvite produces coffin-lid shaped, tabular crystals, whilst abiotic struvite produced was more dendritic crystal morphologies (Figure 2). The abiotic struvite produced conformed to other studies, where pH exceeded 9 (Ronteltap et al., 2010; Ye et al., 2014). The crystal morphologies of bio-struvite (Figure 2) have been demonstrated more suitable for direct land application, as the reduced surface area

422 from more euohedral crystals, improving its soil retention time as a fertiliser (Shaddel et al.,
423 2019).

424 **Table 4 Summary of biochemical properties of investigated microorganisms and comparison**
 425 **with existing literature (based on Table 1)**

	New	Agreement
Enzyme production	<ul style="list-style-type: none"> • <i>B. pumilus</i>, <i>M. xanthus</i> and <i>H. salinarum</i> produce urease 	<ul style="list-style-type: none"> • <i>B. antiquum</i> produce urease
Electron acceptor	<ul style="list-style-type: none"> • <i>I. loihiensis</i> – O₂ and NO₃-N (facultative anaerobe) • <i>B. pumilus</i> and <i>M. xanthus</i> are facultative anaerobes ^c 	<ul style="list-style-type: none"> • All the tested microorganism can use O₂ as electron acceptor ^a
Carbon source	<ul style="list-style-type: none"> • <i>I. loihiensis</i> cannot directly use carbohydrates, but can use proteins • <i>B. pumilus</i> and <i>M. xanthus</i> cannot directly use carbohydrates ^{b, c} 	<ul style="list-style-type: none"> • <i>B. antiquum</i>, <i>I. loihiensis</i> and <i>H. salinarum</i> cannot directly use carbohydrates • <i>B. pumilus</i>, <i>M. xanthus</i>, <i>H. salinarum</i> and <i>B. antiquum</i> can use amino acids/proteins
Growth temperature	<ul style="list-style-type: none"> • <i>H. salinarum</i> prefer mesophilic temperature (24°C) ^c 	<ul style="list-style-type: none"> • <i>B. pumilus</i> <i>M. xanthus</i> and <i>I. loihiensis</i> prefer high mesophilic temperature (34°C) • <i>B. antiquum</i> prefer mesophilic temperature (22°C)
Growth pH	<ul style="list-style-type: none"> • <i>I. loihiensis</i> can grow within pH 5.5–8.5, and prefer mild alkaline pH 8 	<ul style="list-style-type: none"> • <i>B. pumilus</i>, <i>M. xanthus</i>, <i>B. antiquum</i> and <i>H. salinarum</i> prefer neutral pH (7.1–7.3)
Growth NaCl	<ul style="list-style-type: none"> • <i>M. xanthus</i> prefer 1% w/v NaCl • <i>B. antiquum</i> and <i>H. salinarum</i> prefer 0.5% w/v NaCl ^c 	<ul style="list-style-type: none"> • <i>I. loihiensis</i> prefer 3.5% w/v NaCl • <i>B. pumilus</i> prefer 0.5 % w/v NaCl
Growth Ca ²⁺	<ul style="list-style-type: none"> • Positive effect of Ca²⁺ of 28 mg/L on <i>I. loihiensis</i> growth • Negative effect of Ca²⁺ of 28 mg/L on <i>M. xanthus</i> growth 	

426 a - Microorganisms produced bio-struvite only under aerobic conditions.

427 b - The *B. pumilus* were 33% positive for glucose utilisation.

428 c – Different findings from previous studies.

4 Conclusion

- Proteins/amino acids were the preferred organic carbon sources for the five microorganisms investigated.
- *B. pumilus*, *M. xanthus*, *H. salinarum* and *B. antiquum* were able to produce urease.
- *I. loihiensis* was found to be a facultative anaerobe able to use O₂ and NO₃-N as an electron acceptor.
- The preferred temperature for all selected microorganisms was within the mesophilic range (22–34 °C); most microorganisms preferred a neutral pH and NaCl concentrations less than 1% w/v, whereas *I. loihiensis* preferred a mild alkaline pH 8, high NaCl of 3.5% w/v and the presence of Ca²⁺.
- The selected microorganisms produced bio-struvite crystals under aerobic conditions. The morphology of crystals produced was dominantly coffin-lid and long-bar shapes.
- The bio-struvite production and PO₄-P removal highly depended on the microbial growth and DO level. At the investigated optimal growing conditions, in the presence of DO, the bio-struvite crystal (>10 µm) yield and PO₄-P removal varied between 1,521–1,746 mg/L and 55–76%, respectively.

5 Reference

- Alcaraz, G., 2015. Gisella Alcaraz Bacillus Pumilus [WWW Document]. URL https://microbewiki.kenyon.edu/index.php/Gisella_Alcaraz-Bacillus_Pumilus (accessed 1.1.19).
- Arias, D., Cisternas, L., Rivas, M., 2017. Biomineralization mediated by ureolytic bacteria

451 applied to water treatment: a review. *Crystals* 7, 345.
 452 <https://doi.org/10.3390/cryst7110345>

453 Bailey, J. V., Joye, S.B., Kalanetra, K.M., Flood, B.E., Corsetti, F.A., 2007. Evidence of
 454 giant sulphur bacteria in Neoproterozoic phosphorites. *Nature* 445, 198–201.
 455 <https://doi.org/10.1038/nature05457>

456 Bezerra, M.A., Santelli, R.E., Oliveira, E.P., Villar, L.S., Escaleira, L.A., 2008. Response
 457 surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*
 458 76, 965–977. <https://doi.org/10.1016/j.talanta.2008.05.019>

459 Claus, D., 1992. A standardized Gram staining procedure. *World J. Microbiol. Biotechnol.* 8,
 460 451–452. <https://doi.org/10.1007/BF01198764>

461 Coe, F.L., Evan, A., Worcester, E., 2005. Kidney stone disease. *J. Clin. Invest.* 115, 2598–
 462 2608. <https://doi.org/10.1172/JCI26662.2598>

463 Da Silva, S., Bernet, N., Delgenès, J.P., Moletta, R., 2000. Effect of culture conditions on the
 464 formation of struvite by *Myxococcus xanthus*. *Chemosphere* 40, 1289–1296.
 465 [https://doi.org/10.1016/S0045-6535\(99\)00224-6](https://doi.org/10.1016/S0045-6535(99)00224-6)

466 Gassama, U.M., Puteh, A. Bin, Abd-Halim, M.R., Kargbo, B., 2015. Influence of municipal
 467 wastewater on rice seed germination, seedling performance, nutrient Uptake, and
 468 chlorophyll content. *J. Crop Sci. Biotechnol.* 18, 9–19. [https://doi.org/10.1007/s12892-](https://doi.org/10.1007/s12892-014-0091-4)
 469 014-0091-4

470 Gavrish, E.I., Krauzova, V.I., Potekhina, N. V, Karasev, S.G., Plotnikova, E.G., Altyntseva,
 471 O. V, Korosteleva, L. a, Evtushenko, L.I., 2004. Three new species of *brevibacteria*,
 472 *Brevibacterium antiquum* sp. nov., *Brevibacterium aurantiacum* sp. nov. and

473 *Brevibacterium permense* sp. nov. *Microbiology* 73, 218–225.
 474 <https://doi.org/10.1023/B:MICL.0000023986.52066.1e>

475 González-Muñoz, M.T., De Linares, C., Martínez-Ruiz, F., Morcillo, F., Martín-Ramos, D.,
 476 Arias, J.M., 2008. Ca-Mg kutnahorite and struvite production by *Idiomarina* strains at
 477 modern seawater salinities. *Chemosphere* 72, 465–472.
 478 <https://doi.org/10.1016/j.chemosphere.2008.02.010>

479 González-Muñoz, M.T., Rodríguez-Navarro, C., Martínez-Ruiz, F., Arias, J.M., Merroun,
 480 M.L., Rodríguez-Gallego, M., 2010. Bacterial biomineralization: new insights from
 481 *Myxococcus*-induced mineral precipitation. *Geol. Soc. London, Spec. Publ.* 336, 31–50.
 482 <https://doi.org/10.1144/SP336.3>

483 Grace-Martin, K., 2012. Can a regression model with a small R-squared be useful? [WWW
 484 Document]. *Anal. Factor.* URL <http://www.theanalysisfactor.com/small-r-squared/>
 485 (accessed 2.1.19).

486 Janssen, G.R., Wireman, J.W., Dworkin, M., 1977. Effect of temperature the growth of
 487 *Myxococcus xanthus*. *J. Bacteriol.* 130, 561–562.

488 Le Corre, K.S., Valsami-Jones, E., Hobbs, P., Parsons, S. a., 2005. Impact of calcium on
 489 struvite crystal size, shape and purity. *J. Cryst. Growth* 283, 514–522.
 490 <https://doi.org/10.1016/j.jcrysgr.2005.06.012>

491 Leng, Y. and Soares, A. (2018) Understanding the fundamentals of bio-struvite
 492 biomineralization in wastewater, Cranfield University. PhD thesis, in review

493 Losensky, G., Jung, K., Urlaub, H., Pfeifer, F., Fröls, S., Lenz, C., 2017. Shedding light on
 494 biofilm formation of *Halobacterium salinarum* R1 by SWATH-LC/MS/MS analysis of

495 planktonic and sessile cells. *Proteomics* 17, 1–13.
 496 <https://doi.org/10.1002/pmic.201600111>

497 Maathuis, F.J., 2009. Physiological functions of mineral macronutrients. *Curr. Opin. Plant*
 498 *Biol.* 12, 250–258. <https://doi.org/10.1016/j.pbi.2009.04.003>

499 Massey, M.S., Davis, J.G., Ippolito, J.A., Sheffield, R.E., 2009. Effectiveness of recovered
 500 magnesium phosphates as fertilizers in neutral and slightly alkaline soils. *Agron. J.* 101,
 501 323–329. <https://doi.org/10.2134/agronj2008.0144>

502 Merroun, M.L., Chekroun, K. Ben, Arias, J.M., González-Muñoz, M.T., 2003. Lanthanum
 503 fixation by *Myxococcus xanthus*: Cellular location and extracellular polysaccharide
 504 observation. *Chemosphere* 52, 113–120. [https://doi.org/10.1016/S0045-6535\(03\)00220-](https://doi.org/10.1016/S0045-6535(03)00220-0)
 505 0

506 Mesbah, N., Wiegel, J., 2005. Halophilic thermophiles: A novel group of extremophiles, in:
 507 *Microbial Diversity: Current Perspectives and Potential Applications*. IK Publishing
 508 House, New Delhi, pp. 91–118.

509 Minitab, 2010. Minitab 17 Statistical Software.

510 Mormile, M.R., Biesen, M.A., Gutierrez, M.C., Ventosa, A., Pavlovich, J.B., Onstott, T.C.,
 511 Fredrickson, J.K., 2003. Isolation of *Halobacterium salinarum* retrieved directly from
 512 halite brine inclusions. *Environ. Microbiol.* 5, 1094–1102.
 513 <https://doi.org/10.1046/j.1462-2920.2003.00509.x>

514 Nocker, A., Cheswick, R., Dutheil de la Rochere, P.M., Denis, M., Léziart, T., Jarvis, P.,
 515 2017. When are bacteria dead? A step towards interpreting flow cytometry profiles after
 516 chlorine disinfection and membrane integrity staining. *Environ. Technol.* 38, 891–900.

517 <https://doi.org/10.1080/09593330.2016.1262463>

518 Orange, F., Westall, F., Disnar, J.R., Prieur, D., Bienvenu, N., Le Romancer, M., D'Éfarge, C.,
519 2009. Experimental silicification of the extremophilic Archaea *Pyrococcus abyssi* and
520 *Methanocaldococcus jannaschii*: applications in the search for evidence of life in early
521 Earth and extraterrestrial rocks. *Geobiology* 7, 403–418. [https://doi.org/10.1111/j.1472-](https://doi.org/10.1111/j.1472-4669.2009.00212.x)
522 [4669.2009.00212.x](https://doi.org/10.1111/j.1472-4669.2009.00212.x)

523 Poza, M., Sieiro, C., Villa, T.G., 2004. Cloning and expression of clt genes encoding milk-
524 clotting proteases from *Myxococcus xanthus* 422. *Appl. Environ. Microbiol.* 70, 1–6.
525 <https://doi.org/10.1128/AEM.70.10.6337>

526 Prywer, J., Torzewska, A., 2010. Biomineralization of struvite crystals by *Proteus mirabilis*
527 from artificial urine and their mesoscopic structure. *Cryst. Res. Technol.* 45, 1283–1289.
528 <https://doi.org/10.1002/crat.201000344>

529 Prywer, J., Torzewska, A., 2009. Bacterially induced struvite growth from synthetic urine:
530 experimental and theoretical characterization of crystal morphology. *Cryst. Growth Des.*
531 9, 3538–3543. <https://doi.org/10.1021/cg900281g>

532 Rivadeneyra, M.A., Delgado, R., Párraga, J., Ramos-Cormenzana, A., Delgado, G., 2006.
533 Precipitation of minerals by 22 species of moderately halophilic bacteria in artificial
534 marine salts media: Influence of salt concentration. *Folia Microbiol. (Praha)*. 51, 445–
535 453. <https://doi.org/10.1007/BF02931589>

536 Robinson, R.K., 2014. *Encyclopaedia of food microbiology* (1st ed.), Academic press.

537 Ronteltap, M., Maurer, M., Hausherr, R., Gujer, W., 2010. Struvite precipitation from urine -
538 Influencing factors on particle size. *Water Res.* 44, 2038–2046.

539 <https://doi.org/10.1016/j.watres.2009.12.015>

540 Sadowski, R.R., Prywer, J., Torzewska, A., 2014. Morphology of struvite crystals as an
 541 evidence of bacteria mediated growth. *Cryst. Res. Technol.* 49, 478–489.
 542 <https://doi.org/10.1002/crat.201400080>

543 Schultze-Lam, S., Fortin, D., Davis, B., Beveridge, T., 1996. Mineralization of bacterial
 544 surfaces. *Chem. Geol.* 132, 171–181. [https://doi.org/10.1016/S0009-2541\(96\)00053-8](https://doi.org/10.1016/S0009-2541(96)00053-8)

545 Shaddel, S., Ucar, S., Andreassen, J., Sterhus, S., 2019 Engineering of struvite crystals by
 546 regulating supersaturation- Correlation with phosphorus recovery, crystal, morphology
 547 and process efficiency. *J. Environ. Chem. Eng.* Elsevier Ltd, 7(1).
 548 <https://doi.org/10.1016/j.jece.2019.102918>

549 Shivaji, S., Chaturvedi, P., Suresh, K., Reddy, G.S.N., Dutt, C.B.S., Wainwright, M.,
 550 Narlikar, J. V., Bhargava, P.M., 2006. *Bacillus aerius* sp. nov., *Bacillus aerophilus* sp.
 551 nov., *Bacillus stratosphericus* sp. nov. and *Bacillus altitudinis* sp. nov., isolated from
 552 cryogenic tubes used for collecting air samples from high altitudes. *Int. J. Syst. Evol.*
 553 *Microbiol.* 56, 1465–1473. <https://doi.org/10.1099/ijs.0.64029-0>

554 Silva, T. Da, Cássia, V. De, Amanda Lais de Souza Coto, Rafael de Carvalho Souza,
 555 M.B.S.N., Gomes, E., Bonilla-Rodriguez, G.O., 2016. Effect of pH, temperature, and
 556 chemicals on the endoglucanases and β -glucosidases from the thermophilic fungus
 557 *Myceliophthora heterothallica* F.2.1.4. obtained by solid-state and submerged cultivation.
 558 *Biochem. Res. Int.* 2016. <https://doi.org/10.1155/2016/9781216>

559 Simoes, F., Vale, P., Stephenson, T., Soares, A., 2017. Understanding the growth of the bio-
 560 struvite production *Brevibacterium antiquum* in sludge liquors. *Environ. Technol.* 1–10.

561 <https://doi.org/10.1080/09593330.2017.1411399>

562 Sinha, A., Singh, A., Kumar, S., Khare, S.K., Ramanan, A., 2014. Microbial mineralization
563 of struvite: a promising process to overcome phosphate sequestering crisis. *Water Res.*
564 54, 33–43. <https://doi.org/10.1016/j.watres.2014.01.039>

565 Smirnov, A., Suzina, N., Chudinova, N., Kulakovskaya, T., Kulaev, I., 2005. Formation of
566 insoluble magnesium phosphates during growth of the archaea *Halorubrum distributum*
567 and *Halobacterium salinarum* and the bacterium *Brevibacterium antiquum*. *FEMS*
568 Microbiol. Ecol. 52, 129–137. <https://doi.org/10.1016/j.femsec.2004.10.012>

569 Soares, A., Veeram, M., Simoes, F., Wood, E., Parsons, S. a., Stephenson, T., 2014. Bio-
570 Struvite: A new route to recover phosphorus from wastewater. *Clean - Soil, Air, Water*
571 42, 994–997. <https://doi.org/10.1002/clen.201300287>

572 Syberg, S., 2016. Reducing the edge effect in cell culture microplates [WWW Document].
573 Thermo Fish. Sci. URL [https://www.rdmag.com/article/2016/10/reducing-edge-effect-](https://www.rdmag.com/article/2016/10/reducing-edge-effect-cell-culture-microplates)
574 cell-culture-microplates (accessed 2.1.19).

575 Tansel, B., Lunn, G., Monje, O., 2018. Struvite formation and decomposition characteristics
576 for ammonia and phosphorus recovery: A review of magnesium-ammonia-phosphate
577 <https://doi.org/10.1016/j.chemosphere.2017.12.004>

578 Tchobanoglous, G., Burton, F.L., Stensel, H.D., Metcalf&Eddy, 2003. Wastewater
579 engineering: Treatment and reuse, 4th ed. McGraw-Hill Education.

580 Thomas, S.H., Wagner, R.D., Arakaki, A.K., Skolnick, J., Kirby, J.R., Shimkets, L.J.,
581 Sanford, R.A., Löffler, F.E., 2008. The mosaic genome of *Anaeromyxobacter*
582 dehalogenans strain 2CP-C suggests an aerobic common ancestor to the delta-

583 proteobacteria. PLoS One 3. <https://doi.org/10.1371/journal.pone.0002103>

584 Torzewska, A., Stączek, P., Różalski, A., 2003. Crystallization of urine mineral components
585 may depend on the chemical nature of *Proteus* endotoxin polysaccharides. J. Med.
586 Microbiol. 52, 471–477. <https://doi.org/10.1099/jmm.0.05161-0>

587 Trinh, C.T., Srienc, F., 2009. Metabolic engineering of *Escherichia coli* for efficient
588 conversion of glycerol to ethanol. Appl. Environ. Microbiol. 75, 6696–6705.
589 <https://doi.org/10.1128/AEM.00670-09>

590 Trujillo, M.E., Goodfellow, M., 2015. *Brevibacterium*, in: Bergey's Manual of Systematics of
591 Archaea and Bacteria. John Wiley & Sons, Inc., in association with Bergey's Manual
592 Trust, pp. 1–22. <https://doi.org/10.1002/9781118960608.gbm00062>

593 Ye, Z., Shen, Y., Ye, X., Zhang, Z., Chen, S., Shi, J., 2014. Phosphorus recovery from
594 wastewater by struvite crystallization: Property of aggregates. J. Environ. Sci. 26, 991–
595 1000. [https://doi.org/10.1016/S1001-0742\(13\)60536-7](https://doi.org/10.1016/S1001-0742(13)60536-7)

596 Yee, N., Phoenix, V.R., Konhauser, K.O., Benning, L.G., Ferris, F.G., 2003. The effect of
597 cyanobacteria on silica precipitation at neutral pH: Implications for bacterial
598 silicification in geothermal hot springs. Chem. Geol. 199, 83–90.
599 [https://doi.org/10.1016/S0009-2541\(03\)00120-7](https://doi.org/10.1016/S0009-2541(03)00120-7)

600 Zinder, S.H., Dworkin, M., 2013. Morphological and physiological diversity, in: The
601 Prokaryotes: Prokaryotic Biology and Symbiotic Associations. Springer New York, pp.
602 185–220. https://doi.org/10.1007/978-3-642-30194-0_9



DSPACE

<https://dspace.org/>

Understanding the biochemical characteristics of struvite bio-mineralising microorganisms and their future in nutrient recovery

Leng, Yirong; Colston, Robert; Soares, Ana

2020-01-08

Elsevier

<https://doi.org/10.1016/j.chemosphere.2019.125799>

Leng Y, Colston R, Soares A. (2020) Understanding the biochemical characteristics of struvite bio-mineralising microorganisms and their future in nutrient recovery. Chemosphere, Volume 247, May 2020, Article number 125799