

Geographical variation of cuticular hydrocarbon profiles of adult flies and empty puparia amongst three populations of *Calliphora vicina* (Diptera: Calliphoridae)

Canan Kula^{1,*}, Jens Amendt², Falko P. Drijfhout³ and Hannah E. Moore¹

¹Cranfield Forensic Institute, Cranfield University, Defence Academy of the United Kingdom, Shrivenham, Wiltshire SN6 8LA, UK

²Institute of Legal Medicine, University Hospital Frankfurt/Main, Goethe-University, 60323 Frankfurt/Main, Germany

³Chemical Ecology Group, School of Chemical and Physical Science, Keele University, Keele ST5 5BG, UK

*Correspondence: c.kula@cranfield.ac.uk

Abstract

Blowflies (Diptera: Calliphoridae) are of great importance in forensic entomology and in determining the minimum post-mortem interval, as they may be the first group of insects to colonise decomposing remains. Reliable species identification is an essential prerequisite. Classically, morphological characters or DNA sequences are used for this purpose. However, depending on the species and the condition of the specimen, this can be difficult, e.g. in the case of empty fly puparia. Recent studies have shown that cuticular hydrocarbon (CHC) profiles are species-specific in necrophagous taxa and represent another promising tool for identification. However, the population-specific variability of these substances as a function of e.g. local climatic parameters has not yet been sufficiently investigated. The aim of this study was to determine the geographical variation of CHC profiles of the blowfly *Calliphora vicina* depending on different countries of origin. Flies were reared in the UK, Germany and

Turkey in common garden experiments under ambient conditions. CHC profiles of the resulting adult flies and their empty puparia were analysed using gas chromatography-mass spectrometry. Data were visualised by principal component analysis and clustered by population. The populations of the UK and Germany, both having similar climates and being geographically close to each other, showed greater similarities in CHC profiles. However, the CHC profile of the Turkish population, whose climate is significantly different from the other two populations, was very different. Our study confirms the high potential of CHC analysis in forensic entomology but highlights the need to investigate geographical variability in chemical profiles.

Keywords:

Calliphoridae, adult fly, empty puparia, cuticular hydrocarbons, GC-MS, Turkey, Germany, United Kingdom

Introduction

Calliphoridae (Diptera) is a significant family in the field of forensic entomology due to the fact that they are among the primary insects colonised on human bodies. (Benecke, 2001; Amendt et al., 2004). Forensic investigators consider them to be crucial evidence regarding the time and location of death (Catts and Goff, 1992; Amendt et al., 2007; Gennard, 2007; Moore et al., 2017a). A minimum post-mortem interval (minPMI) can be determined by using the arrival of adult insects and the resulting colonisation patterns in conjunction with the age estimation of the immatures developing on the dead body (Michaud and Moreau, 2009; Pechal et al., 2014). Accurate species identification is important for estimating a minPMI, as the duration of the life cycle and the ageing process varies by species (Moore et al., 2013; Paula et al., 2017). For an

accurate calculation, reference growth data obtained in the laboratory are used for the respective species (Moore and Shemilt, 2018). However, a potential source of error in determining minPMI is ignoring the fact that data from the laboratory may not reflect the developmental adaptations of the local population of interest. Therefore, Tarone et al. (2011) emphasised the need to determine how ecological conditions influence physiological responses and can lead to differences in growth rate between populations of the same species that exist many hundreds or thousands of kilometres apart, e.g. on a latitudinal scale. Hwang and Turner (2009) suggested the existence of local adaptations that can be seen on a small scale that different levels of urbanisation may cause for the forensically important blowfly *Calliphora vicina* (Robineau-Desvoidy, 1830), but, e.g. Limsopatham et al. (2018) found similar larval growth rates for this species from the UK and German populations at different temperatures during the major part of development.

Another important aspect of forensic entomology is providing information whether the crime was committed at the place of discovery. If a corpse is colonised with insects typical of a different habitat or geographical region, this may indicate that the corpse has been transported. However, Charabidze et al. (2017) rightly point out that since the majority of European necrophagous species have wide distribution ranges that cover many countries, sampling non-native species inside European regions is improbable. Additionally, although each species has a preferred ecological environment, individuals from the species may sometimes be found outside of the chosen range. Hence, it is difficult to link a certain species with a particular location or environment.

For proving population-specific differences and habitat adaptations, which could be linked to a change in development, the differentiation or assignment of such clusters would be desirable. For this purpose, e.g. molecular biological and morphological analyses are suitable. (Kranz et al., 2017). Zhou et al. (2020) investigated phylogenetic relationships among Calliphoridae

species using DNA-based sequence data and measured the degree of genetic variation. Landmark-based geometric morphometric analysis has also recently been employed in entomology. However, DNA-based techniques require molecular biology experience and extensive equipment to perform gene sequencing and so far do not reveal any evidence of a consistent difference between populations. While morphometric data collection is manageable, statistical and computer-assisted analysis requires skills and experience. Also, different geographical populations could not separate from each other (Limsopatham et al., 2018). An alternative technique that has the potential to distinguish the variations in various populations of the same forensically important species is cuticular hydrocarbon (CHC) analysis.

For a few years now, the chemistry of the insect cuticle has been studied as a new tool in forensic entomology. The cuticle is coated with a waxy layer consisting of hydrocarbons, fatty acids, and alcohols (Gibbs, 1998a). The primary function of the cuticular layer is to maintain water balance and prevent desiccation; as insects adapt to new habitats with different temperatures and climate types, the composition of the layer changes (Chung and Carroll, 2015). The hydrocarbons are found as saturated (alkanes) and unsaturated forms (alkenes and alkynes), which can have one or more methyl groups. Unsaturated forms can contain one, two, or three double bonds along the chain length (Moore et al., 2013). Long-chain hydrocarbons with higher boiling points provide more efficient protection at higher temperatures than methyl-branched and unsaturated hydrocarbons with lower boiling points (Drijfhout et al., 2010). However, methyl-branched and unsaturated hydrocarbons are present in the epicuticular layer even though they reduce the ability of the protective barrier, as they are required for signal exchange during chemical communication in nestmate recognition and sex determination (Toolson, 1984). need more mobility and flexibility, especially in the early immature stages. To increase the flexibility, branched methyl alkanes and alkenes with lower melting points are

needed. As the larvae grow and develop, especially in the post-feeding stage, mobility and flexibility decrease and longer chain hydrocarbons are needed (Drijfhout et al., 2010).

The biophysical properties of surface epicuticular lipids also vary among the same species, and this variation is probably due in part to genetic and environmental factors (Gibbs et al., 1991). Noorman and Den Otter (2002) investigated the effects of relative humidity, temperature, and population density on the CHC profiles of adult *Musca domestica* L. (house flies) and showed that the CHC profile was affected by relative humidity and temperature, and the change was traceable.

The pupae are immobile, and protection against dehydration is vital; when the larvae face threats such as sunlight or lower relative humidity, they can move away, but pupae cannot escape exposure (Paula et al., 2017). Therefore, a cuticle that provides effective waterproofing to the pupae may avoid fatal impacts. Toolson (1982) examined the effects of larval and puparial thermal regimes on transcuticular water loss rates and epicuticular composition in adult *Drosophila pseudoobscura* (fruit flies) and showed that the temperatures exposed during the puparial period are influential in shaping the cuticular profiles of adult flies. Toolson and Hadley (1979) examined the cuticle permeability and lipid composition of *Centruroides sculpturatus* Ewing 1928 (scorpions), which they collected in the summer and winter seasons, and revealed the seasonal effect on the structure. Kruger and Pappas (1993) demonstrated the geographic variation of the CHC profile among fourteen populations of *Aedes albopictus* (Skuse, 1895) (mosquitoes). Chung and Carroll (2015) investigated how ecological variation in insect CHC profiles can lead to differentiation in mating signals and reproductive isolation. They noted that the evolution of CHC profiles might be the best method for understanding ecological speciation.

Previously, geographical differences were studied by examining the CHC profiles of beetles (Chown et al., 2011; Hadley, 1978; Howard et al., 1995), wasps (Espelie et al., 1990; Michelutti et al., 2018), cockroaches (Brown et al., 2000; Everaerts et al., 1997), grasshoppers (Gibbs et al., 1991; Tregenza et al., 2000), sandflies (Kamhawi et al., 1987), ants (Martin et al., 2008; Menzel et al., 2017), mosquitoes (Anyanwu et al., 2000; Kruger et al., 1991; Kruger and Pappas, 1993; Rosa-Freitas et al., 1992) and fruit flies (Bontonou et al., 2013; Chung and Carroll, 2015; Ingleby et al., 2013; Markow and Toolson, 2013; Rajpurohit et al., 2017; Rouault et al., 2001, 2004). However, studies on blowfly species are rather limited (Byrne et al., 1995; Moore et al., 2022; Paula et al., 2017, 2018, 2020).

This study aims to investigate the variation between the chemical profiles of the forensically important blowfly, *Calliphora vicina* (Diptera: Calliphoridae), and their empty puparia reared outdoors in different habitats in the UK, Germany and Turkey. It will also determine if any potential local adaptations have an influence on their chemical profiles. Differentiation will be based on qualitative and quantitative differences in CHC profiles.

Materials and methods

Insect materials

C. vicina were obtained from three locations: Swindon/United Kingdom (51°36'23.2"N 1°38'06.5"W), Frankfurt/Germany (50°05'39.2"N 8°39'52.4"E) and Ankara/Turkey (39°52'24.9"N 32°43'48.7"E) (see in Figure 1). Straight-line distances are over 740 km between populations of the UK and Germany, over 2190 km between populations of Germany and Turkey, and over 2930 km between populations of the UK and Turkey. The rearing of adult flies and creating of the empty puparia were carried out separately in the relevant countries. The adult flies were collected from the wild in the summer months and placed outdoors in rearing

cages under ambient environmental conditions and were supplied with blood, sugar, water and milk powder. Fresh lamb liver was used as an oviposition medium. Once fly eggs were laid on the oviposition medium, the Petri dish containing the eggs and meat was transferred into small rearing boxes. The boxes were also placed in outdoor cages. During the pupariation period, the cages were checked daily. On the first day of metamorphosis, the first generation of adult flies and their empty puparia were collected and transported to the laboratory in the relevant countries for the extraction process.

Sample preparation

The liquid extraction method described by Moore et al. (2017a) was chosen for CHCs extraction. For the extraction of CHCs, a non-polar solvent (hexane) was used. The extractions were carried out ten times ($n = 10$) without gender separation for each population. For adult fly and their empty puparia extractions, a specimen was placed into a GC vial with hexane, ensuring that the specimen was fully submerged (500 μL for adult flies, 350 μL for empty puparia) and left for 10-15 minutes. The hexane extract containing the CHC profile was directly transferred to a clean vial. Then, the hexane was left to evaporate until the extract could be transferred to 300 μL glass inserts and left to dry down completely. Finally, all samples were stored in the refrigerator at 4°C until they were required for analysis. The dried extract was then reconstituted in 30 μL of hexane for GC-MS analysis.

Chemical analysis of extracts

Chemical analysis of extraction samples prepared in the laboratories of the relevant countries was performed in the same laboratory on the same Agilent Technologies 6890N Network GC system with a split/splitless injector at 250°C, a Restek Rxi-1MS capillary column containing an SP of 100% Polydimethylsiloxane (30 m x 0.25 mmID, 0.25 μm film thickness) coupled to

an Agilent 5973 Network Mass Selective Detector. The GC was connected to a computer, and the data was processed with Agilent Chemstation software. Elution was carried out with helium at 1 mL/min. The oven temperature was programmed to be held at 50°C for 2 minutes, then ramped up to 200°C at 25°C/min, then from 200°C to 260°C at 3°C/min and finally from 260°C to 320°C at 20°C/min where it was held for 2 minutes. The mass spectrometer was operated in Electron Ionisation at 70 eV, scanning from 40-500 amu at 1.5 scans/s. CHCs were identified using a library search (NIST08).

Statistical analysis

Datasets from the chromatograms were initially interpreted using PCA to facilitate the visualisation of trends that might be found. For more details, the reader is referred to Moore and Drijfhout (2013; 2015). PCA has already been used by the authors in previous studies (Moore et al., 2013, 2016, 2017a, 2017b), and the same experimental design was used for this study. CHCs with a percentage peak area of less than 0.5% were excluded for statistical analysis. For PCA, peak areas on chromatograms were identified using Agilent Chemstation software, focusing on only CHCs. All data were normalised before the PCA calculation. In most experiments, normalisation is accomplished by dividing the absolute abundance (concentration) of each component by the total abundance of all compounds, and a relative abundance is calculated, therefore minimising the effects of the high within-treatment variation (Moore and Drijfhout, 2015). The n-alkanes used for this study ranged from C23:H to C31:H. The NCSS 2020 software was used to construct the 3D graphic plane using the dataset acquired by PCA.

Results

CHC profile analysis and chemical identification

187 **Adult flies**

188 A total of 46 CHC compounds were identified in the CHC profiles of the adult *C. vicina*
189 collected from both the UK and Germany, and 31 CHCs in those collected from Turkey. The
190 CHCs detected in the chromatograms are given in Table 1, with the total percentage peak area
191 and standard deviation of each compound. All three populations had the same number of n-
192 alkanes. One of the most noticeable differences between the three populations was the number
193 of branched methyl alkanes and alkenes. The Turkish population showed fewer branched
194 methyl alkanes than the UK and German populations. While one alkene compound was detected
195 in the Turkish population, the alkenes detected in the UK and German populations were
196 excluded from the comparison because of insufficient concentrations. In the CHC profiles, n-
197 alkanes with odd-chain-length (OLA) were detected in higher concentrations than those with
198 even-chain-length n-alkanes (ELA) (Roux et al., 2008). C27:H (Table 1, compound 20) was the
199 dominant n-alkane for all three populations, and its percentage concentration was similar in all
200 the populations. The next predominant compound with a higher concentration in all profiles
201 was C29:H (Table 1, compound 35), and its percentage concentration in the Turkish population
202 was 4% higher than in the UK and German populations, where the percentage concentrations
203 of C29:H were similar.

204 When looking at the difference between populations, it was found that 9-MeC25 (Table 1,
205 compound 7) was detected exclusively in the Turkish population, 3-MeC25 (Table 1, compound
206 11) only in the UK, and 5-MeC26 (Table 1, compound 16) only in the German population.
207 11+13-MeC27 (Table 1, compound 21) was the most predominant branched methyl alkane in
208 all three populations. However, differences in its concentration differed between populations.
209 Its concentration in the UK and Germany populations was almost the same but lower than in
210 the Turkish population. In the UK and German populations, which have almost similar CHC
211 profiles, 3-MeC27 (Table 1, compound 27) and 11+13-MeC29 (Table 1, compound 37) also

212 had higher but different quantitative concentrations than other branched methyl alkanes, while
213 7-MeC29 had the second-highest concentration in the Turkish population.

214 Figure 2 shows the GC chromatograms of adult flies of *C. vicina* from the UK, German and
215 Turkish populations. The CHC profiles were extracted on the first day immediately after
216 eclosion. While the chemical profiles of the UK and German populations were quite similar,
217 the number and concentration of CHCs with a shorter carbon chain were much higher in the
218 Turkish population than in the other populations. In addition, there were CHCs with higher
219 retention times (also known as longer chain CHCs) in the Turkish population that were not
220 found in the UK and German populations.

221 Figure 3 compares the total concentrations of CHCs detected in the chemical profiles of adult
222 flies against the populations. The concentrations of branched methyl alkanes were significantly
223 higher than the concentrations of n-alkanes. The Turkish population had greater total
224 concentrations of n-alkanes than the other populations, although all populations had the same
225 total number of n-alkanes found in their profiles. The UK and German populations already had
226 a higher number of branched methyl alkanes, and their total concentration was higher than that
227 of the Turkish flies. In the UK and German populations, no acceptable concentrations of alkenes
228 were detected. Only the Turkish population had a substantial concentration of alkenes in their
229 chemical profiles.

230 To visualise the findings and to statistically differentiate the populations, PCA computations
231 were carried out with the CHC data presented in Table 1. PCA was performed using six
232 principal components, describing 95.7% of the dataset variation within the data set with the first
233 three principal components, comprising 86.1%, 7.0% and 2.6%, respectively. Each sample was
234 represented by a single data point in the graph, and each population had ten replicates. The 3D
235 plot has been chosen because it allows for a more visual representation of the distinction

between the three different populations. In order to create the PCA plot seen in Figure 4, PC2, PC3, and PC4 scores were used.

Discriminant analysis was used to discover which CHCs were most significant for segregating populations of *C. vicina* adult flies from the UK, Germany, and Turkey. Compounds exhibiting large loading scores will be the most significant in the separation seen in the PCA plot. The main compounds with significant loading values in the PCA dataset were 4-Methylhexacosane, 3-Methylheptacosane, 7-Methylheptacosane, and 3-Methyloctacosane compounds. Furthermore, since branched methyl alkanes had the highest total percentage among the CHC groups, it proved that these compounds were discriminant among the populations.

Empty puparia

The chain lengths of the detected CHCs range from C21:H to C32:H. In the chemical profile of the UK population, 28 CHC compounds were detected, compared to 27 CHCs in the German population and 24 CHCs in the Turkish population (Table 2). The same number of n-alkanes was found in all three groups. The concentration of n-alkanes was much greater than that of the other CHC groups. Nevertheless, while the number of branched methyl alkanes was similar in the UK and German populations, fewer branched methyl alkanes were detected in the Turkish population. Only one alkene compound with sufficient concentration (>0.5%) was detected in the German population.

C27:H and C29:H (Table 2, compounds 8 and 22) were the most abundant n-alkanes in the populations of the UK and Turkey. While the C27:H in the UK population was the most highly concentrated than found in others, the C29:H concentration in the Turkish population was higher than in other populations. C25:H and C27:H (Table 2, compounds 4 and 8), on the other hand, were the most prevalent n-alkanes in the German population. The concentration of C25:H

found in the German population was significantly greater than those found in the populations of the UK and Turkey.

The dominant branched methyl alkanes were the same in all three populations, 11+13-MeC27, 3-MeC27 and 11+13-MeC29 (Table 2, compounds 9, 13 and 24), but their concentrations varied between populations. While a branched methyl alkane could not be specifically detected for the Turkish population, 3-MeC21 and 5-MeC28 (Table 2, compounds 1 and 18) were detected only in the UK population, and 3-MeC30 and 3-MeC31 (Table 2, compounds 29 and 32) were detected only in the German population. Likewise, 13+15-MeC28 (Table 2, compound 15) was detected only in UK and German populations, and 11+13-MeC28 and 13+15-MeC29 (Table 2, compounds 16 and 23) were detected only in UK and Turkey populations. The presence or absence of branched methyl alkanes in the profiles or differences in their concentrations can be considered an indicator of discrimination between populations.

The chromatograms in Figure 5 represent the chemical profiles of *C. vicina* empty puparia collected from the UK, Germany and Turkey. Chromatograms were similar in general appearance. However, compound concentrations differed between populations. Whereas the CHC profiles of the populations of the UK and Germany were similar, the CHC concentrations in the Turkish population were different from them. The CHC concentrations detected in the Turkish population were higher than in the other two populations.

Figure 6 shows the total concentrations of CHC compounds detected in the CHC profiles of all three populations. According to the graph, the total n-alkane concentration detected in the Turkish population was 8% higher than in the UK, which had the lowest n-alkane concentration. On the other hand, the branched methyl alkane concentration detected in the UK population was 11% higher than the German population, with the lowest branched methyl alkane concentrations. Sufficient alkene concentrations (>0.5%) could not be detected in the UK and

283 Turkish populations. Only one alkene was detected at a 5% concentration in the German
284 population.

285 PCA calculations were carried out using the CHC data presented in Table 2 to aid the
286 visualisation of the findings and statistically discriminate between the different populations.
287 PCA was performed using six principal components, describing 99.4% of the dataset variation
288 within the data set with the first three principal components, comprising 85.4%, 12.8% and
289 1.2%, respectively. The PC2, PC3, and PC4 components were used to create the PCA plot
290 shown in Figure 7.

291 Discriminant analysis was used to discover which compounds were most significant for
292 separating *C. vicina* empty puparia from the UK, Germany, and Turkey populations.
293 Compounds exhibiting large loading scores will be the most significant in the separation seen
294 in the PCA plot. The main compounds with significant loading values in the PCA dataset were
295 pentacosane (C25:H), heptacosane (C27:H), and nonacosane (C29:H). As n-alkanes, especially
296 OLAs, had the highest total percentage among the CHC groups, it showed that these n-alkanes
297 were discriminant among the populations.

298 **Discussion**

299 The results showed qualitative and quantitative differences between the three populations of *C.*
300 *vicina* in both adult flies and their empty puparia. These differences are also visible in the
301 chromatograms showing the chemical profiles of the populations. The differences in compound
302 concentrations across chromatograms suggested that CHC profiles of the same species are
303 retained in a unique pattern and can also be used to distinguish populations of the same species
304 living in different geographical areas. PCA analysis helps to visualise data from GC
305 chromatograms that allow differentiating between populations. It was unnecessary to separate

the samples by sex for this research since the purpose was to investigate the variations in chemical profiles across various populations of the same species.

Moore et al. (2017a) determined the age-dependent chemical changes that occur within the adult profiles of *C. vicina*, *Calliphora vomitoria* (Linnaeus, 1758) and *Lucilia sericata* (Meigen, 1826) over a period of 30 days, allowing for the age of the adult flies to be estimated. Therefore, to reduce the influence of age on the chemical profiles and examine the sole geographical effect, the adult fly populations used in this research were chosen as the first generation, derived from the *C. vicina* females collected from the wild and reared outdoors.

Howard and Blomquist (2005) reported that, in addition to being species-specific, geographically segregated populations might have qualitative and quantitative differences in CHC profiles depending on genetic factors and environmental conditions. Chapman et al. (1995) pointed out higher proportions of linear alkanes in populations living in environments with higher temperatures than other compounds. The general pattern is that species living in warmer and drier habitats are expected to lose water more rapidly than populations living in cooler and moister habitats. CHCs with longer carbon chains provide great potential against water loss (Gibbs, 1998b). Branched methyl alkanes prevent tight molecular packing, which reduces membrane fluidity and permeability (Drijfhout et al., 2010). Hot and dry ambient temperatures cause the loss of CHC with a short carbon chain that volatilises quickly. The presence of compounds with longer carbon chains in the CHC profiles of insects living in warmer climates supports the hypothesis that the epicuticular layer is regulated against water loss (Gibbs, 1998a, 1998b).

Studies focusing mainly on *Drosophila* spp. (Bontonou et al., 2013; Chung and Carroll, 2015; Ingleby et al., 2013; Markow and Toolson, 2013; Rajpurohit et al., 2017; Rouault et al., 2001, 2004) have revealed differences among populations. However, few studies have been

conducted regarding its importance in forensic entomology; however, these studies were mostly carried out under controlled laboratory conditions. Byrne et al. (1995) examined the CHC profiles of both close and distant populations of *Phormia regina* (Meigen, 1826) adult blowflies collected from three locations. Populations were analysed using GC–MS, and identified the CHCs. Paula et al. (2017) determined the chemotaxonomic profile and intraspecific variation of CHC compounds in adults of the blowfly *Chrysomya megacephala* (Fabricius, 1794) using discriminant function analysis with GC-MS analysis. These studies successfully identified the species and distinguished them between the different locations. The results showed that there were qualitative and quantitative differences between samples from the populations, and the percentage of short carbon chain CHC was higher in the population living in a cooler and humid environment. In another study, Paula et al. (2020) demonstrated that mid-infrared photoacoustic spectroscopy could be used to evaluate the variation in chemical profiles of *C. megacephala* adults collected from different populations. The results obtained were supportive of the present study and supported that the chemical profiles of *C. vicina* adults vary between both close and more distant populations.

The usefulness of empty puparia in criminal investigations is tenuous, as they are difficult to identify depending on the species and are not narrowly defined in terms of age. They only show the completed development of at least the first colonisation wave. (Moore et al., 2022). However, they could be very useful in crimes where death occurred a long time ago, and empty puparia are the only entomological evidence available (Moore et al., 2017b). So far, only a few studies have investigated the identification of empty puparia, as well as how geographical location and the local temperature or environment could influence the chemical profiles of necrophagous flies and their puparia. Paula et al. (2018) developed a new method to determine the PMI based on chemical compounds of the empty puparia from different oviposition cycles of *C. megacephala*. They examined the chemical composition of 50 empty puparia from

different oviposition cycles using GC-MS. The results show that chemical profiles can be used to distinguish between empty puparia. Moore et al. (2022) analysed the chemical profiles of empty puparia from seven forensically important blowfly species. They used CHC profiles for identification and also investigated geographical differences by comparing profiles of the same species from different regions of four countries. Results showed differences between the profiles of *C. vicina* from Germany, Spain, Norway and England, and also geographical locations could be determined from this chemical analysis.

In the present study, the changes in the CHC composition of *C. vicina* adult flies and empty puparia between the UK, German, and Turkish populations could be explained by the variances in environmental circumstances, distance and climate. While the difference between the CHC profiles of the UK and Turkish populations was greater, the German population shared several characteristics with both of these populations. Between the CHC profiles of the UK and German populations, there was a larger degree of resemblance than between the CHC profiles of the German and Turkish populations. This might be attributed to the fact that the distance between the UK and Germany is relatively shorter and that the two countries have comparable climatic types. The Turkish population differed from the other two populations due to its geographical distance and different climate type. Species from humid climates like the UK have more alkenes and branched methyl alkanes than species living in drier habitats, such as in Turkey.

The changes in CHCs concentrations in the chromatograms of populations are primarily related to the complete chemical profile rather than specific compounds. PCA was applied to the dataset instead, as it would be rather difficult to distinguish the differences in the chromatograms visually. PCA has the advantage of not focusing on specific compounds, as it is not known in advance which compounds are the most indicative for identifying populations (Moore et al., 2013). However, there are a few peaks that stand out due to their significant PCA scores, indicating they are effective within the dataset. These values belonged to branched

380 methyl alkanes, the dominant group in adult flies, and n-alkanes, which were the most dominant
381 compounds in empty puparia.

382 Limsopatham et al. (2018) examined the biological differences between the *C. vicina*
383 populations in Germany and UK by comparing developmental rates, wing morphometrics, and
384 molecular analyses. The separation of populations at a smaller geographic scale, however,
385 remained unclear. Molecular phylogenetic analysis by the maximum likelihood method failed
386 to distinguish between distinct geographic populations on a national or local level. DNA-based
387 approaches may be a viable alternative to morphological analysis, as genotyping is often more
388 rapid and straightforward. However, DNA analysis has several inherent risks, including DNA
389 degradation, ineffective amplification, and contamination (Mazzanti et al., 2010).

390 The results present in this preliminary study revealed that the cuticular hydrocarbon analysis is
391 a method capable of providing more information in forensic entomology. It could also serve as
392 an indication of possible population differences beneficial in forensic casework. On a small
393 scale, identifying variations between distinct populations of the same species at the national
394 level will be the pioneer contribution to forensic entomology in determining the populations in
395 different regions of the same country. Future studies that will include more species, locations
396 and larger datasets are needed to understand the intraspecific variability better.

397 **Acknowledgements** Canan Kula is supported by the Ministry of National Education of the
398 Republic of Turkey.

399 **References**

400 Amendt, J., Krettek, R., Zehner, R. (2004) 'Forensic entomology', *Naturwissenschaften*, 91(2),
401 pp. 51–65.

402 Amendt, J., Campobasso, C.P., Gaudry, E., Reiter, C., LeBlanc, H.N., Hall, J.R.M. (2007) 'Best
 403 practice in forensic entomology - Standards and guidelines', *International Journal of Legal*
 404 *Medicine*, 121(2), pp. 90–104.

405 Anyanwu, G.I., Molyneux, D.H. and Phillips, A. (2000) 'Variation in cuticular hydrocarbons
 406 among strains of the *Anopheles gambiae* sensu stricto by analysis of cuticular hydrocarbons
 407 using gas liquid chromatography of larvae', *Memorias do Instituto Oswaldo Cruz*, 95(3), pp.
 408 295–300.

409 Benecke M. (2001), 'A brief history of forensic entomology', *Forensic Science International*,
 410 120, pp. 2–14.

411 Bontonou, G., Denis, B. and Wicker-Thomas, C. (2013) 'Interaction between temperature and
 412 male pheromone in sexual isolation in *Drosophila melanogaster*', *Journal of Evolutionary*
 413 *Biology*, 26(9), pp. 2008–2020.

414 Brown, W. V., Rose, H.A., Lacey, M.J. and Wright, K. (2000) 'The cuticular hydrocarbons of
 415 the giant soil-burrowing cockroach *Macropanesthia rhinoceros* Saussure (Blattodea:
 416 Blaberidae: Geoscapheinae): Analysis with respect to age, sex and location', *Comparative*
 417 *Biochemistry and Physiology - B Biochemistry and Molecular Biology*, 127(3), pp. 261–277.

418 Byrne, A.L., Camann, M.A., Cyr, T.L., Catts, E.P. and Espelie, K.E. (1995) 'Forensic
 419 implications of biochemical differences among geographic populations of the black blow fly,
 420 *Phormia regina* (Meigen)', *Journal of Forensic Science*, 40(3), pp. 372–77.

421 Catts E., Goff M., (1992) 'Forensic entomology in criminal investigation', *Annual Review of*
 422 *Entomology*, 37, pp. 253–272.

423 Chapman, R.F., Espelie, K.E. and Sword, G.A. (1995) 'Use of cuticular lipids in grasshopper
 424 taxonomy: A study of variation in *Schistocerca gossypi* (Thomas)', *Biochemical Systematics*
 425 *and Ecology*, 23(4), pp. 383–398.

426 Charabidze, D., Gosselin, M. and Hedouin, V. (2017) 'Use of necrophagous insects as evidence
 427 of cadaver relocation: Myth or reality?', *PeerJ*, pp. 1–32.

428 Chown, S.L., Sørensen, JG and Terblanche, JS (2011) 'Water loss in insects: An environmental
 429 change perspective', *Journal of Insect Physiology*, 57(8), pp. 1070–1084.

430 Chung, H. and Carroll, S.B. (2015) 'Wax, sex and the origin of species: Dual roles of insect
 431 cuticular hydrocarbons in adaptation and mating', *BioEssays*, 37(7), pp. 822–830.

432 Drijfhout, F.P., (2010) 'Cuticular hydrocarbons: A new tool in forensic entomology?', in: J.
 433 Amendt, Campobasso, C.P., Goff, M.L. and Grassberger, M. *Current Concepts in Forensic*
 434 *Entomology*', *Springer*, pp. 179–204.

435 Espelie, K.E., Berisford, C.W. and Dahlsten, D.L. (1990) 'Cuticular hydrocarbons of
 436 geographically isolated populations of *Rhopalicus pulchripennis* (Hymenoptera:
 437 Pteromalidae): Evidence for two species', *Comparative Biochemistry and Physiology - Part B:*
 438 *Biochemistry*, 96(2), pp. 305–308.

439 Everaerts, C., Farine, J.P. and Brossut, R. (1997) 'Changes of species specific cuticular
 440 hydrocarbon profiles in the cockroaches *Nauphoeta cinerea* and *Leucophaea maderae* reared
 441 in heterospecific groups', *Entomologia Experimentalis et Applicata*, 85(2), pp. 145–150.

442 Gennard, D.E. (2007) 'Forensic entomology: An introduction', *John Wiley & Sons*.

443 Gibbs, A., Mousseau, T.A. and Crowe, J.H. (1991) 'Genetic and acclimatory variation in
 444 biophysical properties of insect cuticle lipids.', *Proceedings of the National Academy of*
 445 *Sciences*, 88(16), pp. 7257–7260.

446 Gibbs, A.G. (1998a) 'Water-proofing properties of cuticular lipids', *American Zoologist*, 38(3),
 447 pp. 471–482.

448 Gibbs, A.G. (1998b) 'The role of lipid physical properties in lipid barriers', *American Zoologist*,
 449 38(2), pp. 268–279.

450 Hadley, N.F. (1978) 'Cuticular permeability of desert Tenebrionid beetles: Correlations with
 451 epicuticular hydrocarbon composition', *Insect Biochemistry*, 8(1), pp. 17–22.

452 Howard, R.W. and Blomquist, G.J. (2005) 'Ecological, behavioral, and biochemical aspects of
 453 insect hydrocarbons', *Annual Review of Entomology*, 50, pp. 371–393.

454 Howard, R.W., Howard, C.D. and Colquhoun, S. (1995) 'Ontogenetic and environmentally
 455 induced changes in cuticular hydrocarbons of *Oryzaephilus surinamensis* (Coleoptera:
 456 Cucujidae)', *Annals of the Entomological Society of America*, 88(4), pp. 485–495.

457 Hwang, C.C. and Turner, B.D. (2009) 'Small-scaled geographical variation in life-history traits
 458 of the blowfly *Calliphora vicina* between rural and urban populations', *Entomologia*
 459 *Experimentalis et Applicata*, 132(3), pp. 218–224.

460 Ingleby, F.C., Hosken, D.J., Flowers, K., Hawkes, M.F., Lane, S.M., Rapkin, J., Dworkin, I.
 461 and Hunt, J. (2013) 'Genotype-by-environment interactions for cuticular hydrocarbon
 462 expression in *Drosophila simulans*', *Journal of Evolutionary Biology*, 26(1), pp. 94–107.

463 Kamhawi, S., Molyneux, D.H., Killick-Kendrick, R., Milligan, P.J.M., Phillips, A., Wilkes, T.J.
 464 and Killick-Kendrick, M. (1987) 'Two populations of *Phlebotomus ariasi* in the Cévennes

465 focus of leishmaniasis in the south of France revealed by analysis of cuticular hydrocarbons’,
 466 *Medical and Veterinary Entomology*, 1(1), pp. 97–102.

467 Kranz, W., Carroll, C., Dixon, D.A., Goodpaster, J. V. and Picard, C.J. (2017) ‘Factors affecting
 468 species identifications of blow fly pupae based upon chemical profiles and multivariate
 469 statistics’, *Insects*, 8(2).

470 Kruger, E.L. and Pappas, C.D. (1993) ‘Geographic variation of cuticular hydrocarbons among
 471 fourteen populations of *Aedes albopictus* (Diptera: Culicidae).’, *Journal of medical*
 472 *entomology*, 30(3), pp. 544–548.

473 Kruger, E.L., Pappas, C.D. and Howard, R.W. (1991) ‘Cuticular hydrocarbon geographic
 474 variation among seven North American populations of *Aedes albopictus* (Diptera: Culicidae).’,
 475 *Journal of medical entomology*, 28(6), pp. 859–864.

476 Limsopatham, K., Hall, M.J.R., Zehner, R., Zajac, B.K., Verhoff, M.A., Sontigun, N.,
 477 Sukontason, K., Sukontason, K.L. and Amendt, J. (2018) ‘A molecular, morphological, and
 478 physiological comparison of English and German populations of *Calliphora vicina* (Diptera:
 479 Calliphoridae)’, *PLoS ONE*, 13(12), pp. 1–22.

480 Markow, T.A. and Toolson, EC (2013) ‘Temperature effects on epicuticular hydrocarbons and
 481 sexual isolation in *Drosophila mojavensis*’, *Ecological and Evolutionary Genetics of*
 482 *Drosophila*, pp. 315–331.

483 Martin, S.J., Helanterä, H. and Drijfhout, F.P. (2008) ‘Colony-specific hydrocarbons identify
 484 nest mates in two species of Formica ant’, *Journal of Chemical Ecology*, 34(8), pp. 1072–1080.

485 Mazzanti, M., Alessandrini, F., Tagliabracci, A., Wells, J.D. and Campobasso, C.P. (2010)
 486 ‘DNA degradation and genetic analysis of empty puparia: Genetic identification limits in
 487 forensic entomology’, *Forensic Science International*, 195(1–3), pp. 99–102.

488 Menzel, F., Blaimer, B.B. and Schmitt, T. (2017) ‘How do cuticular hydrocarbons evolve?
 489 Physiological constraints and climatic and biotic selection pressures act on a complex functional
 490 trait’, *Proceedings of the Royal Society B: Biological Sciences*, 284(1850).

491 Michaud, J.P., Moreau, G. (2009) ‘Predicting the visitation of carcasses by carrion-related
 492 insects under different rates of degree-day accumulation’, *Forensic Science International*,
 493 185(1–3), pp. 78–83.

494 Michelutti, K.B., Soares, E.R.P., Sguarizi-Antonio, D., Piva, R.C., Suárez, Y.R., Cardoso,
 495 C.A.L. and Antonialli-Junior, W.F. (2018) ‘Influence of temperature on survival and cuticular
 496 chemical profile of social wasps’, *Journal of Thermal Biology*, 71, pp. 221–231.

497 Moore, H., Lutz, L., Bernhardt, V., Drijfhout, F.P., Cody, R.B. and Amendt, J. (2022) ‘Cuticular
 498 hydrocarbons for the identification and geographic assignment of empty puparia of forensically
 499 important flies’, *International Journal of Legal Medicine*, 1–10.

500 Moore, H. and Shemilt, S. (2018) ‘Cuticular hydrocarbon analysis in forensic entomology: A
 501 review’, *Archaeological and Environmental Forensic Science*, 1(2), pp. 127–138.

502 Moore, H.E., Adam, C.D. and Drijfhout, F.P. (2013) ‘Potential use of hydrocarbons for aging
 503 *Lucilia sericata* blowfly larvae to establish the postmortem interval’, *Journal of Forensic*
 504 *Sciences*, 58(2), pp. 404–412.

505 Moore, H.E., Butcher, J.B., Day, CR and Drijfhout, F.P. (2017a) ‘Adult fly age estimations
 506 using cuticular hydrocarbons and Artificial Neural Networks in forensically important
 507 Calliphoridae species’, *Forensic Science International*, 280, pp. 233–244.

508 Moore, H.E. and Drijfhout, F.P. (2015) ‘Surface hydrocarbons as min-PMI indicators. Fit for
 509 purpose?’, *Forensic Entomology: International Dimensions and Frontiers*, pp. 361–380.

510 Moore, H.E., Pechal, J.L., Benbow, M.E. and Drijfhout, F.P. (2017b) ‘The potential use of
 511 cuticular hydrocarbons and multivariate analysis to age empty puparial cases of *Calliphora*
 512 *vicina* and *Lucilia sericata*’, *Scientific Reports*, 7(1), pp. 1–11.

513 Noorman, N. and Den Otter, C.J. (2002) ‘Effects of relative humidity, temperature, and
 514 population density on production of cuticular hydrocarbons in housefly *Musca domestica* L.’,
 515 *Journal of Chemical Ecology*, 28(9), pp. 1819–1829.

516 Paula, M.C., Antonialli-Junior, W.F., Mendonça, A., Michelutti, K.B., Eulalio, ADMM,
 517 Cardoso, C.A.L., de Lima, T. and Von Zuben, C.J. (2017) ‘Chemotaxonomic profile and
 518 intraspecific variation in the blow fly of forensic interest *Chrysomya megacephala* (Diptera:
 519 Calliphoridae)’, *Journal of medical entomology*, 54(1), pp. 14–23.

520 Paula, M.C., Michelutti, K.B., Eulalio, ADMM, Mendonça, A., Cardoso, C.A.L., Andrade,
 521 L.H.C., Lima, S.M. and Antonialli-Junior, W.F. (2020) ‘New approach to application of mid-
 522 infrared photoacoustic spectroscopy in forensic analysis: Study with the necrophagous blow fly
 523 *Chrysomya megacephala* (Diptera: Calliphoridae)’, *Journal of Photochemistry and*
 524 *Photobiology B: Biology*, 209, p. 111934.

525 Paula, M.C., Michelutti, K.B., Eulalio, ADMM, Piva, R.C., Cardoso, C.A.L. and Antonialli-
 526 Junior, W.F. (2018) ‘New method for estimating the post-mortem interval using the chemical

527 composition of different generations of empty puparia: Indoor cases', *PLoS ONE*, 13(12), pp.
528 1–13.

529 Pechal, J.L., Moore, H., Drijfhout, F. and Benbow, M.E. (2014) 'Hydrocarbon profiles
530 throughout adult Calliphoridae aging: A promising tool for forensic entomology', *Forensic*
531 *Science International*, 245, pp. 65–71.

532 Rajpurohit, S., Hanus, R., Vrkoslav, V., Behrman, E.L., Bergland, A.O., Petrov, D., Cvačka, J.
533 and Schmidt, P.S. (2017) 'Adaptive dynamics of cuticular hydrocarbons in *Drosophila*',
534 *Journal of Evolutionary Biology*, 30(1), pp. 66–80.

535 Rosa-Freitas, M.G., Broomfield, G., Priestman, A., Milligan, P.J.M., Momen, H. and
536 Molyneux, D.H. (1992) 'Cuticular hydrocarbons, isoenzymes and behavior of three populations
537 of *Anopheles darlingi* from Brazil', *Journal of the American Mosquito Control Association*,
538 8(4), pp. 357–366.

539 Rouault, J., Cappy, P. and Jallon, JM (2001) 'Variations of male cuticular hydrocarbons with
540 geoclimatic variables: An adaptative mechanism in *Drosophila melanogaster*?', *Genetica*,
541 110(2), pp. 117–130.

542 Rouault, J.D., Marican, C., Wicker-Thomas, C. and Jallon, JM (2004) 'Relations between
543 cuticular hydrocarbon (HC) polymorphism, resistance against desiccation and breeding
544 temperature; a model for HC evolution in *D. melanogaster* and *D. simulans*', *Springer*, pp. 195–
545 212.

546 Roux, O., Gers, C. and Legal, L. (2008) 'Ontogenetic study of three Calliphoridae of forensic
547 importance through cuticular hydrocarbon analysis', *Medical and Veterinary Entomology*,
548 22(4), pp. 309–317.

549 Tarone, A.M., Picard, C.J., Spiegelman, C. and Foran, D.R. (2011) ‘Population and temperature
550 effects on *Lucilia sericata* (Diptera: Calliphoridae) body size and minimum development time’,
551 *Journal of Medical Entomology*, 48(5), pp. 1062–1068.

552 Toolson, EC (1984) ‘Interindividual variation in epicuticular hydrocarbon composition and
553 water loss rates of the cicada *Tibicen dealbatus* (Homoptera: Cicadidae)’, *Physiological*
554 *Zoology*, 57(5), pp. 550–56.

555 Toolson, EC (1982) ‘Effects of rearing temperature on cuticle permeability and epicuticular
556 lipid composition in *Drosophila pseudoobscura*’, *Journal of Experimental Zoology*, 222(3), pp.
557 249–253.

558 Toolson, EC and Hadley, N.F. (1979) ‘Seasonal effects on cuticular permeability and
559 epicuticular lipid composition in *Centruroides sculpturatus* Ewing 1928 (Scorpiones:
560 Buthidae)’, *Journal of comparative physiology*, 129(4), pp. 319–325.

561 Tregenza, T., Buckley, S.H., Pritchard, V.L. and Butlin, RK (2000) ‘Inter- and intrapopulation
562 effects of sex and age on epicuticular composition of meadow grasshopper, *Chorthippus*
563 *parallelus*’, *Journal of Chemical Ecology*, 26(1), pp. 257–278.

564 Zabala, J., Díaz, B. and Saloña-Bordas, M.I. (2014) ‘Seasonal blowfly distribution and
565 abundance in fragmented landscapes. Is it useful in forensic inference about where a corpse has
566 been decaying?’, *PLoS ONE*, 9(6)

567 Zhou, A., Tian, P., Li, Z., Li, X., Tan, X., Zhang, Z., Qiu, L., He, H., Ding, W. and Li, Y. (2020)
568 ‘Genetic diversity and differentiation of populations of *Chlorops oryzae* (Diptera,
569 Chloropidae)’, *BioMed Central Ecology*, 20(1), pp. 1–14.




Fig. 1. Locations of collection sites for *C. vicina* populations.




Fig. 2: GC chromatograms showing CHC profiles of *C. vicina* adult flies collected from a) Turkey, b) Germany, c) the United Kingdom, respectively. The marked areas show the population-specific key regions within the CHC profiles.




Fig. 3. Plot showing the total percentage concentrations of CHC classes detected in *C. vicina* adult flies.




Fig. 4. PCA plot showing PC3 vs PC2 vs PC4 for *C. vicina* adult flies clusters, using n-alkanes, alkenes, and methyl branched alkanes. Clusters were grouped by population.




Fig. 5. GC chromatograms showing CHC profiles of *C. vicina* empty puparia collected from a) Turkey, b) Germany, c) the United Kingdom, respectively.




Fig. 6. Plot showing the total percentage concentrations of CHCs detected in *C. vicina* empty puparia.




Fig. 7. PCA plot showing PC3 vs PC2 vs PC4 for *C. vicina* empty puparia clusters, using n-alkanes, alkenes, and methyl branched alkanes. Clusters were grouped by population.

Table 1. List of the compounds extracted from adult files of Turkish (TR), British (UK), and German (DE) populations of *C. vicina*, along with the total percentage peak area of each compound present and the percentage standard deviation

Peak number	Retention Time	Peak ID	<i>C. vicina</i> (UK)	<i>C. vicina</i> (DE)	<i>C. vicina</i> (TR)
1	17.779	C23	0.92 ± 0.34	0.68 ± 0.14	1.42 ± 1.05
2	18.587	7-MeC23	tr	tr	0.85 ± 0.62
3	19.693	C24	tr	tr	0.79 ± 0.59
4	21.041	3-MeC24	1.70 ± 1.15	ND	ND
5	21.802	C25	3.50 ± 1.58	2.81 ± 0.54	4.32 ± 1.65
6	22.555	11 + 13-MeC25	3.29 ± 1.24	1.57 ± 0.41	4.71 ± 2.97
7	21.151	9-MeC25	ND	ND	1.38 ± 0.40
8	22.691	7-MeC25	1.70 ± 2.09	0.68 ± 0.17	1.09 ± 0.26
9	23.079	5-MeC25	tr	tr	1.80 ± 1.05
10	23.163	4-MeC25	tr	tr	0.83 ± 0.50
11	23.314	3-MeC25	1.66 ± 0.64	1.28 ± 0.32	ND
12	23.846	C26	1.12 ± 0.47	1.33 ± 0.30	1.39 ± 0.31
13	24.087	13 + 15-MeC26	1.28 ± 0.33	1.24 ± 0.49	ND
14	24.583	11 + 13-MeC26	1.09 ± 0.23	0.91 ± 0.32	ND
15	24.690	8-MeC26	0.87 ± 0.57	0.58 ± 0.24	ND
16	23.876	5-MeC26	ND	3.18 ± 0.49	ND
17	25.296	4-MeC26	5.16 ± 3.13	0.64 ± 0.16	5.26 ± 3.13
18	25.813	3-MeC26	tr	0.71 ± 0.42	1.30 ± 1.31
19	24.169	Heptacosene	ND	ND	0.62 ± 0.18
20	26.145	C27	11.13 ± 6.11	10.99 ± 1.78	11.03 ± 2.76
21	26.440	13 + 15-MeC27	tr	0.72 ± 0.55	ND
22	26.890	11 + 13-MeC27	9.77 ± 3.10	9.94 ± 3.75	11.73 ± 2.54
23	27.024	7-MeC27	4.41 ± 7.72	2.59 ± 0.45	ND
24	27.375	6-MeC27	1.06 ± 0.99	ND	ND
25	27.421	5-MeC27	2.10 ± 0.96	2.47 ± 0.51	ND
26	27.682	4-MeC27	1.10 ± 0.76	2.32 ± 2.18	2.23 ± 0.89
27	27.844	3-MeC27	5.47 ± 1.89	6.22 ± 1.42	1.38 ± 0.59
28	28.138	C28	1.54 ± 0.73	2.07 ± 1.30	2.63 ± 0.76
29	28.363	13 + 15-MeC28	2.23 ± 0.67	3.62 ± 0.77	ND
30	28.708	11 + 13-MeC28	2.09 ± 0.94	2.10 ± 0.43	ND
31	28.799	8-MeC28	1.56 ± 1.03	1.52 ± 0.44	ND
32	28.886	6-MeC28	1.13 ± 0.62	1.50 ± 1.64	ND
33	29.171	4-MeC28	5.61 ± 3.96	4.90 ± 1.14	5.00 ± 1.57
34	29.305	3-MeC28	1.98 ± 0.95	3.06 ± 2.38	3.89 ± 2.65
35	29.590	C29	6.51 ± 3.42	6.20 ± 2.16	9.06 ± 2.56
36	29.763	13 + 15-MeC29	1.40 ± 0.73	2.31 ± 2.54	ND
37	29.940	11 + 13-MeC29	6.50 ± 3.25	7.71 ± 1.11	2.77 ± 0.87
38	30.001	7-MeC29	1.37 ± 0.24	1.95 ± 0.31	8.36 ± 2.45
39	30.134	5-MeC29	tr	0.58 ± 0.30	ND
40	30.185	4-MeC29	0.92 ± 0.50	0.82 ± 0.27	1.12 ± 0.40
41	30.251	3-MeC29	1.82 ± 0.90	2.19 ± 0.55	3.99 ± 1.91
42	30.527	C30	0.70 ± 0.28	tr	0.82 ± 0.21
43	30.736	8-MeC30	1.15 ± 0.69	1.11 ± 0.44	2.33 ± 0.83
44	30.910	4-MeC30	0.97 ± 0.69	1.10 ± 0.48	3.46 ± 0.98
45	30.985	3-MeC30	0.66 ± 0.21	tr	ND
46	31.106	C31	tr	0.82 ± 0.40	1.59 ± 0.72
47	31.142	13 + 15-MeC31	tr	0.65 ± 0.22	ND
48	29.993	11 + 13-MeC31	ND	0.56 ± 0.19	ND
49	31.360	9 + 11-MeC31	0.95 ± 0.43	1.61 ± 0.55	1.60 ± 0.64
50	31.590	3-MeC31	0.53 ± 0.20	1.20 ± 0.46	1.25 ± 0.37

nd: Not Detected.

tr: Detected in trace amounts (<0.5%).

Table 2. List of the compounds extracted from empty puparia of Turkish (TR), British (UK), and German (DE) populations of *C. vicina*, along with the total percentage peak area of each compound present and the percentage standard deviation

Peak Number	Retention Time	Peak ID	<i>C. vicina</i> (UK)	<i>C. vicina</i> (DE)	<i>C. vicina</i> (TR)
1	15.708	3-MeC21	1.44 ± 1.11	ND	ND
2	17.751	C23	tr	0.87 ± 0.49	0.83 ± 0.09
3	19.675	C24	tr	ND	0.55 ± 0.07
4	21.716	C25	4.52 ± 0.87	31.15 ± 6.66	4.96 ± 0.42
5	23.224	3-MeC25	0.57 ± 0.11	0.51 ± 0.21	0.73 ± 0.13
6	23.788	C26	3.74 ± 1.34	1.70 ± 0.50	2.23 ± 0.16
7	25.127	3-MeC26	0.93 ± 0.35	0.57 ± 0.17	0.90 ± 0.21
8	26.024	C27	36.66 ± 8.55	23.44 ± 5.36	32.02 ± 2.73
9	26.625	11 + 13-MeC27	4.19 ± 1.01	1.97 ± 0.76	3.04 ± 0.68
10	26.805	7-MeC27	1.31 ± 0.55	0.70 ± 0.25	1.21 ± 0.23
11	27.261	5-MeC27	1.13 ± 0.25	0.61 ± 0.20	0.98 ± 0.21
12	27.494	4-MeC27	0.99 ± 0.19	0.53 ± 0.17	0.79 ± 0.30
13	27.669	3-MeC27	8.13 ± 1.45	5.69 ± 1.68	7.50 ± 1.36
14	28.020	C28	4.71 ± 1.04	2.99 ± 0.96	5.99 ± 0.82
15	28.226	13 + 15-MeC28	1.21 ± 0.46	1.06 ± 0.20	ND
16	28.603	11 + 13-MeC28	1.37 ± 0.22	ND	0.62 ± 0.12
17	28.701	7-MeC28	0.58 ± 0.15	ND	0.57 ± 0.15
18	29.043	5-MeC28	1.74 ± 0.36	ND	ND
19	29.113	4-MeC28	0.32 ± 0.13	0.77 ± 0.25	1.68 ± 0.22
20	29.218	3-MeC28	2.24 ± 0.62	tr	0.90 ± 0.19
21	29.412	Nonacosene	ND	4.70 ± 1.21	ND
22	29.507	C29	14.10 ± 3.44	9.03 ± 2.26	21.34 ± 2.85
23	29.703	13 + 15-MeC29	0.72 ± 0.15	ND	1.27 ± 0.21
24	29.832	11 + 13-MeC29	3.92 ± 0.84	4.96 ± 1.15	3.06 ± 0.30
25	29.915	7-MeC29	1.18 ± 0.26	0.87 ± 0.28	0.94 ± 0.13
26	30.140	4-MeC29	0.83 ± 0.68	tr	ND
27	30.204	3-MeC29	1.64 ± 0.46	1.28 ± 0.37	2.38 ± 0.67
28	30.412	C30	0.36 ± 0.12	tr	1.66 ± 0.26
29	30.893	3-MeC30	ND	1.23 ± 3.08	ND
30	31.165	C31	0.96 ± 0.41	1.78 ± 0.59	3.82 ± 0.76
31	31.188	11 + 13-MeC31	ND	0.53 ± 0.17	ND
32	31.589	3-MeC31	ND	1.04 ± 0.85	ND
33	32.301	3-MeC32	ND	0.95 ± 0.38	ND

nd: Not Detected.

tr: Detected in trace amounts (<0.5%).



DSPACE

<https://dspace.org/>

Geographical variation of cuticular hydrocarbon profiles of adult flies and empty puparia amongst three populations of *Calliphora vicina* (Diptera: Calliphoridae)

Kula, Canan; Amendt, Jens; Drijfhout, Falko P.; Moore, Hannah E.

2023-01

Oxford University Press (OUP)

<https://doi.org/10.1093/jme/tjac167>

Kula C, Amendt J, Drijfhout FP, Moore HE. (2023) Geographical variation of cuticular hydrocarbon profiles of adult flies and empty puparia amongst three populations of *Calliphora vicina* (Diptera: Calliphoridae). *Journal of Medical Entomology*, Volume 60, Issue 1, January 2023, pp. 14–23