

**Effect of deficit irrigation and methyl jasmonate application on the
composition of strawberry (*Fragaria x ananassa*) fruit and leaves**

Jordi Giné-Bordonaba¹ and Leon A. Terry*

¹Plant Science Laboratory, Cranfield University, Bedfordshire, MK43 0AL, UK

²*Present address:* Institute for Food and Agricultural Research and Technology (IRTA),
Postharvest Programme, Parc Científic i Tecnològic Agroalimentari de Lleida, Parc de
Gardeny, Lleida, 25003, Spain.

* Corresponding author. Tel.: +44-7500 766490; Fax: +44-1234 758380.

E-mail address: l.a.terry@cranfield.ac.uk (L.A. Terry).

Abstract

Drought stress is among the most severe environmental risks threatening strawberry production. In the present study, the effect of deficit irrigation (DI; 50 mL/day) and/or elicitation with methyl jasmonate (MeJA; 0.1 mM) on the composition of secondary fruit and leaves from three strawberry pre-commercial cultivars (253/29, 279/4 and 279/5) was investigated and compared to plants kept at or near field capacity (200 mL/day). For certain cultivars (253/29), DI applied at green stage of fruit development resulted in a considerable reduction in berry size (1.7-fold). In other cultivars (279/4 and 279/5), fruit size was comparable in DI-treated and fully irrigated plants. Changes in the major sugars and organic acids of strawberry leaves and fruit were cultivar and organ dependent and were associated to an osmotic adjustment strategy within the plant to counteract the effects of drought. Overall, elicitation with MeJA had a minimal effect on plant growth and morphological traits. Nevertheless, MeJA increased fructose content of DI-treated leaves and palliated the differences in glucose content of fruit from different water treatments. The most pronounced effect of MeJA was related to an enhanced synthesis and accumulation of pelargonidin-3-glucoside (nearly 2-fold) in red-ripe fruit from cultivar 279/5.

Keywords: anthocyanins, berry size, drought, organic acids, sugars.

1. Introduction

Berries have long been recognised to play an important role in human nutrition, providing health-benefits against a wide range of diseases, mainly due to their elevated content in certain bioactives including ascorbate, anthocyanins, phenolic acids, carotenoids, etc. (Giné Bordonaba and Terry, 2011a; Manganaris et al., 2013). Most bioactive compounds within the plants are secondary metabolites whose synthesis can be triggered in response to biotic and abiotic stresses, such as UV radiation, drought, wounding as well as infections (Terry and Joyce, 2004; Terry et al., 2007a; Jahangir et al., 2009). In the particular case of strawberries, several studies have demonstrated the effect that certain preharvest treatments or cultivation practices have on strawberry fruit biochemistry (Terry et al., 2007a; Keutgen and Pawelzik, 2008; Crespo et al., 2010; Giné Bordonaba and Terry, 2010), including the effect on the concentration of certain taste- and health-related compounds. For instance, earlier works demonstrated that besides the positive environmental effects (i.e. water savings), deficit irrigation (DI) in strawberry plants resulted in berries with higher concentrations of anthocyanins and antioxidant capacity (Terry et al., 2007a) as well as other markers of strawberry fruit quality. Later studies revealed that such effects were cultivar dependent (Giné Bordonaba and Terry, 2010). Nonetheless, DI applied to strawberry plants has been linked with a significant reduction in fruit size and yield (Blatt, 1984; Serrano et al., 1992; Krüger et al., 1999; Liu et al., 2007; Terry et al., 2007a) which also seems to be cultivar dependent (Giné Bordonaba and Terry, 2010).

Jasmonic acid (JA) and its methyl ester methyl jasmonate (MeJA) are naturally occurring plant hormones which have been shown to regulate a wide range of physiological and biological processes (Cheong and Choi, 2003; Rohwer and Erwin, 2008), including responses to drought stress. Given the capacity of MeJA to act as an elicitor and considering that this

compound is already classified as Generally Recognise As Safe (GRAS) substance by the U.S. Food and Drug Administration (Wang et al., 2009) it may have a potential for enhancing the synthesis of bioactive compounds (Perez-Balibrea et al., 2011) and increase fruit quality whilst palliating the negative effects of DI. In this context, Wang (1999) reported that preharvest application of MeJA (0.01-0.1 mM) resulted in changes in plant metabolism that rendered strawberry leaves to better withstand in vitro water stress. Preharvest application of MeJA seems to alter stomatal opening in strawberry (Wang, 1999) and other crops (Horton, 1991) resulting in better transpiration control and hence potentially improving water stress tolerance. In soybean, MeJA has been shown to ameliorate the damaging effects of drought stress by modifying endogenous phytohormones and polyamines (Hassanein et al., 2009). Moreover, both preharvest and postharvest application of MeJA has been associated with greater antioxidant capacity in Chinese red bayberry (Wang et al., 2009) and strawberry (Ayala-Zavala et al., 2005) as well as enhanced anthocyanin synthesis in apples (Rudell and Mattheis, 2008) or other berries (Wang, 2003; Wang et al., 2008).

This study was conducted to further understand the response mechanisms of strawberry plants to water stress conditions and to elucidate whether or not elicitation with MeJA may be a suitable alternative to minimise the negative effects of DI on berry weight of three different pre-commercial strawberry cultivars whilst maintaining/enhancing the taste- and health-related composition of the fruit. Special attention was given to quantifying sugar and organic acids in both fruit and leaves as major respiratory substrates, and anthocyanins and ascorbic acid concentrations in fruit.

2. Materials and methods

2.1 Plant materials and experimental design

Three different maiden year cold-stored strawberry pre-commercial cultivars (*viz.* 253/29, 279/4 and 279/5) were supplied by Redeva (Surrey, UK) and grown in a glasshouse (April to July) in 1 L capacity pots containing commercial standard compost. Cultivars were selected by Redeva breeders to assess their potential adaptation to dry climates and drought. A completely randomised design was adopted considering cultivar, water treatments (50 or 200 ml day⁻¹) and MeJA treatments (none or 0.1 mM) as the principal sources of variation. Prior to commencing water treatments plants were kept at or near field capacity (*ca.* 0.7 m³ of water per m³ of soil; conductivity *ca.* 850 mV) for approximately three weeks following the methodology described in earlier works (Terry et al., 2007a; Giné Bordonaba and Terry, 2010). Water treatments started once the majority of primary fruit from the primary truss were at green I stage of development (prior to the second fruit expansion growth phase; Terry et al., 2004). Then, plants were irrigated daily (*ca.* 09:00 h) with either 50 or 200 ml day⁻¹ over an eight-week period. Methyl jasmonate (Sigma, Dorset, UK) treatments at 0.1 mM + 0.05% Tween-20 (Wang, 1999) were applied as a foliar spray to incipient runoff every 72h. MeJA treatments started when the majority of the primary fruit from the primary truss were at white stage of development. Similarly, control plants were sprayed with a 0.05% Tween-20 solution.

2.2 Soil moisture content and environmental monitoring

Soil moisture content, recorded as the conductivity from the growing media (mV), was measured daily (*ca.* 16.00h) by time-domain-reflectometry (TDR) using a Thetaprobe (ThetaKit type TK3, Delta-T devices, Cambs., UK). Hourly temperatures within the

glasshouse were recorded by means of a Tiny Tag Ultra 2 data logger (Gemini Data Logger, Sussex, UK), shielded from solar radiation. Mean temperature inside the glasshouse through the growing period was 21 °C.

2.3 Fruit, leaf and runner sampling

From each plant, all fruit from the primary truss were harvested at red stage. Four fully expanded leaves of similar size and age per plant were excised towards the end of the trial (30 days after initiation of water treatments and when all experimental fruit had been harvested) and the length and surface area of the leave recorded. On the last day of the experiment, following leaf sampling, the length as well as the total runner density (g) was determined for each plant. After harvest or excision, objective colour of fruit and leaves was measured using a Minolta CR-400 colorimeter and a DP-400 data processor (Minolta Co. Ltd., Japan) with an 8 mm light-path aperture, respectively (Terry et al., 2007b). Berry and leave weight was measured and recorded and thereafter immediately snap-frozen in liquid nitrogen and stored briefly at -40°C before being freeze-dried in an Edwards Modulyo freeze drier (W. Sussex, UK) for 6 and 4 days at 0.015kPa, respectively. Lyophilized samples were subsequently ground in a pestle and mortar, weighed and returned to the freezer until use. All reagents were purchased from Sigma (Dorset, UK) unless otherwise stated.

2.4 Extraction and quantification of sugars and organic acids

Sugars from both freeze-dried berries and leaves were extracted using 62.5% (v/v) aqueous methanol as described elsewhere (Terry et al. 2007a). Sugar content was determined using an Agilent 1200 series HPLC binary pump system (Agilent, Berks., UK), equipped with an Agilent refractive index detector (RID) G1362A. Strawberry extracts (20 µL) were diluted

(1:10), and injected into a Rezex RCM monosaccharide Ca⁺ (8%) column of 300 mm x 7.8 mm diameter (Phenomenex, CA, USA; Part no. 00H-0130-K0) with a Carbo-Ca²⁺ guard column of 4 mm x 3 mm diameter (Phenomenex,; Part no. AJ0-4493). Column and oven temperature as well as the mobile phase conditions were those reported earlier (Giné Bordonaba and Terry, 2010). Extracts for organic acids determination were prepared as described elsewhere from both berry or leave freeze-dried samples (Giné Bordonaba and Terry, 2010). L-ascorbic, citric, and malic acid contents in extracts were detected at 210 nm using the same HPLC system as described above equipped with an Agilent DAD G1315B/G1365B photodiode array with multiple wavelength detector. The mobile phase (1.0 mL min⁻¹) was analytical grade degassed 0.2% (w/v) metaphosphoric acid in H₂O (Giné Bordonaba and Terry, 2009). The presence and abundance of individual sugars or organic acids were automatically calculated by comparing sample peak area to standards (0.025-2.5 mg mL⁻¹) using ChemStation Rev. B.02.01.

2.5 Antioxidant capacity of strawberry leaves

Antioxidant capacity from strawberry leaves was measured using the FRAP assay as described in earlier works (Terry et al., 2007a; Giné Bordonaba and Terry, 2012) with some modifications. A 50 µL aliquot of diluted sample extract (1:9; v/v) or Fe²⁺ (FeSO₄ · 7H₂O) standards (0 – 5.0 mM) was added to 3.6 mL of freshly prepared FRAP working solution (viz. 5 mL of 10 mM TPTZ (2,4,6-tripyridyl-2-triazine) in 40 mM HCl + 5 mL of 10 mM FeCl₃ in 50 mL of 300 mM acetate buffer). The reaction mixture was incubated at 37°C for 10 min and absorbance measured spectrophotometrically at 593 nm using a Camspec M501 UV/Vis spectrophotometer. Antioxidant capacity was expressed as the concentration of antioxidants having a ferric reducing ability (mmols Fe²⁺ g⁻¹ DW).

2.6 Analysis of individual anthocyanins

Individual anthocyanins were extracted using the methodology described in earlier works (Giné Bordonaba and Terry, 2011b) by mixing 150 mg of freeze-dried fruit sample with 3 ml of 70% (v/v) methanol and 0.5% (v/v) HCl in HPLC-grade water. The slurry obtained was held at 35 °C in a water bath with constant shaking for 1.5 h; mixing the samples every 15 min. Finally, the flocculate obtained was filtered through a 0.2 µm Millex-GV syringe driven filter unit (Millipore Corporation, MA) and the clear extract analyzed by HPLC coupled to a Diode Array Detector (DAD), using the same equipment as described for sugars and organic acids. The anthocyanin profile of strawberry fruit was determined according to the method described by Gine Bordonaba et al. (2011).

2.7 Data analysis

All statistical analysis were carried out using Genstat for Windows, Version 10 (VSN International Ltd., Herts., UK). Data were subjected to analysis of variance (ANOVA) tests based on a completely randomised design. Least significant difference values (LSD; $P = 0.05$) were calculated for mean separation using critical values of t for two-tailed tests.

3. Results and Discussion

3.1 Morphological and physiological changes in response to drought or elicitation

In agreement with earlier works (Terry et al., 2007a; Giné Bordonaba and Terry, 2009; Grant et al., 2010) soil water content differed between treatments but also between cultivars according to the water stress conditions (**Figure 1**). In all cultivars, deficit irrigation resulted, in average, in 2-fold lower water content in the growing media if compared to fully irrigated plants. Soil water content for DI-treated plants declined following similar water-soil dynamics to that previously reported (Liu et al., 2007; Terry et al., 2007a; Savić et al., 2008; Giné Bordonaba and Terry, 2010; Grant et al., 2010). This said, greater differences between normal (200 mL day⁻¹) or deficit (50 mL day⁻¹) irrigated plants were encountered for cultivar 253/29 (less drought-tolerant cultivar; **Figure 1**). Accompanying the observed changes in plant water uptake, changes in berry weight of secondary strawberry fruit, from the primary truss, in response to different irrigation conditions were cultivar depended and therefore in agreement with earlier findings from Giné Bordonaba and Terry (2010) and Grant et al. (2010, 2012). Whereas no significant changes were observed in fruit weight of cvs. 279/4 and 279/5 subjected or not to DI conditions, cv. 253/29 held under drought conditions produced nearly 1.7-fold smaller fruit (**Figure 2**). This result together with the observed soil water conductivity values suggest that both cvs. 279/4 and 279/5 were more drought-tolerant than cv. 253/29. In comparison with other studies (Liu et al., 2007; Terry et al., 2007a; Savić et al., 2008; Giné Bordonaba and Terry, 2010) the lesser reduction of berry weight in fruit from cvs. 279/4 and 279/5 grown under DI conditions may be related to not only the capacity of these cultivars to better withstand water stress conditions but also may be associated to initiating water stress at later fruit developmental stages (i.e. green stage instead of flower initiation as reported earlier (Terry et al., 2007a; Giné Bordonaba and Terry, 2010)).

205 In contrast to the clear effect of water treatments on soil water content, plants from different
206 cultivars responded differently to foliar treatment with MeJA. Whereas preharvest application
207 of MeJA did not have an effect on the amount of water extracted from the growing media in
208 cv. 253/29, 1.5-fold lower soil water content was observed in MeJA-treated plants from cv.
209 279/4, regardless of the irrigation treatment (**Figure 1**). In cultivar 279/5, MeJA increased
210 (1.6-fold) and reduced (1.2-fold) soil water content for DI-treated and non-water stressed
211 plants, respectively. Differences in the amount of water extracted by the plant from the
212 growing media may be related to the direct effect of MeJA on root growth (Staswick et al.,
213 1992), which seems to be cultivar dependent, as well as a direct effect on root hydraulic
214 conductivity (Savić et al., 2008). For instance, in *Arabidopsis thaliana*, primary root growth
215 was inhibited by 50% when seedlings were grown on medium containing MeJA (Staswick et
216 al., 1992). Accordingly, visual inspection of root development at the end of the trial for each
217 plant indicated a positive correlation between the capacity of the plant to extract water and
218 root growth (data not shown) but supported only partially the associations between MeJA and
219 root development found by others (Staswick et al., 1992; Maksymiec and Krupa, 2007).
220 Differences between this and early studies may also be the result of the different species being
221 investigated as well as different methodologies used in the application of MeJA. In addition,
222 in the present study, MeJA was applied at an advanced fruit phenological stage when most of
223 the root development has already occurred. Elicitation with MeJA had a minimal effect on
224 berry weight and was cultivar dependent. In fruit from cv. 279/4 treatment with MeJA
225 resulted in a slight reduction of fruit weight regardless of the water treatment whereas no
226 effect was noticed in fruit from the rest of cultivars (**Figure 2**).

227 Fruit dry matter as a proportion of fresh weight was not significantly affected by the
228 conditions imposed in this study (**Figure 2**). In contrast, earlier works have shown that DI

applied at earlier fruit developmental stages resulted in fruit with greater dry matter content (Serrano et al., 1992; Krüger et al., 1999; Liu et al., 2007; Terry et al., 2007a; Giné Bordonaba and Terry, 2010). Dry matter content from leaves was neither affected by water treatments and hence it may be plausible to speculate that the concentration effect reported in earlier works (Terry et al., 2007a; Giné Bordonaba and Terry, 2010) is likely to have been related to the limitation of water uptake from the plants rather than to an enhanced import of solutes into the fruit from other parts of the plant (i.e. leaves).

As expected, greater water supplied to the plant resulted in greater runner biomass in all except in plants from the less drought-tolerant cv. 253/29. Accordingly, it is well accepted that greater water supply to strawberry plants generally resulted in higher vegetative growth, including runners fresh mass (Grant et al., 2010 and 2012). No other morphological traits (leaves length, foliar density) except some differences in objective colour of leaves were altered by the different treatments applied (data not shown).

3.2 Leaf biochemical changes

Fructose content of leaves was significantly different between cultivars, water treatments and the interaction between water treatments and MeJA (**Table 1**). Excised leaves from cultivar 279/5 had the greatest fructose content (20.26 mg g⁻¹ FW) followed by cv. 279/4 (15.41 mg g⁻¹ FW) and cv. 253/29 (13.24 mg g⁻¹ FW) and regardless of the water treatments. Under DI conditions, leaves had on average, 1.2-fold greater fructose content than leaves from plants kept at or near field capacity. Similar trends were observed for glucose and hence these results suggest that hexose sugars accumulate in response to drought as also reported for tomato plants submitted to other abiotic stress (Khelil et al., 2007). This specific behaviour may represent a beneficial plant response by avoiding metabolic inhibitions while

concomitantly contributing to an osmotic adjustment (solute potential) within the plant (Balibrea et al., 2000). Elicitation with MeJA in combination with DI resulted in all cases in higher glucose content as compared to non-elicited plants (**Table 1**). Nor sucrose or the total sugar content was affected by the conditions imposed in this study rather than genotypic differences. AsA concentrations were significantly different between cultivars, irrigation regimes but were not affected by MeJA. Specifically, AsA was greater in leaves from plants grown under reduced irrigation ($1.53 \text{ mg g}^{-1} \text{ DW}$) as compared to plants kept at or near field capacity ($1.15 \text{ mg g}^{-1} \text{ DW}$). Greater AsA content in DI-treated leaves may be a plant defence strategy to scavenge hydroxyl radicals and detoxify the accumulation of H_2O_2 resulting from water stress. The greatest increase (2-fold) in AsA as a result of drought was observed in leaves from cv. 279/5. Contrasting results were found by Wang et al (1999) where leaves of drought-stressed strawberry plants had lower AsA content if compared to leaves from control plants. However, different cultivars and experimental conditions were tested in each study. Malic acid content in leaves was affected by the interaction between water treatments and genotypes. Except for cv. 279/5, where malic acid was lower (1.7-fold) in leaves from DI-treated plants, reduced water resulted in greater (1.15-fold in average) amounts of malic acid in leaves from cultivars 279/4 and 253/29. Increased malic acid content in leaves, as also observed for certain sugars, may result from lowered respiration and hence confirm the osmotic adjustment (solute potential) strategies within different plant tissues or organs to counteract the effects of drought. Citric acid was the only acid affected by the interaction between cultivar, water treatments and elicitation with MeJA (**Table 1**). The differential osmotic adjustments observed among cultivars warrants further research aiming to obtain new strawberry cultivars with increase resistance to drought.

In terms of the leaves antioxidant capacity (AC), cultivar was the main source of variation, with cultivar 253/29 having the greatest AC values (133.5 mg Fe²⁺ g⁻¹ DW) followed by cvs. 279/5 (131.8 mg Fe²⁺ g⁻¹ DW) and 279/4 (111.2 mg Fe²⁺ g⁻¹ DW), respectively. Whereas none of the treatments alone had a significant effect on leaf AC, the interaction between water treatment and MeJA was significant. In plants kept at or near field capacity, elicitation with MeJA resulted in fairly similar AC values, whereas greater values were observed in plants subjected to DI conditions (**Figure 3**). This result is of particular interest since MeJA may partly alleviate drought stress by increasing the capacity of the plant to scavenge free radicals and thus limiting the oxidative stress.

3.3 Fruit biochemical changes

High variability in the sugar concentration was observed between fruit from different cultivars but overall values were in the range of those reported by others (Terry et al., 2007a; Giné Bordonaba and Terry, 2009; Giné Bordonaba and Terry, 2010; Crespo et al., 2010). Fruit from 279/4 and 279/5 had *ca.* 1.2-fold greater glucose and fructose concentrations than that of 253/29. Greater fructose or glucose content in these cultivars may be related to greater activity of some sugar mobilising enzymes (Khelil et al., 2007) or may be just a characteristic trait associated to greater drought tolerance. Sucrose concentrations were greater in cultivar 279/5 (254.6 mg g⁻¹ DW) followed by 253/29 (240.4 mg g⁻¹ DW) and 279/4 (203.9 mg g⁻¹ DW). None of the treatments applied had a marked effect on fruit sugar concentration. A significant interaction between MeJA, irrigation treatments and cultivar was only observed for glucose concentrations whereby DI resulted in lower glucose content only for fruit from cv. 279/5. MeJA minimised the differences in glucose content of fruit receiving different amounts of water. In agreement with these results, the response of strawberry plants to DI conditions

applied at earlier development stages resulted in different sugar concentrations depending on the cultivar (Giné Bordonaba and Terry, 2010). In other fruit, it has already been demonstrated that DI, unless applied during the major sugar-accumulation period, does not alter final sugar content (Barry et al., 2004). Therefore, the greater extent of DI treatments on strawberry sugars as described in earlier works (Terry et al., 2007a; Giné Bordonaba and Terry, 2010) may either be genotypically regulated or associated to initiating drought conditions at earlier fruit developmental stages (**Table 2**). Future studies should clarify the specific effect of applying DI conditions at different phenological stages on the final fruit composition of a range of cultivars. Wang et al. (2008) found that preharvest application of MeJA resulted in higher soluble solid content and lower titratable acidity of three blackberry cultivars and similar findings were earlier reported by Wang and Zeng (2005) on raspberries. Despite sugar content not always being well correlated with soluble solids measurements in strawberry fruit (Giné Bordonaba and Terry, 2009), the results presented herein demonstrate the differential effect of preharvest application of MeJA combined with full or deficit irrigation on sugar concentrations of strawberry fruit (**Table 2**).

Fruit AsA content was not affected by the conditions imposed in this study except for some cultivar differences. In agreement, earlier works (Giné Bordonaba and Terry, 2010) found AsA to be either higher, lower or not affected in plants submitted to drought stress depending on the cultivar. Besides, high variability in the AsA content among different strawberry cultivars has been extensively described in the literature (Tulipani et al., 2008; Giné Bordonaba and Terry, 2009; Giné Bordonaba and Terry, 2010; Crespo et al., 2010). AsA concentration in fruit from cultivar 253/29 ($17.1 \text{ mg g}^{-1} \text{ DW}$) was 2.3- or 1.5-fold higher than that for fruit from cvs. 279/4 ($7.3 \text{ mg g}^{-1} \text{ DW}$) and 279/5 ($11.7 \text{ mg g}^{-1} \text{ DW}$), respectively. Elicitation with MeJA did not induce significant changes in fruit AsA concentrations and

hence is in agreement with studies conducted on other horticultural crops (Pérez-Balibrea et al., 2011). In contrast, citric acid was quite similar among cultivars (56 mg g⁻¹ DW) but significantly affected by deficit irrigation or MeJA. Plants subjected to drought stress resulted in either higher or lower citric acid content depending on the cultivar which corroborates the genotype-specific response to reduced water supply reported by Giné Bordonaba and Terry (2010). Preharvest foliar application of MeJA resulted in fruit with greater citric acid content (60.2 mg g⁻¹ DW) if compared to fruit from non-treated plants (51.1 mg g⁻¹ DW) whilst opposite results were observed by Wang and Zheng (2005) in raspberries. The concentration of malic acid was affected by differences in genotypes as well as the interactions between irrigation and cultivar or between the different preharvest treatments applied fruit (**Table 2**), yet no clear conclusions may be drawn.

Four different anthocyanins, namely cyanidin-3-glucoside (Cya-3-gluc), pelargonidin-3-glucoside (Pg-3-gluc) and two pg-derivatives, were identified in the present study according to their retention time, UV spectra and comparison with standards (Giné Bordonaba et al., 2011), although their concentration strongly depended on the cultivar and treatment applied. Fruit from cultivar 253/29 were characterised by the presence of these four pigments whereas fruit from cultivars 279/4 and 279/5 lacked the pg-derivative 2 (**Table 3**). Significant differences between cultivars were observed for each individual anthocyanin hence corroborating the strong influence of genetic background in determining anthocyanin content of strawberry fruit (Carbone et al., 2009; Crespo et al., 2010). In detail, fruit from cultivar 279/5 had *ca.* 2-fold greater pg-3-gluc concentrations than that of fruit from cultivars 279/4 and 253/29. DI had little effect on the anthocyanin concentrations of the different cultivars investigated. In contrast, earlier works (Terry et al., 2007) found anthocyanin concentrations to be greater in fruit from plants (cv. Elsanta) that received less water. In the above-

mentioned study deficit irrigation was applied at flower initiation and hence it may be possible to speculate that even though anthocyanin accumulation occurs during later developmental stages (Carbone et al., 2009) the enhanced synthesis of these pigments as a result of drought stress may be up-regulated or triggered at much earlier development stages (i.e. from flower initiation up to green-stage). In addition, results from the present study revealed the differential effect that water deficit irrigation may have on strawberry anthocyanins of different cultivars as also observed for other metabolites herein and elsewhere (Giné Bordonaba and Terry, 2010). Preharvest elicitation with MeJA has been shown to enhance anthocyanin concentrations in berries (Pérez et al., 1997; Wang and Zheng, 2005; Wang et al., 2009) and other fruit (Rudell et al., 2008). In a recent study using *Fragaria Chiloensis* ripened *in vitro*, MeJA (10 and 100 μ M) was shown to induce ripening through its involvement of anthocyanin accumulation, cell wall degradation and the biosynthesis of ethylene (Concha et al., 2013). In the present study, anthocyanin concentration of the fruit was also positively influenced by preharvest treatment with MeJA; this was especially noticeable in fruit from cv. 279/5 where the concentration of cya-3-gluc, Pg-3-gluc and Pg-derivatives was 1.8, 2.7 and 1.8-fold greater, respectively, in MeJA-treated plants if compared to control. The enhanced synthesis of anthocyanins following elicitation with MeJA has been related to the direct effect of this hormone on plant secondary metabolism and especially on the activation of the enzyme phenylalanine ammonia lyase (Gundlach et al., 1992; Concha et al., 2013).

CONCLUSIONS

Plants have developed different mechanisms to survive periods of drought. The results from this study suggest that osmotic adjustment involving changes in sugar and acid

metabolism is one mechanism which in turn depends not only on the cultivar but also on the plant organ (namely leaf and/or fruit). Accordingly, this is the first report describing changes in the major sugar and organic acids in both fruit and leaves of strawberry plants subjected to deficit irrigation. Elicitation with MeJA, under the conditions described in this study, had a minimal effect on plant growth and some morphological traits (fruit weight, runners biomass, dry matter of leaves and fruit, etc.) which in turn was cultivar dependent. Nonetheless, MeJA resulted in increased fructose content of DI-treated leaves and minimised the differences in glucose content between fruit from different water treatments. Under the conditions investigated, MeJA did not alter fruit ripening (sugar and acid accumulation) but had a pronounced effect on the plant secondary metabolism by increasing the synthesis and accumulation of the main strawberry anthocyanin (Pg-3-gluc) in cultivar 279/5. In less drought-tolerant cultivars (namely 253/29), the negative effects of DI on fruit size were not mitigated and anthocyanin accumulation was not enhanced by elicitation with MeJA. Further studies are encouraged to investigate greater doses of this natural hormone or more continued applications aiming to further understand the relationship between drought resistance and MeJA in strawberry plants.

AKNOWLEDGEMENTS

The authors thank the Horticultural Development Company (CP 43) and GlaxoSmithKline for funding. Redeva, formerly the Summer Fruit Company, Total Berry (UK) is gratefully acknowledge for supplying the plants.

392 **REFERENCES**

- 393 Ayala-Zavala, J.F., Wang, S.Y., Wang, C.Y., Gonzalez-Aguilar, G.A., 2005. Methyl
394 jasmonate in conjunction with ethanol treatment increases antioxidant capacity,
395 volatile compounds and postharvest life of strawberry fruit. *Eur. Food Res. Technol.*
396 221, 731-738.
- 397 Barry, G.H., Castle, W.S., Davies, F.S., 2004. Rootstocks and plant water relations affect
398 sugar accumulation of citrus fruit via osmotic adjustment. *J. Amer. Soc. Hort. Sci.* 129,
399 881-889.
- 400 Blatt, C.R., 1984. Irrigation, mulch, and double row planting related to fruit size and yield of
401 Bounty strawberry. *HortScience* 19, 826-827.
- 402 Carbone, F., Preuss, A., De Vos, R.C.H., D'amico, E., Perrotta, G., Bovy, A.G., Martens, S.
403 and Rosati, C., 2009. Developmental, genetic and environmental factors affect the
404 expression of flavonoid genes, enzymes and metabolites in strawberry fruit. *Plant Cell*
405 *Environ.* 32, 1117-1131.
- 406 Cheong, J.J., Choi, Y.D., 2003. Methyl jasmonate as a vital substance in plants. *Trends Genet.*
407 19, 409-413.
- 408 Concha, C.M., Figueroa, N.E., Poblete, L.A., Oñate, F.A., Schwab, W. and Figueroa, C.R.
409 Methyl jasmonate treatment induces changes in fruit ripening by modifying the
410 expression of several ripening genes in *Fragaria chiloensis* fruit. *Plant Physiol.*
411 *Biochem.* 70, 433-444.
- 412 Crespo, P., Giné Bordonaba, J., Terry, L.A., Carlen, C., 2010. Characterisation of major taste
413 and health-related compounds of four strawberry genotypes grown at different Swiss
414 production sites. *Food Chem.* 122, 16-24.
- 415 Grant, O.M., Johnson, A.W., Davies, M.J., James, C.M., Simpson, D.W., 2010. Physiological
416 and morphological diversity of cultivated strawberry (*Fragaria x ananassa*) in response
417 to water deficit. *Environ. Exp. Bot.* 68, 264-272.
- 418 Grant, O.M., Davies, M.J., Johnson, A.W., Simpson, D.W., 2012. Physiological and growth
419 responses to water deficit in cultivated strawberry (*Fragaria x ananassa*) and in one of
420 its progenitors, *Fragaria chiloensis*. *Environ. Exp. Bot.* 83, 23-32.
- 421 Giné Bordonaba, J., Terry, L.A., 2009. Development of a glucose biosensor for rapid
422 assessment of strawberry quality: Relationship between biosensor response and fruit
423 composition. *J. Agric Food Chem.* 57. 8220-8226.

- 424 Giné Bordonaba, J., Terry, L.A., 2010. Manipulating the taste-related composition of
425 strawberry fruit (*Fragaria x ananassa*) from different cultivars using deficit irrigation.
426 *Food Chem.* 122, 1020-1026.
- 427 Giné Bordonaba, J., Terry, L.A., 2011a. Strawberry. In: Terry, L.A. (Ed.) Health-promoting
428 properties of fruit and vegetables. CABI, Wallingford, Oxfordshire, UK, pp. 291-320.
- 429 Giné Bordonaba, J., Crespo, P., Terry, L.A., 2011b. A new acetonitrile-free mobile phase for
430 HPLC-DAD determination of individual anthocyanins in blackcurrant and strawberry
431 fruit: a comparison and validation study. *Food Chem.* 129, 1265-1273.
- 432 Giné Bordonaba, J., Terry, L.A., 2012. Electrochemical behaviour of polyphenol rich fruit
433 juices using disposable screen-printed electrodes: Towards a rapid sensor for
434 antioxidant capacity and individual antioxidants. *Talanta*, 90, 38-45.
- 435 Gundlach, H., Müller, M.J., Kutchan, T.M., Zenk, M.H., 1992. Jasmonic acid is a signal
436 transducer in elicitor-induced plant cell cultures. *Proc. Natl. Acad. Sci. USA*, 89,
437 2389-2393.
- 438 Hassanein, R.A., Hassanein, A.A., El-din, A.B., Salama, M., Hashem, H.A., 2009. Role of
439 jasmonic acid and abscisic acid treatments in alleviating the adverse effects of drought
440 stress and regulating trypsin inhibitor production in soybean plant. *Aust. J. Bas. Appl.*
441 *Sci.* 3, 904-919.
- 442
- 443 Horton, R.F., 1991. Methyl jasmonate and transpiration in barley. *Plant Physiol.* 96, 1376-
444 1378.
- 445
- 446 Jahangir, M., Abdel-Farid, I. B., Kim, H. K., Choi, Y. H., Verpoorte, R., 2009. Healthy and
447 unhealthy plants: The effect of stress on the metabolism of brassicaceae. *Environ.*
448 *Exp. Bot.* 67, 23-33.
- 449 Keutgen, A. J., Pawelzik, E., 2008. Quality and nutritional value of strawberry fruit under
450 long term salt stress. *Food Chem.* 107, 1413-1420.
- 451 Krüger, E., Schmidt, G. and Brückner, U., 1999. Scheduling strawberry irrigation based upon
452 tensiometer measurement and a climatic water balance model. *Sci. Hortic.* 81(4), 409-
453 424.
- 454 Liu, F., Savić, S., Jensen, C. R., Shahnazari, A., Jacobsen, S. E., Stiki, R., Andersen, M.N.,
455 2007. Water relations and yield of lysimeter-grown strawberries under limited
456 irrigation. *Sci. Hortic.* 111, 128-132.

458 Manganaris, G.A., Goulas, V., Vicente, A.R. and Terry, L.A. 2013. Berry antioxidants: small
459 fruits providing large benefits. *J Sci. Food Agric.* 94, 825-833.

460 Maksymiec, W., Krupa, Z., 2007. Effects of methyl jasmonate and excess copper on root and
461 leaf growth. *Biol. Plantarum.* 51, 322-326.

462 Pérez, A.G., Sanz, C., Olías, R., Olías, J.M., 1997. Effect of methyl jasmonate on in vitro
463 strawberry ripening. *J. Agric. Food Chem.* 45, 3733-3737.

464 Pérez-Balibrea, S., Moreno, D.A., García-Viguera, C., 2011. Improving the phytochemical
465 composition of broccoli sprouts by elicitation. *Food Chem.* 129, 35-44.

466 Ravnikar, M., Vilhar, B., Gogala, N., 1992. Stimulatory effects of jasmonic acid in potato
467 stem node and protoplast culture. *J. Plant Growth Regul.* 11, 29-33.

468 Rohwer, C.L., Erwin, J.E., 2008. Horticultural applications of jasmonates: a review. *J. Hortic.*
469 *Sci. Biotech.* 83, 283-304.

470 Rudell, D.R., Mattheis, J. P., 2008. Synergism exists between ethylene and methyl jasmonate
471 in artificial light-induced pigment enhancement of 'Fuji' apple fruit peel. *Postharv.*
472 *Biol. Technol.* 47, 136-140.

473 Savić, S., Stiki, R., Radović, B.V., Bogievi, B., Jovanović, Z., Šukalović, V.H., 2008.
474 Comparative effects of regulated deficit irrigation (RDI) and partial root-zone drying
475 (PRD) on growth and cell wall peroxidase activity in tomato fruit. *Sci. Hortic.* 117, 15-
476 20.

477 Serrano, L., Carbonell, X., Savé, R., Marfà, O., Peñuelas, J., 1992. Effects of irrigation
478 regimes on the yield and water use of strawberry. *Irrig. Sci.*, 13, 45-48.

479 Staswick, P. E., Su, W., Howell, S. H., 1992. Methyl jasmonate inhibition of root growth and
480 induction of a leaf protein are decreased in an arabidopsis thaliana mutant. *Proc. Natl.*
481 *Acad. Sci. USA*, 89, 6837-6840.

482 Terry, L.A. and Joyce, D.C. 2004. Influence of growing condition and associated variable
483 efficacy of acibenzolar in suppression of *Botrytis cinerea* on strawberry fruit. *Adv.*
484 *Straw. Res.* 23, 11-19

485 Terry, L.A., Joyce, D.C., Adikaram, N.K.B., Khambay, B.P.S., 2004. Preformed antifungal
486 compounds in strawberry fruit and flower tissues. *Postharv. Biol. Technol.* 31, 201-
487 212.

- 488 Terry, L.A., Chope, G.A., Giné Bordonaba, J., 2007a. Effect of water deficit irrigation and
489 inoculation with *Botrytis cinerea* on strawberry (*Fragaria x ananassa*) fruit quality. *J.*
490 *Agric. Food Chem.* 55, 10812-10819.
- 491 Terry, L.A., Ilkenhans, T., Poulston, S., Rowsell, L., Smith, A.W.J., 2007b. Development of
492 new palladium-promoted ethylene scavenger. *Postharv. Biol. Technol.* 45, 214-220.
- 493 Tulipani, S., Mezzetti, B., Capocasa, F., Bompadre, S., Beekwilder, J., De Vos, C.H.R.,
494 Capanoglu, E., Bovy, A. Battino, M., 2008. Antioxidants, phenolic compounds, and
495 nutritional quality of different strawberry genotypes. *J. Agric. Food Chem.* 56, 696-
496 704.
- 497 Wang, S. Y., 1999. Methyl jasmonate reduces water stress in strawberry. *J. Plant Growth*
498 *Regul.* 18, 127-134.
- 499 Wang, C.Y., 2003. Maintaining postharvest quality of raspberries with natural volatile
500 compounds. *Int. J. Food Sci. Technol.* 38, 869-875.
- 501 Wang, S.Y., Zheng, W., 2005. Preharvest application of methyl jasmonate increase fruit
502 quality and antioxidant capacity in raspberries. *Int. J. Food Sci. Technol.* 40, 187-195.
- 503 Wang, S.Y., Bowman, L., Ding, M., 2008. Methyl jasmonate enhances antioxidant capacity
504 and flavonoid content in blackberries (*Rubus sp.*) and promotes antiproliferation of
505 human cancer cells. *Food Chem.* 107, 1261-1269.
- 506 Wang, K., Jin, P., Cao, S., Shang, H., Yang, Z., Zheng, Y., 2009. Methyl jasmonate reduces
507 decay and enhances antioxidant capacity in Chinese bayberries. *J. Agric. Food Chem.*
508 57, 5809-5815.
- 509 Wolfe, K.L., Kang, X., He, X., Dong, M., Zhang, Q., Liu, R.H., 2008. Cellular antioxidant
510 activity of common fruit. *J. Agric Food Chem.* 56, 8418-8426.

511

512

513

514

515

516

517

518

519

520

521

522 **Table captions:**

523 **Table 1:** Sugar concentration (mg g^{-1} DW) in strawberry secondary fruit and leaves of three
524 different cultivars in response to drought and/or elicitation with 0 (-MJ) or 0.1 mM MeJA
525 (+MJ).

526 **Table 2:** Organic acid concentration (mg g^{-1} DW) in strawberry secondary fruit and leaves of
527 three different cultivars in response to drought and/or elicitation with 0 (-MJ) or 0.1 mM
528 MeJA (+MJ).

529 **Table 3:** Concentration of individual anthocyanins (cyanidin-3-glucoside (Cya-3-gluc),
530 pelargonidin-3-glucoside (Pg-3-gluc) and pelargonidin derivatives (Pg-deriv 1 and 2) in
531 strawberry secondary fruit from three different cultivars in response to drought and/or
532 elicitation with 0 (-MJ) or 0.1 mM MeJA (+MJ).

533

534 **Figure captions:**

535 **Figure 1:** Average water conductivity of the growing media in plants from three strawberry
536 pre-commercial cultivars (A: 253/29; B: 279/4; C: 279/5) grown under deficit irrigation (DI;
537 50 mL day⁻¹) or full irrigation (CT; 200 mL day⁻¹) and treated with 0 or 0.1 mM foliar
538 application of MeJA. (D) Dynamic changes in soil water content (expressed as soil
539 conductivity) of the growing media grown under deficit irrigation (50 mL day⁻¹; empty

symbols) or full irrigation (200 mL day⁻¹; full symbols) and treated with 0 (round) or 0.1 mM (triangle) foliar application of MeJA. Error bar depicts LSD values (p<0.05) for the interaction between water treatment*MeJA (A, B and C) or water treatment*MeJA*day (D).

Figure 2: Runners biomass (g plant⁻¹) and weight characteristics of fruit and leaves from three strawberry pre-commercial cultivars grown under deficit irrigation (DI; 50 mL day⁻¹) or full irrigation (CI; 200 mL day⁻¹) and treated with 0 (MJ-) or 0.1 mM foliar application of MeJA(+MJ). Error bars depict standard deviation for n=3.

Figure 3: Antioxidant capacity (mg Fe²⁺ g⁻¹ DW) of strawberry leaves from three strawberry pre-commercial cultivars grown under deficit irrigation (DI; 50 mL day⁻¹) or full irrigation (CI; 200 mL day⁻¹) and treated with 0 (MJ-) or 0.1 mM foliar application of MeJA (+MJ). Error bars depict standard deviation for n=3.

Table 1: Sugar and organic acid concentration (mg g⁻¹ DW) in leaves of strawberry plants from three different pre-commercial cultivars in response to drought and/or elicitation with 0 (-MJ) or 0.1 mM MeJA (+MJ).

Cv	Irrigation	Fructose		Glucose		Sucrose		Ascorbic		Citric		Malic	
		-MJ	+MJ	-MJ	+MJ	-MJ	+MJ	-MJ	+MJ	-MJ	+MJ	-MJ	+MJ
253/29	50	35.5	42.3	41.3	48.7	59.3	64.3	1.7	1.5	11.3	10.5	27.2	25.5
	200	36.9	33	51.9	41.2	65.1	48.5	1.3	1.1	8.4	8.0	24.6	15.8
279/4	50	45.1	55.9	51.8	65.2	82.4	75.9	1.1	1.4	10.9	7.6	28.0	32.9
	200	32.4	36.1	41.4	45.1	66.8	75.6	1.0	1.0	9.0	10.8	26.3	20.1
279/5	50	51.4	69	53.4	70.3	84.7	87.6	2.8	2.0	12.2	16.0	16.6	19.5
	200	52.8	50.3	56.3	56.4	82.3	87.2	1.2	1.3	16.4	12.4	28.9	27.1
LSD _{cv}		7.53		3.61		10.22		0.39		2.51		7.50	
LSD _{cv*irrigation}		10.64		10.56		14.46		0.55		3.56		10.61	
LSD _{cv*irrigation*MeJA}		15.05		15.00		20.45		0.78		5.04		14.00	

Table 2: Sugar and organic acid concentration (mg g⁻¹ DW) in strawberry secondary fruit of three different pre-commercial cultivars in response to drought and/or elicitation with 0 (-MJ) or 0.1 mM MeJA (+MJ).

Cv	Irrigation	Fructose		Glucose		Sucrose		Ascorbic		Citric		Malic	
		-MJ	+MJ	-MJ	+MJ	-MJ	+MJ	-MJ	+MJ	-MJ	+MJ	-MJ	+MJ
253/29	50	189.5	191.2	183.6	190.2	213.5	262.4	15.6	25.5	62.4	82.4	8.3	14.1
	200	178.9	191.3	174.4	185.3	264.3	221.4	16.6	10.5	56.6	44.7	13.1	10.5
279/4	50	209.2	201.3	213.9	194.8	210.7	193.1	5.4	5.0	42.7	50.1	6.1	6.6
	200	204.2	211.4	206.7	210.2	210	201.9	8.5	10.5	54.9	60.9	13.9	8.5
279/5	50	248	226.4	181.5	228.8	249.8	232.7	12.5	15.0	59.4	68.1	16.9	13.0
	200	216.5	209.4	218.9	204.4	284.3	251.6	5.5	13.9	34.8	54.9	13.6	11.6
LSD _{cv}		24.14		15.34		25.61		5.14		9.54		2.59	
LSD _{cv*irrigation}		30.13		21.78		36.22		7.26		13.49		3.67	
LSD _{cv*irrigation*MeJA}		38.27		30.68		51.22		10.27		19.07		5.19	

Table 3: Concentration ($\mu\text{g g}^{-1}$ DW) of individual anthocyanins (cyaniding-3-glucoside (Cya-3-gluc), pelargonidin-3-glucoside (Pg-3-gluc) and pelargonidin derivatives (Pg-deriv 1 and 2) in strawberry secondary fruit from three different cultivars in response to drought and/or elicitation with 0 or 0.1M MeJA.

Cv	Irrigation	Cya-3-gluc		Pg-3-gluc		Pg-deriv. 1		Pg-deriv. 2	
		-MJ	+MJ	-MJ	+MJ	-MJ	+MJ	-MJ	+MJ
253/29	50	44.2	33.5	1559.2	880.2	148.3	89.8	377.7	227.2
	200	41.4	33.3	776	874	68.1	94.8	166.6	237.2
279/4	50	20.1	30.8	764	587	47.5	42.6	-	-
	200	23.2	27.9	1252	1054	59.7	58.8	-	-
279/5	50	26	48.3	1174	3198	72.4	132.3	-	-
	200	22.9	31.5	890	2782	57.6	123.6	-	-
LSD _{cv}		5.11		356.27		15.49		-	
LSD _{cv*irrigation}		7.23		503.82		21.91		75.84	
LSD _{cv*irrigation*MeJA}		10.22		712.51		30.99		102.13	




Figure 1: Giné Bordonaba and Terry




Figure 2: Giné Bordonaba and Terry





Figure 3: Giné Bordonaba and Terry

RESEARCH HIGHLIGHTS

- The effects of deficit irrigation from green stage to ripe fruit on strawberry fruit size and quality are cultivar dependent.
- Changes in sugar and organic acids in response to drought were differently regulated in fruit than in leaves.
- Elicitation with MeJA had a minimal effect on plant growth.
- MeJA increased the concentration of pelargonidin-3-glucoside in ripe red fruit from one cultivar.

Graphical Abstract





DSPACE

<https://dspace.org/>

Effect of deficit irrigation and methyl jasmonate application on the composition of strawberry (*Fragaria x ananassa*) fruit and leaves

Giné-Bordonaba, Jordi; Terry, Leon A.

2015-12-31

Elsevier

<http://dx.doi.org/10.1016/j.scienta.2015.12.026>

Jordi Giné-Bordonaba, Leon A. Terry, Effect of deficit irrigation and methyl jasmonate application on the composition of strawberry (*Fragaria x ananassa*) fruit and leaves, *Scientia Horticulturae*, Volume 199, 16 February 2016, pp63-70